

# Tevelev degrees

Alessio Cela

ETH Zurich

## Introduction and main definitions

Let  $X$  be a smooth projective variety,  $g \geq 0$  a genus,  $n \geq 0$  and  $\beta \in H_2(X, \mathbb{Z})$  a curve class. Fix  $x_1, \dots, x_n \in X$  general points of  $X$ . We are interested in counting maps from  $C$  to  $X$  in class  $\beta$  and passing through  $x_1, \dots, x_n$ .

Assume  $2g - 2 + n > 0$ , so that the moduli stack  $\overline{\mathcal{M}}_{g,n}$  of stable curves is well-defined and let  $\overline{\mathcal{M}}_{g,n}(X, \beta)$  be the moduli stack of  $n$ -pointed genus  $g$  stable maps in class  $\beta$  to  $X$ .

There is a map

$$\begin{aligned} \bar{\tau} : \overline{\mathcal{M}}_{g,n}(X, \beta) &\longrightarrow \overline{\mathcal{M}}_{g,n} \times X^{\times n} \\ [f : (C, p_1, \dots, p_n) \rightarrow X] &\mapsto ((\bar{C}, \bar{p}_1, \dots, \bar{p}_n), (f(p_1), \dots, f(p_n))) \end{aligned}$$

recalling the stabilized domain curve and the image of the marked points under the morphism. One way to formulate our problem is by looking at the degree of  $\bar{\tau}$ .

Note that  $\overline{\mathcal{M}}_{g,n}(X, \beta)$  has virtual dimension equal to the dimension of  $\overline{\mathcal{M}}_{g,n} \times X^{\times n}$  if and only if

$$c_1(X) \cdot \beta = r(n + g - 1). \quad (1)$$

### Definition of the virtual count

Assume condition (1) is satisfied. Then the **virtual Tevelev degree**  $\text{vTev}_{g,n,\beta}^X \in \mathbb{Q}$  is defined by

$$\bar{\tau}_*([\overline{\mathcal{M}}_{g,n}(X, \beta)]^{\text{vir}}) = \text{vTev}_{g,n,\beta}^X [\overline{\mathcal{M}}_{g,n} \times X^{\times n}].$$

Here  $[\ ]^{\text{vir}}$  and  $[ ]$  denote the virtual and the usual fundamental classes.

One can also define the geometric count as follows. Let  $\mathcal{M}_{g,n} \subset \overline{\mathcal{M}}_{g,n}$  and  $\mathcal{M}_{g,n}(X, \beta) \subset \overline{\mathcal{M}}_{g,n}(X, \beta)$  be the loci where the curve  $C$  is smooth and let

$$\tau : \mathcal{M}_{g,n}(X, \beta) \rightarrow \mathcal{M}_{g,n} \times X^{\times n}$$

be the restriction of  $\bar{\tau}$ .

### Definition of the geometric count

Assume condition (1) is satisfied. Assume further that for the general point  $((C, p_1, \dots, p_n), (x_1, \dots, x_n)) \in \mathcal{M}_{g,n} \times X^{\times n}$  the fiber under  $\tau$  consists of finitely many reduced (necessarily non-stacky) points. Then we define the **Geometric Tevelev degrees**  $\text{Tev}_{g,n,\beta}^X \in \mathbb{Z}$  by

$$\text{Tev}_{g,n,\beta}^X = \#\text{general fiber of } \tau.$$

## Projective line

Using a slightly different point of view, Tevelev [10] computed some Geometric Tevelev degrees of  $\mathbb{P}^1$ . The full description of Geometric Tevelev of  $\mathbb{P}^1$  have been obtained in [5] via intersection theory on Hurwitz spaces, the case of  $\mathbb{P}^n$  is instead treated in [6] via limit linear series. Building on these two approaches, these counts are generalized in [4] for  $X = \mathbb{P}^1$  to the situation where the covers are constrained to have arbitrary ramification profiles. The following is [5, Theorem 6].

### Explicit formulas for $\mathbb{P}^1$

Let  $g \geq 0$ ,  $\ell \in \mathbb{Z}$ , and call

$$d[g, \ell] = g + 1 + \ell, \quad \text{and} \quad n[g, \ell] = g + 3 + 2\ell.$$

Assume  $n[g, \ell] \geq 3$  and  $d[g, \ell] \geq 1$ . Then we have:

$$\text{Tev}_{g,n[g,\ell],d[g,\ell]}^{\mathbb{P}^1} = 2^g - 2 \sum_{i=0}^{-\ell-2} \binom{g}{i} + (-\ell - 2) \binom{g}{-\ell - 1} + \ell \binom{g}{-\ell},$$

**Sketch of Proof** Let  $\overline{\mathcal{H}}_{g,d[g,\ell],n[g,\ell]}$  be the moduli stack of degree  $d[g, \ell]$  and  $n[g, \ell]$  marked admissible covers [8] and

$$\bar{\tau} : \overline{\mathcal{H}}_{g,d[g,\ell],n[g,\ell]} \rightarrow \overline{\mathcal{M}}_{g,n} \times \overline{\mathcal{M}}_{0,n[g,\ell]}$$

be the map recalling the marked domain curve (the ramification points are forgotten) and the marked target curve (the branch points are forgotten). The advantage of replacing  $\overline{\mathcal{M}}_{g,n}(X, d[\mathbb{P}^1])$  with  $\overline{\mathcal{H}}_{g,d,n}$  is that the boundary of the Hurwitz stack has a very nice stratification.

Up to a combinatorial factor, we want to find the degree of  $\bar{\tau}$  and we do this by computing the degree of the zero cycle


$$\bar{\tau}^*[(C, D)] \in \mathbb{A}_0(\overline{\mathcal{H}}_{g,d[g,\ell],n[g,\ell]}).$$

where the point

$$(C, D) \in \overline{\mathcal{M}}_{g,n[g,\ell]} \times \overline{\mathcal{M}}_{0,n[g,\ell]}$$

is chosen to have the following form:  $C$  is obtained by gluing at two points a smooth genus  $g-1$  curve containing  $n[g, \ell] - 1$  marked points and a smooth genus 0 curve containing 1 marked point,  $D$  is a smooth  $n[g, \ell]$ -pointed genus 0 curve.

The actual fiber  $\bar{\tau}^{-1}[(C, D)]$  will have excess dimension, so some care must be taken in the analysis. Fixed  $(C, D)$ , the Hurwitz cover can degenerate only in one of the following three ways:



**Explanation of the picture:** degrees of the map are written in green, the last marking is in blue and the first  $n[g, \ell]$  markings are in red.

From this one deduces the following recursion:

$$\text{Tev}_{g,n[g,\ell],d[g,\ell]} = \text{Tev}_{g-1,n[g-1,\ell],d[g-1,\ell]} + \text{Tev}_{g-1,n[g-1,\ell+1],d[g-1,\ell+1]}$$

reducing the problem to the genus 0 case. Finally the genus 0 case is treated by hand:

$$\text{Tev}_{0,n[0,\ell],d[0,\ell]} = 1 \text{ for all } \ell \geq 0.$$

## Application : Castelnuovo's classical count of $g_d^1$ 's

Let  $C$  be a general smooth genus  $g$  curve. Fix a degree  $d \geq 1$  and consider the Brill-Noether locus

$$G_d^1(C) = \{g_d^1\text{'s on } C\}$$

which is smooth of dimension  $\rho = g - 2(g - d + 1)$ . Assume  $\rho = 0$ . Then we can write  $g = -2\ell$  and  $d = g + \ell + 1$  for some  $\ell \in \mathbb{Z}$  and

$$G_d^1(C) = W_d^1(C) = \{L \in \text{Pic}^d(C) \mid h^0(C, L) \geq 2\} \subseteq \text{Pic}^d(C)$$

In his famous paper [2] Castelnuovo proved that

$$\deg([W_d^1(C)]) = \frac{1}{1 + |\ell|} \binom{2|\ell|}{|\ell|}$$

which agrees (after some algebraic manipulations) with  $\text{Tev}_{g,3,d}^{\mathbb{P}^1}$ .

## The quantum Euler class

Denote by  $(QH^*(X, \mathbb{Q}), \star)$  the small quantum cohomology ring of  $X$  (see [7] for an introduction) and let

$$H^*(X, \mathbb{Q}) \otimes H^*(X, \mathbb{Q}) \xrightarrow{\star} QH^*(X, \mathbb{Q}).$$

be the multiplication map.

### Definition of the Quantum Euler class

The **quantum Euler class**  $E$  of  $X$  is the image of the diagonal class  $[\Delta]$  under the multiplication map above (note that  $[\Delta]$  lives naturally in  $H^*(X, \mathbb{Q}) \otimes H^*(X, \mathbb{Q})$  via the Künneth isomorphism).

This class plays a central role in the computation of Virtual Tevelev degrees. Indeed, we have the following equality (see [1, Theorem 1.3]):

$$\text{vTev}_{g,n,\beta}^X = \text{Coeff}(\mathbf{P}^{\star n} \star \mathbf{E}^{\star g}, q^{\beta} \mathbf{P}) \quad (2)$$

where  $\mathbf{P}$  is the point class.

## Comparison between the virtual and the geometric count

Enumerativity results of Virtual Tevelev Degrees have been studied in [9]. To state their main result [9, Theorem 24] we require additional notation.

Assume  $X$  is a **Fano** variety of dimension  $r$ . Define  $s(X) > 0$  to be the smallest positive integer for which there exists an effective curve class  $\beta \in H_2(X, \mathbb{Z})$  such that

$$s(X) = c_1(X) \cdot \beta$$

and such that the evaluation map  $\text{ev}_1 : \overline{\mathcal{M}}_{0,1}(X, \beta) \rightarrow X$  is surjective.

Define  $t(X) > 0$  to be the smallest positive integer for which there exists an effective curve class  $\beta \in H_2(X, \mathbb{Z})$  such that

$$t(X) = c_1(X) \cdot \beta.$$

## Enumerativity

Fix a genus  $g \geq 0$ . Assume that:

- there exists  $k > 0$  such that for all  $\beta$  satisfying  $c_1(X) \cdot \beta > k$  we have  $c_1(X) \cdot \beta > (r - s(X))h^1(f^*T_X)$  for all  $[f] \in \mathcal{M}_g(X, \beta)$ ;
- $s(X) + t(X) \geq r + 1$ .

Then there exist  $d[g, X] > 0$  such that for all  $\beta$  such that  $c_1(X) \cdot \beta > d[g, X]$  and  $n = n[g, X, \beta] \geq 0$  such that Equation (1) is satisfied, the Geometric Tevelev degree  $\text{Tev}_{g,n,\beta}^X$  is well-defined and coincides with the Virtual Tevelev degree  $\text{vTev}_{g,n,\beta}^X$ .

**Simple Example** For  $X = \mathbb{P}^r$  we have

$$\text{vTev}_{g,n,dL}^{\mathbb{P}^r} = (r + 1)^g$$

where  $L$  is the class of a line (see [1, Example 2.2]). In particular, for  $r = 1$  and  $\ell \geq 0$  we see that

$$\text{vTev}_{g,n[g,\ell],d[g,\ell]}^{\mathbb{P}^1} = \text{Tev}_{g,n[g,\ell],d[g,\ell]}^{\mathbb{P}^1} = 2^g.$$

## Fano Hypersurfaces

Let  $X \subset \mathbb{P}^{r+1}$  be a smooth Fano hypersurface of dimension  $r \geq 3$  and degree  $m \geq 2$ . Note that  $X$  is Fano precisely when  $m \leq r + 1$ . Also, by Lefschetz Hyperplane theorem we have

$$H_2(X, \mathbb{Z}) = \mathbb{Z}L$$

where  $L$  is the class of a line in  $X$ . In particular

$$QH^*(X, \mathbb{Q}) = H^*(X, \mathbb{Q}) \otimes_{\mathbb{Q}} \mathbb{Q}[q]$$

as  $\mathbb{Q}[q]$ -module. Although the definition of  $E$  involves also the primitive cohomology of  $X$ , in [3, Theorem 5], we were able to obtain explicit simple formulas for  $E$ .

### The Quantum Euler class for Fano Hypersurfaces

The following equalities hold:

- if  $m \leq r$  then

$$E = m^{-1} \chi(X) H^{*r} + (r + 2 - m - \chi(X)) m^{m-1} q H^{*m-2},$$

- if  $m = r + 1$  then

$$E = m^{-1} \chi(X) H^{*r} + \sum_{j=1}^r m^{-1} (j - \chi(X)) \binom{r}{j-1} (m!)^{j-1} \left[ m^m - \frac{m!}{j} (r+1) \right] q^j H^{*r-j}.$$

Using this and Equation (2) it is also possible to obtain formulas for  $\text{vTev}_{g,n,dL}$  (in terms of  $\mathbf{P}$ ). In particular, for low degree hypersurfaces we have (see [1, Theorem 5.19] and [9, Theorem 11]):

### Explicit formulas for low degree Fano Hypersurfaces

If  $r > \max(2m - 4, 2)$  and  $X$  is not a quadric, then

$$\text{vTev}_{g,n,dL}^X = ((m-1)!)^n (r+2-m)^g m^{(d-n)m-g+1}$$

If in addition  $r > (m+1)(m-2)$ , then  $\text{Tev}_{g,n,dL}$  are well-defined for  $d \geq d[g, X]$  and coincide with  $\text{vTev}_{g,n,dL}$ .

## References

- A. Buch and R. Pandharipande, *Tevelev degrees in Gromov-Witten theory*, arXiv:2112.14824.
- G. Castelnuovo, *Numero delle involuzioni razionali giacenti sopra una curva di dato genere*, Rendiconti R. Accad. Lincei 5 (1889), 130-133.
- A. Cela, *Quantum Euler class and virtual Tevelev degrees of Fano complete intersections*, https://arxiv.org/abs/2204.01151.
- A. Cela, C. Lian, *Generalized Tevelev degrees of  $\mathbb{P}^1$* , arXiv:2111.05880.
- A. Cela, R. Pandharipande, J. Schmitt *Tevelev degrees and Hurwitz moduli spaces*, Math. Proc. Cambridge Philos. Soc..
- G. Farkas, C. Lian *Linear series on general curves with prescribed incidence conditions*, arXiv:2105.09340.
- W. Fulton, R. Pandharipande *Notes on stable maps and quantum cohomology*, Algebraic geometry-Santa Cruz 1995, volume 62 of Proc. Sympos. Pure Math., pages 45-96. Amer. Math. Soc., Providence, RI, 1997.
- J. Harris and D. Mumford, *On the Kodaira dimension of the moduli space of curves* With an appendix by William Fulton, Invent. Math. 67 (1982), 23-88.
- C. Lian, R. Pandharipande *Enumerativity of virtual Tevelev degrees*, arXiv:2110.05520.
- J. Tevelev, *Scattering amplitudes of stable curves*, arXiv:2007.03831.