

Reducibility and nonlinear stability for a quasi-periodically forced NLS

E. Haus* B. Langella†, A. Maspero‡ M. Procesi§

January 4, 2023

Abstract

Motivated by the problem of long time stability vs. instability of KAM tori of the Nonlinear cubic Schrödinger equation (NLS) on the two dimensional torus $\mathbb{T}^2 := (\mathbb{R}/2\pi\mathbb{Z})^2$, we consider a quasi-periodically forced NLS equation on \mathbb{T}^2 arising from the linearization of the NLS at a KAM torus. We prove a reducibility result as well as long time stability of the origin. The main novelty is to obtain the precise asymptotic expansion of the frequencies which allows us to impose Melnikov conditions at arbitrary order.

1 Introduction and main results

The nonlinear cubic Schrödinger equation on the two dimensional torus $\mathbb{T}^2 := (\mathbb{R}/2\pi\mathbb{Z})^2$

$$i\partial_t v = -\Delta v + |v|^2 v, \quad x \in \mathbb{T}^2 \tag{1.1}$$

is one of the fundamental equations in mathematical physics. The pioneering work by Bourgain [11] ensures that the equation is locally well-posed for any data in $H^s(\mathbb{T}^2)$ for all $s > 0$, where

$$\|u\|_{H^s} := \left(\sum_{n \in \mathbb{Z}^2} (1 + |n|)^{2s} |\hat{u}_n|^2 \right)^{\frac{1}{2}}.$$

Much less is known on the long time behaviour of the solutions, and in the last ten years a rich and diverse dynamics has been discovered for (1.1). Such dynamics comprise both “regular” dynamics, such as periodic [25], quasiperiodic [16, 26, 38], almost-periodic [13, 10] trajectories contained in invariant tori, and more “irregular” one, such as weakly turbulent trajectories undergoing energy cascade phenomena [14, 24, 22, 27, 21].

A particularly interesting long term goal is to understand the interplay of these two behaviours, in particular whether there are weakly turbulent trajectories connecting distinct invariant tori. An important intermediate step is to understand the local dynamics close to these invariant

*Dipartimento di Matematica e Fisica, Università di Roma Tre, Largo S.L. Murialdo, 00100 Roma, Italy, ehaus@mat.uniroma3.it

†International School for Advanced Studies (SISSA), Via Bonomea 265, 34136, Trieste, Italy beatrice.langella@sissa.it

‡International School for Advanced Studies (SISSA), Via Bonomea 265, 34136, Trieste, Italy alberto.maspero@sissa.it

§Dipartimento di Matematica e Fisica, Università di Roma Tre, Largo S.L. Murialdo, 00100 Roma, Italy, procesi@mat.uniroma3.it

sets, namely if one takes an initial datum close to an invariant torus, for how long does its trajectory stay close to the torus? The next step is to show that, beyond the stability times, unstable phenomena occur, and one may explicitly construct orbits which start δ -close to an invariant torus and then slowly drift away by a finite but arbitrarily large factor. A convenient way of ensuring that trajectories do not stay close to a given torus is to show that they exhibit a Sobolev norm explosion, namely that after some (very long) time their Sobolev norm, in some fixed $H^s(\mathbb{T}^2)$ with $s \neq 0, 1$, becomes larger than an arbitrary number K (recall that on a given invariant torus the Sobolev norms are essentially constant).

The simplest invariant object is clearly the fixed point $v = 0$. In this case stability results were proved for example in [5, 17, 9, 4, 6], while instability results in [12, 32, 33, 34], the landmark result in [14] and subsequent generalizations [24, 28, 22]. For these problems a fundamental role is played by the linearized frequencies at $v = 0$ of (1.1), namely

$$\Omega_j = |j|^2, \quad j \in \mathbb{Z}^2, \quad (1.2)$$

and the interest is either to avoid resonances between them to prove stability results or to exploit them to create instability phenomena.

A much more challenging problem is to study stability/instability of more complicated invariant objects. The simplest nontrivial invariant objects are the invariant tori of dimension 1 filled with plane wave solutions of the form $u(t, x) = \rho e^{i(jx - \omega t)}$, $\omega = |j|^2 + \rho^2$. Their long time stability was proved in [18], whereas their instability in H^s , $s \in (0, 1)$, in [27]. Again a fundamental role is played by the resonance properties of the linearized frequencies at plane waves solutions, which are easily computed to be

$$\Omega_j = \sqrt{|j|^4 + 2|j|^2\rho^2} = |j|^2 + \frac{c_j(\rho)}{|j|^2}, \quad j \in \mathbb{Z} \setminus \{0\}. \quad (1.3)$$

Here it is important to notice how the correction to the unperturbed frequencies (1.2) decays in $|j|$, namely whatever is the direction of the vector $j \in \mathbb{Z}^2$.

The next studied invariant objects have been the so-called finite gap solutions, which are solutions for the integrable 1d cubic NLS. In particular, they are solutions of (1.1) depending only on 1 variable, say x_1 , but they are quasi-periodic in time and fill invariant tori of finite dimension $\mathbf{d} \in \mathbb{N}$. Their long time stability has been studied in [35] (see also [29]) whereas their instability, again in H^s with $s \in (0, 1)$ in [21]. As the reader might guess, again a fundamental role is played by the linearized frequencies at finite gap solutions. In this case it is quite involved to compute their expression, and it requires partial Birkhoff normal form, a sort of “1d pseudodifferential normal form” and KAM reducibility techniques. The final result is that they expand as

$$\Omega_j = |j|^2 + \mu(j) + \frac{\mathbf{a}(m)}{\langle m \rangle^2} + \frac{R_j}{\langle j \rangle^2}, \quad j = (m, n) \in \mathbb{Z}^2 \setminus \{0\}, \quad (1.4)$$

where $\mu(j)$ has a finite number of possible values while \mathbf{a}, R are of size ε . Note the first difference with (1.2) and (1.3): the correction to the unperturbed frequencies depends on the direction of the vector $j = (m, n) \in \mathbb{Z}^2$.

Towards stability/instability of KAM tori. The next natural step is to study the long time stability vs instability of more general (truly bidimensional) quasiperiodic tori of Procesi-Procesi [38]. There are a number of very serious difficulties to overcome in order to deal with this more general case. Again the main challenge is to obtain the correct asymptotic expansion of the linearized frequencies at KAM tori, namely the analogous of formulas (1.3), (1.4). This is a very hard problem which requires new ideas with respect to the finite gap case.

The goal of this paper is to settle the machinery to obtain such asymptotics and to apply them to study a simplified NLS-like equation which, despite not having a particular physical interest, contains the main difficulties to be overcome in order to tackle the full problem of stability/instability of KAM tori.

In order to justify the model and the simplifications that we will introduce, let us first give a brief description of how the KAM-solutions appear. The first step is to choose an appropriate approximate solution, typically a solution of the linearized equation at $v = 0$, which is the starting point of a quadratic algorithm which converges to a true solution. The idea in [26, 37, 43] is to fix a finite set $\mathcal{S} = \{\mathbf{k}^{(1)}, \dots, \mathbf{k}^{(d)}\} \subset \mathbb{Z}^2$ (of maximal rank), look for solutions of (1.1) close to

$$v_{\text{lin}}(t, x) = \sum_{j \in \mathcal{S}} \sqrt{\xi_j} e^{ij \cdot x + i|j|^2 t},$$

and show that for *most* choices of \mathcal{S} and *many* choices of $\xi \in \mathbb{R}_+^d$ (in some small ball of radius ε) there exists a true solution q^{qp} of (1.1) essentially supported on the Fourier modes in \mathcal{S} which is quasi-periodic in time with frequency

$$\omega = (\omega_i)_{i=1}^d, \quad \omega_i = |\mathbf{k}^{(i)}|^2 + \mathcal{O}(\varepsilon).$$

Since the NLS equation is Gauge and translation invariant it turns out that the solutions in [38] have the form of *covariant quasi-periodic traveling waves*, according to the following definition:

Definition 1.1 (Quasi-periodic traveling waves). *Let $\mathbf{K} \in \text{Mat}_{d \times 2}(\mathbb{Z})$ (of maximal rank) with*

$$\mathbf{K}^T = \begin{pmatrix} \mathbf{k}_1^{(1)} & \dots & \mathbf{k}_1^{(d)} \\ \mathbf{k}_2^{(1)} & \dots & \mathbf{k}_2^{(d)} \end{pmatrix}. \quad (1.5)$$

A function $q(\varphi, x)$ is called a quasi-periodic traveling wave if it has the form

$$q(\varphi, x) := Q(\varphi + \mathbf{K}x)$$

where $Q: \mathbb{T}^d \rightarrow \mathbb{C}$ is a real analytic function.

We say that a traveling wave is Gauge covariant if Q is a function with Fourier coefficients supported in the set $\{\ell: \ell \cdot \vec{\mathbf{1}} = 1\}$, i.e.

$$Q(\varphi + t\vec{\mathbf{1}}) = e^{it} Q(\varphi), \quad \vec{\mathbf{1}} = (1, \dots, 1) \in \mathbb{Z}^d. \quad (1.6)$$

We say it is Gauge invariant if supported in the set $\{\ell: \ell \cdot \vec{\mathbf{1}} = 0\}$. Finally we set

$$\|Q\|_{\mathbf{a}, \mathbf{p}}^2 := \sum_{\ell} e^{2\mathbf{a}|\ell|} |Q(\ell)|^2 \langle \ell \rangle^{2\mathbf{p}} < \infty, \quad \text{for } \mathbf{a} > 0, \mathbf{p} > \frac{d}{2}. \quad (1.7)$$

Note that Gauge covariant, quasi-periodic traveling waves satisfy

$$q(\varphi + \mathbf{K}\zeta, \cdot) = \tau_{\zeta} q(\varphi, \cdot), \quad \forall \zeta \in \mathbb{R}^2 \quad (1.8)$$

$$q(\varphi + t\vec{\mathbf{1}}, \cdot) = e^{it} q(\varphi, \cdot), \quad \forall t \in \mathbb{R}, \quad (1.9)$$

where τ_{ζ} is the translation operator $(\tau_{\zeta} u)(x) := u(x + \zeta)$. In this sense, the solutions in [38] are covariant quasiperiodic traveling waves of the form

$$q^{\text{qp}}(\omega t, x) = q^{\text{qp}}(\varphi, x)|_{\varphi=\omega t} = Q(\omega t + \mathbf{K}x) \quad (1.10)$$

where $\omega \in \mathbb{R}^d$ belongs to a Cantor-like set of positive measure and \mathbf{K} as in (1.5).

Linearizing (1.1) at q^{qp} , one gets the equation

$$i\partial_t u = -\Delta u + A(\omega t)u + \mathcal{N}(\omega t, u) \quad (1.11)$$

with

$$\begin{aligned} A(\varphi)u &:= 2|q^{\text{qp}}(\varphi, \cdot)|^2 u + q^{\text{qp}}(\varphi, \cdot)^2 \bar{u} , \\ \mathcal{N}(\varphi, u) &:= 2q^{\text{qp}}(\varphi, \cdot)|u|^2 + \bar{q}^{\text{qp}}(\varphi, \cdot)u^2 + |u|^2 u . \end{aligned} \quad (1.12)$$

As the original equation (1.1) is gauge and translation covariant and $q^{\text{qp}}(\varphi, x)$ is a gauge covariant quasi-periodic traveling wave, the linear operator $A(\varphi)$ and the nonlinearity $\mathcal{N}(\varphi, u)$ fulfill the following covariant properties, as we show in Appendix C: for any $(\varphi, \zeta, t) \in \mathbb{T}^d \times \mathbb{R}^2 \times \mathbb{R}$

$$A(\varphi + t\vec{1}) \circ e^{it} = e^{it} \circ A(\varphi), \quad A(\varphi + \mathbf{K}\zeta) \circ \tau_\zeta = \tau_\zeta \circ A(\varphi), \quad (1.13)$$

$$\mathcal{N}(\varphi + t\vec{1}, e^{it}u) = e^{it} \circ \mathcal{N}(\varphi, u), \quad \mathcal{N}(\varphi + \mathbf{K}\zeta, \tau_\zeta u) = \tau_\zeta \circ \mathcal{N}(\varphi, u). \quad (1.14)$$

The fact that one may reduce to constant coefficients the operator $A(\varphi)$ was already discussed in [37, 38] (and is actually simpler to prove than in the case of finite gap solutions [35]). On the other hand the methods of [37, 38] do not give any information on the asymptotics of the frequencies, and actually one can prove that an expansion as the one in (1.4) cannot hold. Then one is not able to prove that the non-resonance conditions hold and one may not prove neither the stability result nor the successive instability one.

The results. As we already mentioned, the goal of this paper is to initiate the analysis of stability/instability of the general case by studying the nonlinear stability problem for a simplified *non-resonant* model, which contains only the difficulties related to the Melnikov conditions and avoids the further algebraic complications (which one expects to deal with exactly as in [37] and [35]). To this purpose we study the following simplified model:

$$iu_t - \Delta u + \mathcal{V}(\omega t, x)u + \partial_{\bar{u}}\mathcal{P}(t, x, u, \bar{u}) = 0, \quad x \in \mathbb{T}^2. \quad (1.15)$$

We make the following assumptions on \mathcal{V} and \mathcal{P} :

(H \mathcal{V}) $\mathcal{V}(\omega t, x)$ is the multiplication operator by an analytic gauge invariant traveling wave (see Definition 1.1), namely

$$\mathcal{V}(\omega t, x) = V(\omega t + \mathbf{K}x), \quad V(\varphi) = \sum_{\ell \in \mathbb{Z}^d} V(\ell)e^{i\ell \cdot \varphi}, \quad (1.16)$$

with the constraint that

$$V(\ell) = 0 \quad \forall \ell: \pi(\ell) = 0, \quad \text{or} \quad \sum_{i=1}^d \ell_i \neq 0. \quad (1.17)$$

(H \mathcal{P}) The non linear term $\mathcal{P}(t, x, y_1, y_2) = P(\omega t + \mathbf{K}x, y_1, y_2)$ with P an analytic function on $\mathbb{T}^d \times B_{\mathbf{r}_0}(0)$,

$$P(\varphi, y_1, y_2) = \sum_{\ell, d_1, d_2} P_{\ell, d_1, d_2} e^{i\ell \cdot \varphi} y_1^{d_1} y_2^{d_2}, \quad \|P\|_{\mathbf{a}, \mathbf{r}_0} = \sum_{\ell, d_1, d_2} |P_{\ell, d_1, d_2}| e^{|\ell|} \mathbf{r}_0^{d_1 + d_2} < \infty,$$

with a zero of order at least two and satisfying the invariance property

$$P(\varphi + t\vec{1}, e^{it}y_1, e^{-it}y_2) = P(\varphi, y_1, y_2).$$

Remark 1.2. We model \mathcal{V} on the first term of the operator $A(\varphi)$ in (1.12), namely the multiplication operator by $|q^{\text{qp}}(\omega t, x)|^2$, which is a gauge invariant traveling wave with $\int_{\mathbb{T}} |q^{\text{qp}}(\omega t, x)|^2 dx$ constant in time. So we assume that the same applies to \mathcal{V} .

Remark 1.3. The operator $\mathcal{V}(\varphi, \cdot)$ and the nonlinearity $\mathcal{P}(\varphi, \cdot)$ fulfill (1.13), (1.14).

For any $p \geq 0$ we introduce the space $\mathfrak{H}^p = \mathfrak{H}^p(\mathbb{Z}^2) := \ell^1(\mathbb{Z}^2) \cap h^p(\mathbb{Z}^2)$ endowed with the norm

$$\|u\|_p := \|u\|_{\ell^1(\mathbb{Z}^2)} + \|u\|_{h^p(\mathbb{Z}^2)}.$$

Note that, by Young inequality, the space \mathfrak{H}^p is an algebra with respect to the convolution product, i.e.

$$\|u * v\|_p \leq 2^{2p+1} \|u\|_p \|v\|_p \quad (1.18)$$

We are now ready to state our stability result.

Theorem 1.4. Fix $p > 0$ and consider the equation (1.15). Assume $(H\mathcal{V})$ and $(H\mathcal{P})$. Let $\mathbf{R} := [-1, 1]^d$. There exist $\gamma_*, \epsilon_* > 0$ and for any $\gamma \in (0, \gamma_*)$, and any potential V satisfying

$$\epsilon := \gamma^{-1} \|V\|_{\text{a.p}} < \epsilon_* , \quad (1.19)$$

there exist a Cantor-like set \mathcal{C} with measure $\text{meas}(\mathbf{R} \setminus \mathcal{C}) \leq C\gamma$, and $r_*, T_* > 0$ such that for any $\omega \in \mathcal{C}$, $r \in (0, r_*)$ the following holds true. For any $u \in \mathfrak{H}^p$, with $\|u(0)\|_p \leq r$, the solution $u(t) \in \mathfrak{H}^p$ of (1.15) fulfills

$$\|u(t)\|_p \leq 2r, \quad \forall |t| \leq \frac{T_*}{r^2} . \quad (1.20)$$

As already mentioned, the proof of Theorem 1.4 follows the same general scheme of [35] based on the three steps: 1. Reducibility, 2. Non-resonance conditions, 3. Order three Birkhoff Normal Form. Since the last step is standard (it is briefly discussed in Section 6) we concentrate on the first two.

We start by studying the reducibility for the Schrödinger operator

$$A = i\mathfrak{L}, \quad \mathfrak{L} = -\Delta + \mathcal{V}, \quad (1.21)$$

in the space $\mathfrak{H}^p := \ell^1(\mathbb{Z}^2) \cap h^p(\mathbb{Z}^2)$, where $p > 0$ is given, which amounts to prove that the following property holds:

Definition 1.5 (Reducibility). We say that a time dependent Schrödinger operator \mathfrak{L} is reducible on \mathfrak{H}^p if there exists a quasi-periodic in time bounded and invertible linear operator G such that

$$G_* \mathfrak{L} := -iG^{-1} \dot{G} + G^{-1} \mathfrak{L} G = \text{diag}(\Omega_j), \quad \Omega_j \in \mathbb{R}. \quad (1.22)$$

In [15], Eliasson and Kuksin have proven in the analytic setting the existence of an invertible and time quasi-periodic linear operator G such that $G_* \mathfrak{L}$ as in (1.22) is reduced to a time independent, block diagonal operator Q , under smallness assumptions on ϵ as in (1.19) and for a Borel set of frequencies ω of asymptotically full measure in ϵ , without the constraint that \mathcal{V} is a quasi-periodic traveling wave. Furthermore, it is well known (see for instance [41]) that if \mathfrak{L} is covariant in the sense of (1.13), then Q is diagonal. However, the reducibility result of [15] does not provide an asymptotic expansion of the frequencies $\{\Omega_j\}_{j \in \mathbb{Z}^2}$. Thus here we proceed to prove a more refined result, from which we obtain an asymptotic expansion for $\{\Omega_j\}_{j \in \mathbb{Z}^2}$ that generalizes the one found in [21] in the finite gap case (see (1.4)). To state our result we first need two definitions.

Definition 1.6 (Generators). A vector $v = (v_1, v_2) \in \mathbb{Z}^2 \setminus \{0\}$ shall be called a generator if

$$\gcd(v_1, v_2) = 1 \text{ and } v_1 > 0 \quad \text{or} \quad v = (0, 1) . \quad (1.23)$$

We shall denote by $\mathcal{V} \subset \mathbb{Z}^2 \setminus \{0\}$ the set of generators.

From now on, we fix

$$0 < \delta \ll 1, \quad \text{and} \quad \mu := 1 - 2\delta . \quad (1.24)$$

Definition 1.7. Given $j \in \mathbb{Z}^2 \setminus \{0\}$, we define $v(j) \in \mathcal{V}$ as the first vector w.r.t. the lexicographic ordering that attains the minimum in

$$\min_{v \in \mathcal{V}: |v| \leq |j|^\delta} |v \cdot j| \quad (1.25)$$

and set $b(j) := j \cdot v(j)$. This means that $|b(j)| = \min \{|v \cdot j| \mid v \in \mathcal{V} : |v| \leq |j|^\delta\}$.

With these notations, we prove the following:

Theorem 1.8. There exist $\gamma_*, \epsilon_* > 0$ and for any $\gamma \in (0, \gamma_*)$, any potential V satisfying (1.19), there exist a Cantor-like set \mathcal{C} with measure $\text{meas}(\mathbb{R} \setminus \mathcal{C}) \leq C\gamma$, a sequence of Lipschitz functions $\mathbb{R} \ni \omega \mapsto \Omega_j(\omega)$ and a bounded linear operator G , close to the identity and with quasi-periodic dependence on time, such that for any $\omega \in \mathcal{C}$

$$G_* \mathfrak{L} = \text{diag}(\Omega_j) , \quad (1.26)$$

with

$$\begin{aligned} \Omega_j(\omega) &:= \Omega_j = |j|^2 + \frac{\varpi(v(j), b(j), \omega)}{\langle b(j) \rangle^2} e^{-\kappa|v(j)|} + \frac{\Theta^{(1)}(j, \omega)}{\langle b(j) \rangle \langle j \rangle^\mu} + \frac{\Theta^{(2)}(j, \omega)}{\langle j \rangle^{2\mu}} \quad \forall j, \\ \sup_{\omega \in \mathbb{R}} \sup_{j \in \mathbb{Z}^2} \left(|\varpi(b(j), v(j), \omega)| + |\Theta^{(1)}(j, \omega)| + |\Theta^{(2)}(j, \omega)| \right) &< 2\|V\|_{\mathbf{a}, \mathbf{p}}, \end{aligned} \quad (1.27)$$

with $\kappa > 0$ depending on \mathbf{a} and on the matrix \mathbf{K} only. Moreover, for any integer $N \geq 3$, there exist $\tau_N > 0, c_N^* > 0$ and a Cantor set $\mathcal{C}_N \subseteq \mathcal{C}$, such that $\text{meas}(\mathbb{R} \setminus \mathcal{C}_N) \leq c_N^* \gamma$, and $\forall \omega \in \mathcal{C}_N, \forall (\ell, L) \in \mathcal{G}_N$, where

$$\begin{aligned} \mathcal{G}_N &:= \left\{ (\ell, L) \in \mathbb{Z}^d \times \mathbb{Z}^{\mathbb{Z}^2} : \text{if } \ell = 0, \quad \text{then} \quad \sum_{j \in \mathbb{Z}^2} |j|^2 L_j \neq 0 \right. \\ &\quad \left. |L| := \sum_{j \in \mathbb{Z}^2} |L_j| \leq N, \quad \sum_{i=1}^d \ell_i + \sum_{j \in \mathbb{Z}^2} L_j = 0, \quad \sum_{i=1}^d \ell_i \mathbf{k}^{(i)} + \sum_{j \in \mathbb{Z}^2} j L_j = 0 \right\}, \end{aligned} \quad (1.28)$$

one has

$$|\omega \cdot \ell + \Omega \cdot L| \geq \gamma \langle \ell \rangle^{-\tau_N} . \quad (1.29)$$

The major step in proving non-resonance conditions (1.29) is to show that the final frequencies $\{\Omega_j\}_{j \in \mathbb{Z}^2}$ satisfy the asymptotic expansions (1.27). Actually the problem of exhibiting asymptotic expansions for the eigenvalues of a linear Schrödinger operator

$$\tilde{\mathfrak{L}} = -\Delta + W$$

on $\mathbb{T}^n, n \geq 1$, has been widely investigated already in the case of a time independent W (see for instance [19, 20, 42, 30, 31, 39, 40, 7, 8]). Roughly speaking, in such works one partitions the

spectrum of \mathfrak{L} into two subsets: the stable spectrum, which is composed by the eigenvalues such that

$$\Omega_j = |j|^2 + O\left(\frac{1}{\langle j \rangle^{2\mu}}\right), \quad (1.30)$$

and the unstable one, namely the set of eigenvalues for which (1.30) does not hold. We point out that with our notations, stable eigenvalues correspond to those such that

$$\langle b(j) \rangle \geq \langle j \rangle^\mu. \quad (1.31)$$

Refined asymptotic expansions were available in the time independent case also for eigenvalues in the unstable spectrum, and in [8] it was proven that all the eigenvalues of $\tilde{\mathfrak{L}}$ on \mathbb{T}^2 are of the form

$$\Omega_j = |j|^2 + \sum_{n=1}^N \frac{\varpi_n^*(j)}{\langle b(j) \rangle^{2n}} + \sum_{n=1}^N \frac{\Theta_n^*(j)}{\langle j \rangle^{2n\mu}} + O\left(\frac{1}{\langle b(j) \rangle^{2N}}\right), \quad (1.32)$$

using techniques based on pseudo-differential calculus and a geometric decomposition of the space of indexes j à la Nekhoroshev (see [36, 23]). However, an expansion of the form (1.32) would not suffice in order to show non-resonance conditions (1.29) at any order N : this is the reason why we use a slightly different approach, based on a structure which we refer to as quasi-Töplitz (see Definition 2.16 below), and prove expansions (1.27). Remark that the major difference with respect to the expansions (1.32) is represented by the functions $\varpi(b(j), v(j))$, namely by the fact that the terms which decay with $\langle b(j) \rangle$ depend on the vector j through $v(j)$ and $b(j)$ only and decay exponentially in $v(j)$. This is the key to proving the non-resonance conditions (1.29).

Indeed, rewriting

$$\omega \cdot \ell + \Omega \cdot L = \omega \cdot \ell \pm \Omega_{j_1} \pm \Omega_{j_2} \pm \dots \pm \Omega_{j_k}$$

the asymptotic expansion guarantees that if all the $|j_i| \gg |\ell|^{a_1}$, and either all the $|b(j_i)| \gg |\ell|^{a_2}$ or $|v(j_i)| \gg |\ell|$, then (1.29) follows by imposing Diophantine conditions on ω of the form

$$|\omega \cdot \ell + K| \gtrsim |\ell|^{-\tau} \quad \forall \ell \neq 0, \quad \forall K \in \mathbb{Z}$$

(and it is well known that such conditions hold for a positive measure set of ω provided that τ is small enough).

Following this line of reasoning in (1.29) we can ignore all the terms $\frac{\Theta^{(1)}(j_i)}{\langle b(j_i) \rangle \langle j_i \rangle^\mu}, \frac{\Theta^{(2)}(j_i)}{\langle j_i \rangle^{2\mu}}$ where the $|j_i|$ are sufficiently large. Similarly we can ignore all the terms $\frac{\varpi(v(j_i), b(j_i))}{\langle b(j_i) \rangle^2} e^{-\kappa|v(j_i)|}$ where either $v(j_i)$ or $b(j_i)$ is large. This means that, for $|L| \leq N$, the expression $\omega \cdot \ell + \Omega \cdot L$ assumes (up to a negligible error) only a finite number of distinct values (of cardinality bounded only by ℓ) which are small.

In turn this means that one can surely impose conditions of the type (1.29) provided that τ_N is sufficiently large.

2 Functional setting

2.1 Time-dependent momentum preserving operators.

We shall consider quasi-periodic time dependent operators $t \mapsto M(\omega t) \in \mathcal{L}(\mathfrak{H}^p)$ with frequency $\omega \in \mathbb{R}^d$, where

$$M(\varphi) = \sum_{\ell \in \mathbb{Z}^d} M(\ell) e^{i\varphi \cdot \ell}, \quad M(\ell) \in \mathcal{L}(\mathfrak{H}^p), \quad (2.1)$$

that are analytic and momentum preserving according to the following definition:

Definition 2.1 (Momentum preserving operators). *Here and in the following we shall define the linear map $\pi : \mathbb{Z}^d \rightarrow \mathbb{Z}^2$*

$$\pi(\ell) := \sum_{i=1}^d \ell_i \mathbf{k}^{(i)}, \quad \mathbf{c}^{-1} := \sup_{\ell \neq 0} \frac{|\pi(\ell)|}{|\ell|}. \quad (2.2)$$

We say that a time dependent linear operator $M(\omega t)$ is momentum preserving if

$$j - j' \neq \pi(\ell) \Rightarrow M_j^{j'}(\ell) := \langle M(\ell) e^{ij' \cdot x}, e^{ij \cdot x} \rangle = 0. \quad (2.3)$$

For any $a > 0$ we define the norm

$$|M|_a := \sup_{\|u\|_p \leq 1} \|\underline{M}_a u\|_p, \quad (\underline{M}_a)_j^{j'} := \sum_{\ell: \pi(\ell)=j-j'} e^{a|\ell|} |M_j^{j'}(\ell)|.$$

We denote the space of time dependent momentum preserving operators with finite norm as $\mathcal{L}_a(\mathfrak{H}^p)$.

Definition 2.2 (Gauge covariant operators). *We say that M as above is Gauge covariant if $\sum_{i=1}^d \ell_i \neq 0$ implies $M_j^{j'}(\ell) = 0$.*

A convenient way of envisioning such operators is as *normally analytic* maps from a thickened torus

$$\mathbb{T}_a^d := \{\varphi \in \mathbb{C}^d : \operatorname{Re}(\varphi) \in \mathbb{R}^d / (2\pi\mathbb{Z})^d, \quad |\operatorname{Im}(\varphi)| \leq a\} \rightarrow \mathcal{L}(\mathfrak{H}^p). \quad (2.4)$$

Remark 2.3. *If $M \in \mathcal{L}_a(\mathfrak{H}^p)$, then $M(\varphi) \in \mathcal{L}(\mathfrak{H}^p)$ for all $\varphi \in \mathbb{T}_a^d$ and the operator norm satisfies $\|M(\varphi)\|_{\mathcal{L}(\mathfrak{H}^p)} \leq |M|_a$.*

One may easily verify (see Lemma 2.14) that $\mathcal{L}_a(\mathfrak{H}^p)$ is in fact an algebra with respect to composition. Then it is standard to define the commutator and the adjoint action as follows:

Definition 2.4. *For $M, S \in \mathcal{L}_a(\mathfrak{H}^p)$, we set $\operatorname{ad}S[M](\varphi) := [M(\varphi), S(\varphi)] := M(\varphi)S(\varphi) - S(\varphi)M(\varphi)$, which in matrix form reads*

$$([M, S])_j^{j-\pi(\ell)}(\ell) = \sum_{\ell_1 + \ell_2 = \ell} M_j^{j-\pi(\ell_1)}(\ell_1) S_{j-\pi(\ell_1)}^{j-\pi(\ell)}(\ell_2) - S_j^{j-\pi(\ell_1)}(\ell_1) M_{j-\pi(\ell_1)}^{j-\pi(\ell)}(\ell_2).$$

Definition 2.5. *We say that $M \in \mathcal{L}_a(\mathfrak{H}^p)$ is self-adjoint if it satisfies*

$$M_j^{j-\pi(\ell)}(\ell) = \overline{M_{j-\pi(\ell)}^j(-\ell)} \quad (2.5)$$

so that $M(\varphi)$ is self-adjoint for all $\varphi \in \mathbb{T}^d$. As is standard, we say that a bounded operator $G \in \mathcal{L}_a(\mathfrak{H}^p)$ is symplectic if $G = e^{iA}$ with $A \in \mathcal{L}_a(\mathfrak{H}^p)$ self-adjoint.

Remark 2.6. *If A, B are self-adjoint, then so is $i[A, B]$. Consequently, anti self-adjoint operators form a Lie algebra. Moreover, if M is self-adjoint and $G = e^S$ is symplectic (i.e., iS is self-adjoint) then $G^{-1}MG = \exp(\operatorname{ad}(S))M$ is self-adjoint. Similarly Gauge covariant operators form a Lie algebra, so if S, M are Gauge covariant, so is G and $G^{-1}MG$.*

Time-dependent changes of variables Given the linear PDE $u_t = i\mathfrak{L}u$, where \mathfrak{L} is a time dependent Schrödinger operator as in (1.21) with $P \in \mathcal{L}_a(\mathfrak{H}^p)$, let $G \in \mathcal{L}_a(\mathfrak{H}^p)$ be a quasi-periodic in time bounded invertible change of variables of the form $G = e^S$ with $S \in \mathcal{L}_a(\mathfrak{H}^p)$. Then, defining $u := Gv$, one has $v_t = i(G_*\mathfrak{L})v$, with $G_*\mathfrak{L}$ as in (1.22). The Lie exponentiation formula gives a nice representation of $G_*\mathfrak{L}$ in terms of the adjoint actions:

$$\mathfrak{L}_1 := G_*\mathfrak{L} = e^{\text{ad}S}\mathfrak{L} - i \sum_{k=1}^{\infty} \frac{(\text{ad}S)^{k-1}}{k!} \dot{S}, \quad e^{\text{ad}S}\mathfrak{L} := \sum_{k=0}^{\infty} \frac{(\text{ad}S)^k}{k!} \mathfrak{L}. \quad (2.6)$$

Remark 2.7. *Note that, by Remark 2.6, if \mathfrak{L} is self-adjoint and G is symplectic (or equivalently iS is self-adjoint) then \mathfrak{L}_1 is self-adjoint. The same holds for the Gauge covariance.*

Symplectic structure As we have already explained in the introduction, time dependent Schrödinger operators have a natural Hamiltonian structure on the phase space $\mathbb{T}^d \times \mathbb{R}^d \times \mathfrak{H}^p$, equipped with the symplectic form $d\mathcal{Y} \wedge d\varphi + i \sum_j du_j \wedge d\bar{u}_j$. To this purpose we associate to \mathfrak{L} as in (1.21), with P self-adjoint, the Hamiltonian

$$H_{\mathfrak{L}} := \omega \cdot \mathcal{Y} + \sum_j |j|^2 |u_j|^2 + \sum_{j,\ell} P_j^{j-\pi(\ell)}(\ell) e^{i\ell \cdot \varphi} u_{j-\pi(\ell)} \bar{u}_j, \quad (2.7)$$

with Hamilton equations $\dot{\varphi} = \omega$, $\dot{u} = i\mathfrak{L}u$, $\dot{\mathcal{Y}} = -(u, P_{\varphi}u)_{\ell_2(\mathbb{C})}$. Accordingly, to a symplectic map $G = e^{iA}$ we associate the generating function

$$\mathcal{A}(\varphi, u) := \sum_{j,\ell} A_j^{j-\pi(\ell)}(\ell) e^{i\ell \cdot \varphi} u_{j-\pi(\ell)} \bar{u}_j,$$

whose time one flow $\Phi_{\mathcal{A}}^1$ gives the symplectic change of variables $(u, \varphi, \mathcal{Y}) = \mathcal{G}(v, \varphi', \mathcal{Y}') = \Phi_{\mathcal{A}}^1(v, \varphi', \mathcal{Y}')$

$$\varphi = \varphi', \quad \mathcal{Y} = \mathcal{Y}' + i(Gv, G_{\varphi}v)_{\ell_2(\mathbb{C})}, \quad u = G(\varphi)v. \quad (2.8)$$

Note that (by the Lie exponentiation formula) $\Phi_{\mathcal{A}}^1$ conjugates $(\Phi_{\mathcal{A}}^1)_* H_{\mathfrak{L}} = e^{\{\mathcal{A}, \cdot\}} H_{\mathfrak{L}} = H_{\mathfrak{L}_1}$, with \mathfrak{L}_1 defined in (2.6). Finally if P, A are Gauge covariant then so is \mathfrak{L}_1 .

Basic properties of $\mathcal{L}_a(\mathfrak{H}^p)$. We now list some useful properties of the space $\mathcal{L}_a(\mathfrak{H}^p)$.

Remark 2.8. *We shall systematically use the fact that the norm $|\cdot|_a$ of Definition 2.1 is ordered. Indeed, if $M(\varphi), N(\varphi)$ are such that $(\underline{M}_a)_{j'}^{j'} \leq (\underline{N}_a)_{j'}^{j'}$ for any j, j' , then $|M|_a \leq |N|_a$.*

Lemma 2.9. *Let M be a momentum preserving operator according to Definition 2.1. If there exist $\theta, A > 0$ such that*

$$(\underline{M}_a)_{j'}^{j'} \leq A e^{-\theta|j-j'|}, \quad \forall j, j' \in \mathbb{Z}^2,$$

then there exists $C(p) > 0$ such that

$$|M|_a \leq C(p) A \theta^{-p-2}.$$

Proof. We remark that $\|\underline{M}_a u\|_p \leq \|f * u\|_p$ where $f = (f_j)_{j \in \mathbb{Z}^2}$ has components $f_j := e^{-\theta|j|}$. Then one uses (1.18) and the bound follows. \square

Lemma 2.10. *Let \mathcal{V} be a traveling wave as in Definition 1.1 with $\|V\|_{\mathbf{a}, \mathbf{p}} < \infty$. Then the multiplication operator $M_{\mathcal{V}} : u \mapsto \mathcal{V}u$ belongs to $\mathcal{L}_a(\mathfrak{H}^p)$ for any $a \in [0, \mathbf{a})$ and*

$$|M_{\mathcal{V}}|_a \leq C(p, \mathbf{p}) (\mathbf{a} - a)^{-(p-2)} \|V\|_{\mathbf{a}, \mathbf{p}}.$$

Proof. The multiplication operator is represented by the matrix

$$(M_V)_j^{j'}(\ell) = V_{\ell, j-j'} = \begin{cases} V(\ell) & \text{if } j - j' = \pi(\ell) \\ 0 & \text{otherwise} \end{cases}, \quad (2.9)$$

where we used the fact that V is a traveling wave, see (1.16). Hence M_V is momentum preserving. To compute its norm, remark that $|j - j'| = |\pi(\ell)| \leq c^{-1}|\ell|$ hence, setting $\theta = c(\mathbf{a} - a)$,

$$\begin{aligned} (\underline{M}_{V_a})_j^{j'} &= \sum_{\ell: j-j'=\pi(\ell)} e^{a|\ell|} |V(\ell)| \leq e^{-\theta|j-j'|} \sum_{\ell} e^{a|\ell|} |V(\ell)| \\ &\leq e^{-\theta|j-j'|} \sqrt{\sum_{\ell} \frac{1}{\langle \ell \rangle^{2\mathbf{p}}}} \sqrt{\sum_{\ell} e^{2a|\ell|} |V(\ell)|^2 \langle \ell \rangle^{2\mathbf{p}}} \leq A(\mathbf{p}) e^{-\theta|j-j'|} \|V\|_{\mathbf{a}, \mathbf{p}}. \end{aligned}$$

Then use Lemma 2.9. \square

We now want to define the "order" of a momentum preserving operator (see Definition 2.13 below). On the one hand, an operator of order $-m$ is standardly defined as a linear map $\mathfrak{H}^p \rightarrow \mathfrak{H}^{p+m} \forall p$; on the other hand, if \mathfrak{H}^p represents a space of functions of two variables, $\mathfrak{H}^p(\mathbb{Z}^2) = \mathfrak{H}^p(\mathbb{Z}) \otimes \mathfrak{H}^p(\mathbb{Z})$, one can define a "vectorial order" such that an operator of order $-(n_1, n_2)$ maps

$$\mathfrak{H}^p(\mathbb{Z}) \otimes \mathfrak{H}^p(\mathbb{Z}) \rightarrow \mathfrak{H}^{p+n_1}(\mathbb{Z}) \otimes \mathfrak{H}^{p+n_2}(\mathbb{Z})$$

(see for instance [35]). Here actually the main novelty is that we use a "non standard" set of coordinates, which we define in the following.

2.2 A non-linear coordinate set on \mathbb{Z}^2

In this section we prove some properties of the quantities $b(j)$, $v(j)$ defined for all $j \in \mathbb{Z}^2$ in Definitions 1.6, 1.7.

Lemma 2.11. *Let $v, w \in \mathcal{V}$ be distinct generators with $\max(|v|, |w|) < R$. If $x \in \mathbb{Z}^2$ satisfies $\max(|x \cdot v|, |x \cdot w|) < A$, then $|x| < 2AR$.*

Proof. Denoting by $a := (x \cdot v, x \cdot w)^T$, x is the solution of the linear system $Mx = a$ where M is the matrix with rows v^T and w^T . Then one estimates M^{-1} by Cramer's rule. \square

We shall denote by $B_K(0)$ the ball of radius K and center 0 in \mathbb{Z}^2 .

Lemma 2.12. *There exists $J_\delta > 0$ with the following property. For any $j \notin B_{J_\delta}(0)$ such that $|b(j)| < 2|j|^\mu$ (see (1.24)), the following holds true:*

(i) *For all $w \in \mathcal{V}$ with $|w| \leq |j|^\delta$ and $w \neq v(j)$*

$$|w \cdot j| \geq 2\langle j \rangle^\mu > |v(j) \cdot j|,$$

namely $v(j)$ is the unique vector attaining the minimum in (1.25).

(ii) *If there exists $h \in \mathbb{Z}^2$ with $|j - h| < 2 \max(|j|, |h|)^\delta$ and $v(h) \neq v(j)$, then*

$$|v(j)| > \frac{1}{2}|j|^\delta.$$

Proof. (i) Assume by contradiction that there exists another generator $w \in \mathcal{V}$ such that $|w \cdot j| = |v(j) \cdot j| = |b(j)| < 2|j|^\mu$. Then, we apply Lemma 2.11 with $A = 2|j|^\mu$, $R = |j|^\delta$ and we deduce

$$|j| < 4|j|^{1-\delta} \quad \Rightarrow \quad |j| < 4^{\frac{1}{\delta}}$$

and item (i) follows provided $J_\delta \geq 4^{\frac{1}{\delta}}$.

(ii) We claim that, by taking J_δ large enough, one has

$$\frac{1}{2}|j| < |h| < 2|j|. \quad (2.10)$$

Indeed if $|h| \geq 2|j|$, then by triangular inequality one deduces that $|h| \leq 4|h|^\delta$, so in particular, being $\delta \ll 1$, one has $|h| \leq R_\delta$, same for $|j|$. Then just take $J_\delta > R_\delta$ to get a contradiction. The other inequality is analogous.

By the definition of $v(\cdot)$ we have $|v(j)| < |j|^\delta$, $|v(h)| < |h|^\delta$. We distinguish two cases. If $|v(j)| > |h|^\delta$ then the thesis follows by (2.10). If $|v(j)| \leq |h|^\delta$, then using the bounds on $b(j)$, $|j - h|$ and again (2.10) we get

$$|v(h) \cdot h| \leq |v(j) \cdot h| = |v(j) \cdot (j + h - j)| \leq 2|j|^\mu + 2|j|^\delta \max(|j|, |h|)^\delta < 4|h|^\mu$$

provided J_δ is sufficiently large and δ sufficiently small.

Then we apply Lemma 2.11 with $x = h$, $v = v(j)$, $w = v(h)$, $R = |h|^\delta$ and $A = 4|h|^\mu$. We deduce $|h| < 8|h|^{1-\delta}$, which leads to a contradiction for J_δ large. \square

Next we define the order of a momentum preserving operator.

Definition 2.13 (Order). *Given $\mathbf{N} = (n, m)$ with $n, m \geq 0$ and an operator $M \in \mathcal{L}_a(\mathfrak{H}^p)$, we define the following norm:*

$$|M|_{a;-\mathbf{N}} := \sup_{\|u\|_p \leq 1} \|\underline{M}_{a;-\mathbf{N}}u\|_p, \quad (\underline{M}_{a;-\mathbf{N}})_j^{j'} := \sum_{\ell: j-j'=\pi(\ell)} e^{a|\ell|} |M_j^{j'}(\ell)| \langle j \rangle^{\mu n} \langle b(j) \rangle^m, \quad (2.11)$$

where $\forall j \in \mathbb{Z}^2$, $b(j)$ is the quantity in Definition 1.7 and μ in (1.24). We denote by $\mathcal{L}_{a,-\mathbf{N}}$ the subspace of \mathcal{L}_a with finite $|\cdot|_{a,-\mathbf{N}}$ norm.

These operators form an algebra, as the next lemma shows. Let

$$C(p, \sigma) := c^{-p} \sup_{k \in \mathbb{N}} e^{-\sigma k} k^p \sim \sigma^{-p}. \quad (2.12)$$

Lemma 2.14 (Algebra property). *Given two operators M, N , with $|M|_{a;-\mathbf{N}_1}, |N|_{a;-\mathbf{N}_2} < \infty$, we have*

$$|MN|_{a;-\mathbf{N}_1} \leq |M|_{a;-\mathbf{N}_1} |N|_{a;\bar{0}}. \quad (2.13)$$

Moreover, let $\mathbf{N}_1 = (n_1, m_1)$ and $\mathbf{N}_2 = (n_2, m_2)$; then we have for any $a' < a$

$$|MN|_{a';-(n_1+n_2,0)} \leq 2^{\mu n_2} |M|_{a';-\mathbf{N}_1} |N|_{a';-\mathbf{N}_2} + C(\mu(n_1+n_2)/\delta, a-a') |M|_{a;-\mathbf{N}_1} |N|_{a';-\mathbf{N}_2}. \quad (2.14)$$

The proof, being quite technical, is postponed to Appendix A.

Remark 2.15. *We note that if $M \in \mathcal{L}_a(\mathfrak{H}^p)$ is time independent, then (by formula (2.3), and recalling that $\pi(0) = 0$) it is necessarily diagonal, namely there exists a sequence $\{M_j\}_{j \in \mathbb{Z}^2}$, $M_j \in \mathbb{C}$, with $M = \text{diag}(M_j)_{j \in \mathbb{Z}^2}$. Moreover one has $\sup_{j \in \mathbb{Z}^2} |M_j| = |M|_a$, $\forall a$.*

2.3 Quasi-Töplitz norm

The purpose of this section is to define, for a fixed $m > 0$, a decomposition of operators in $\mathcal{L}_a(\mathfrak{H}^p)$ into a sum of terms of order $-\mathbf{N} = -(m-k, k)$ for all $k = 0, \dots, m$. We shall further require an invariance property on the term of order $-(0, m)$, which we will refer to as line-Töplitz.

Definition 2.16. *An operator $M \in \mathcal{L}_a(\mathfrak{H}^p)$ is said to be:*

- Line-Töplitz if there exists a map $\mathbb{Z}^d \times \mathcal{V} \times \mathbb{Z} \rightarrow \mathbb{C}$, $(\ell, v, b) \mapsto \mathfrak{M}(\ell, v, b)$, such that

$$M_j^{j'}(\ell) = \mathfrak{M}(\ell, v(j), b(j)), \quad \forall \ell \in \mathbb{Z}^d, j, j' \in \mathbb{Z}^2.$$

We denote by $\mathcal{T}_a := \mathcal{T}_a(\mathfrak{H}^p)$ the set of line-Töplitz operators in $\mathcal{L}_a(\mathfrak{H}^p)$.

- Line-Töplitz of order $-m$, $m \geq 0$, if

$$|M|_{a, -m}^{\mathbf{T}} := \sup_{v \in \mathcal{V}, b \in \mathbb{Z}} \sum_{\ell \in \mathbb{Z}^d} e^{a\mathfrak{c}|v| + a|\ell|} |\mathfrak{M}(\ell, v, b)| \langle b \rangle^m < +\infty \quad (2.15)$$

where \mathfrak{c} is defined in (2.2). We denote by $\mathcal{T}_{a, -m}$, $m \geq 0$, the set of line-Töplitz operators of order m .

- Quasi-Töplitz of order $-m$, $m \in \mathbb{N}$, if there exist a line-Töplitz operator $M^{\mathbf{T}} \in \mathcal{T}_{a, -m}$ and m operators $M^{(i)} \in \mathcal{L}_{\frac{a}{2}}(\mathfrak{H}^p)$, $i = 1, \dots, m$ such that

$$M = M^{\mathbf{T}} + \sum_{i=1}^m M^{(i)}, \quad M^{(i)} \in \mathcal{L}_{\frac{a}{2}; -(i, m-i)}. \quad (2.16)$$

We denote by $\mathcal{L}_{a, -m}^{\mathfrak{qT}}$ the set of quasi-Töplitz operators of order m of $\mathcal{L}_a(\mathfrak{H}^p)$, which we endow with the norm

$$|M|_{a, -m}^{\mathfrak{qT}} := \inf \left\{ |M^{\mathbf{T}}|_{a, -m}^{\mathbf{T}} + \sum_{i=1}^m |M^{(i)}|_{\frac{a}{2}; -(i, m-i)} \quad \middle| \quad M = M^{\mathbf{T}} + \sum_{i=1}^m M^{(i)} \right\}.$$

Lemma 2.17. *The line-Töplitz operators are bounded, more precisely for all $a' \in (0, a)$ one has*

$$|M|_{a'; -(0, m)} \leq C(\mathfrak{c}, p)(a - a')^{-p-2} |M|_{a, -m}^{\mathbf{T}}$$

Proof. We have¹

$$(\underline{M}_{a'; -(0, m)})_j^{j'} = \sum_{\ell: j-j'=\pi(\ell)} e^{a|\ell|} \langle b(j) \rangle^m |\mathfrak{M}(\ell, v(j), b(j))| e^{-(a-a')|\ell|} \lesssim |M|_{a, -m}^{\mathbf{T}} e^{-c(a-a')|j-j'|}.$$

Let $g_j := e^{-c(a-a')|j|}$ and $g := \{g_j\}_{j \in \mathbb{Z}^2} \in \mathfrak{H}^p$. We have

$$|g|_p := \sqrt{\sum_j e^{-2c(a-a')|j|} \langle j \rangle^{2p} + \sum_j e^{-c(a-a')|j|}} \lesssim_{\mathfrak{c}, p} (a - a')^{-p-2}$$

It follows from the algebra property of \mathfrak{H}^p w.r.t. convolution that

$$|M|_{a'; -(0, m)} \lesssim |g|_p |M|_{a, -m}^{\mathbf{T}} \lesssim_{\mathfrak{c}, p} (a - a')^{-p-2} |M|_{a, -m}^{\mathbf{T}}.$$

□

¹As is conventional we write $A \lesssim B$ if there exists a universal constant $C > 0$ such that $A \leq CB$. Similarly we write $A \lesssim_{\mathfrak{c}, p} B$ if there exists $C > 0$ depending only on \mathfrak{c}, p such that $A \leq CB$.

We shall need to keep track of Lipschitz dependence on the parameter $\omega \in \mathbb{R}^d$. To this purpose, given a compact set $\mathcal{O} \subset \mathbb{R}^d$, we fix $\gamma > 0$ and define the following norm on the space of Lipschitz maps $f : \mathcal{O} \rightarrow E$ (E a Banach space)

$$|f|_E^{\mathcal{O}} := \sup_{\omega \in \mathcal{O}} |f(\omega)|_E + \gamma \sup_{\omega \neq \omega' \in \mathcal{O}} |\Delta_{\omega, \omega'} f|_E, \quad \Delta_{\omega, \omega'} f := \frac{f(\omega) - f(\omega')}{|\omega - \omega'|}.$$

Definition 2.18. Let $\mathcal{O} \subset \mathbb{R}^d$ be compact. We denote by $\mathcal{L}_{a, -m}^{\text{qT}, \mathcal{O}}$ the set of Lipschitz maps $M : \mathcal{O} \rightarrow \mathcal{L}_{a, -m}^{\text{qT}}$ with the $|\cdot|_{a, -m}^{\text{qT}, \mathcal{O}}$ norm.

$$|M|_{a, -m}^{\text{qT}, \mathcal{O}} := \sup_{\omega \in \mathcal{O}} |M(\omega)|_{a, -m}^{\text{qT}} + \gamma \sup_{\omega \neq \omega' \in \mathcal{O}} |\Delta_{\omega, \omega'} M|_{a, -m}^{\text{qT}}$$

Given a momentum preserving operator $M \in \mathcal{L}_a(\mathfrak{H}^p)$ and $K > 0$, we define the projections $\Pi_{|\ell| \leq K} M, \Pi_{|\ell| > K} M$ by

$$\Pi_{|\ell| \leq K} M = \begin{cases} M_j^{j - \pi(\ell)}(\ell) & \text{if } |\ell| \leq K \\ 0 & \text{otherwise} \end{cases}, \quad \Pi_{|\ell| > K} M := M - \Pi_{|\ell| \leq K} M. \quad (2.17)$$

These projections obviously map the space $\mathcal{L}_a(\mathfrak{H}^p)$ to itself. Moreover, since the symbol $\mathfrak{M}(\ell, v(j), b(j))$ of a line-Töplitz operator has no conditions on ℓ , they are also well-behaved with respect to the line-Töplitz structure. By direct inspection one then gets the following Lemmata.

Lemma 2.19. The projections $\Pi_{|\ell| \leq K}$ (and $\Pi_{|\ell| > K}$) preserve the spaces $\mathcal{T}_{a, m}^{\mathcal{O}}, \mathcal{L}_{a, -m}^{\text{qT}, \mathcal{O}}$, and they are continuous. In particular, if M is quasi-Töplitz then so is its time average $\langle M \rangle_{\mathbb{T}^d} := \Pi_{\ell=0} M$. For all $a' < a$, one has the bounds:

$$\begin{aligned} |\Pi_{|\ell| \leq K} M|_{a, -m}^{\text{T}, \mathcal{O}} &\leq |M|_{a, -m}^{\text{T}, \mathcal{O}}, & |\Pi_{|\ell| \leq K} M|_{a, -m}^{\text{qT}, \mathcal{O}} &\leq |M|_{a, -m}^{\text{qT}, \mathcal{O}}, \\ |\Pi_{|\ell| > K} M|_{a', -m}^{\text{T}, \mathcal{O}} &\leq e^{-(a-a')K} |M|_{a, -m}^{\text{T}, \mathcal{O}}, & |\Pi_{|\ell| > K} M|_{a', -m}^{\text{qT}, \mathcal{O}} &\leq e^{-\frac{a-a'}{2}K} |M|_{a, -m}^{\text{qT}, \mathcal{O}}. \end{aligned}$$

Lemma 2.20. Given a time independent (and hence diagonal) operator $A = \text{diag}(A_j) \in \mathcal{L}_{a, -m}^{\text{qT}, \mathcal{O}}$, there exists a decomposition of the eigenvalues

$$A_j = \mathfrak{a}(v(j), b(j)) + \sum_{k=1}^m r_j^{(k)}$$

so that

$$\sup_{v \in \mathcal{V}, b \in \mathbb{Z}} |\mathfrak{a}(v, b)|^{\mathcal{O}} e^{ac|v|} \langle b \rangle^m + \sum_{k=1}^m \sup_j |r_j^{(k)}|^{\mathcal{O}} \langle j \rangle^{k\mu} \langle b(j) \rangle^{m-k} \leq 2|A|_{a, -m}^{\text{qT}, \mathcal{O}} \quad (2.18)$$

Moreover, if $A_j \in \mathbb{R}$, then also $\mathfrak{a}, r_j^{(k)} \in \mathbb{R}$ for $k = 1, \dots, m$.

Proof. Since $A \in \mathcal{L}_{a, -m}^{\text{qT}, \mathcal{O}}$ there exists a decomposition

$$A = A^{\text{T}} + \sum_{k=1}^m A^{(k)}, \quad \text{such that} \quad |A^{\text{T}}|_{a, -m}^{\text{T}, \mathcal{O}} + \sum_{i=1}^m |A^{(i)}|_{\frac{a}{2}; -(i, m-i)}^{\mathcal{O}} \leq 2|A|_{a, -m}^{\text{qT}, \mathcal{O}}.$$

Recalling that A is time-independent, we have

$$A = \langle A \rangle_{\mathbb{T}^d} = \langle A^{\text{T}} \rangle_{\mathbb{T}^d} + \sum_{k=1}^m \langle A^{(k)} \rangle_{\mathbb{T}^d}.$$

Now, the operators $\langle A^T \rangle_{\mathbb{T}^d}, \langle A^{(k)} \rangle_{\mathbb{T}^d}$ are all time-independent and hence diagonal; moreover, $\langle A^T \rangle_{\mathbb{T}^d}$ is still line Töplitz. The bounds follow from Remark 2.8. The reality condition follows by taking the real part in both sides of the equality above. \square

We now discuss the algebra properties of $\mathcal{L}_{a,-m}^{\text{qT},\mathcal{O}}$. Our purpose is to get a result analogous to the one in Lemma 2.14. With this in mind, we give two propositions on the product of operators in $\mathcal{L}_{a,-m}^{\text{qT},\mathcal{O}}$.

Proposition 2.21. *Let $M_1 \in \mathcal{L}_{a,-m_1}^{\text{qT},\mathcal{O}}$ and $M_2 \in \mathcal{L}_{a,-m_2}^{\text{qT},\mathcal{O}}$, and set $m = \min\{m_1, m_2\}$. Then we have $M_1 M_2 \in \mathcal{L}_{a,-m}^{\text{qT},\mathcal{O}}$, with the bound*

$$|M_1 M_2|_{a,-m}^{\text{qT},\mathcal{O}} \leq C a^{-q_0} |M_1|_{a,-m_1}^{\text{qT},\mathcal{O}} |M_2|_{a,-m_2}^{\text{qT},\mathcal{O}}. \quad (2.19)$$

Moreover, setting $m_1 = m_2 = 1$, one also has $\forall 0 < a' < a$ that $M_1 M_2 \in \mathcal{L}_{a',-2}^{\text{qT},\mathcal{O}}$, with

$$|M_1 M_2|_{a',-2}^{\text{qT},\mathcal{O}} \leq C (a - a')^{-q_1} |M_1|_{a,-1}^{\text{qT},\mathcal{O}} |M_2|_{a,-1}^{\text{qT},\mathcal{O}}. \quad (2.20)$$

Here C, q_0, q_1 are positive constants depending only on $m_1, m_2, \mu, \delta, \mathbf{c}, p$.

The proof, being quite technical, is postponed to Appendix B.

Corollary 2.22 (Exponentials). *Given $a, m > 0$ and $A \in \mathcal{L}_{a,-m}^{\text{qT},\mathcal{O}}$ such that (the constants C, γ are defined in Proposition 2.21)*

$$|A|_{a,-m}^{\text{qT},\mathcal{O}} \leq \delta := \frac{a^{q_0}}{4C} \quad (2.21)$$

one has for any sequence $c_k \in \ell_\infty$, and for all $M \in \mathcal{L}_{a,-m}^{\text{qT},\mathcal{O}}$

$$\left| \sum_{k=d}^{\infty} c_k \text{ad}(A)^k M \right|_{a,-m}^{\text{qT},\mathcal{O}} \leq 2|c|_\infty \left(\frac{|A|_{a,-m}^{\text{qT},\mathcal{O}}}{2\delta} \right)^d \quad (2.22)$$

Proof. It follows directly by iterating the bounds in Proposition 2.21. \square

Remark 2.23. *Estimate (2.19) of Proposition 2.21 and Corollary 2.22 ensure that the well known fact that smoothing operators generate bounded changes of variables holds also in the quasi-Töplitz context. On the other hand, estimate (2.20) shows that the product of two operators of order -1 is actually of order -2 , up to a small loss of analyticity.*

3 The homological equation

We consider a diagonal operator

$$D = \text{diag}(\Omega_j)_{j \in \mathbb{Z}^2}, \quad \Omega_j = |j|^2 + \tilde{\Omega}_j, \quad \tilde{\Omega} := \text{diag}(\tilde{\Omega}_j) \in \mathcal{L}_{a,-2}^{\text{qT},\mathbf{R}}. \quad (3.1)$$

Recalling Lemma 2.20, we have $\tilde{\Omega}_j = \mathbf{a}(v(j), b(j)) + r_j^{(1)} + r_j^{(2)}$, with $\mathbf{a}(v, b), r^{(1)}, r^{(2)}$ satisfying (2.18) for $m = 2$. For all $0 < |\ell| \leq K$, all $j \in \mathbb{Z}^2$ and all $v \in \mathcal{V}$ such that $\pi(\ell) \parallel v$ we define

$$\mathbf{d}(\ell, j) := \omega \cdot \ell + \Omega_j - \Omega_{j - \pi(\ell)} \quad (3.2)$$

$$\mathfrak{d}(\ell, v, b) := \omega \cdot \ell + 2 \frac{|\pi(\ell)|}{|v|} b - |\pi(\ell)|^2 + \mathbf{a}(v, b) - \mathbf{a}(v, b - v \cdot \pi(\ell)) \quad (3.3)$$

We define the (possibly empty) sets for $\gamma \geq 0$, $K \in \mathbb{N}$

$$\mathcal{C}^{(1)} \equiv \mathcal{C}_{D,K}^{(1)}(\gamma) := \{\omega \in \mathcal{O} : |\mathfrak{d}(\ell, j)| \geq \gamma |\ell|^{-\tau}, \forall 0 < |\ell| \leq K, j \in \mathbb{Z}^2\} \quad (3.4)$$

$$\mathcal{C}^{(2)} \equiv \mathcal{C}_{D,K}^{(2)}(\gamma) := \{\omega \in \mathcal{O} : |\mathfrak{d}(\ell, v, b)| \geq 2\gamma |\ell|^{-\tau}, \forall 0 < |\ell| \leq K, v \in \mathcal{V} \text{ s.t. } \pi(\ell) \parallel v\}, \quad (3.5)$$

$$\mathcal{O}_{D,K}(\gamma) := \mathcal{C}_{D,K}^{(1)}(\gamma) \cap \mathcal{C}_{D,K}^{(2)}(\gamma). \quad (3.6)$$

Proposition 3.1. Fix $\gamma, a > 0$, $K \in \mathbb{N}$ and a compact set $\mathcal{O} \subseteq \mathcal{O}_0 \subset \mathbb{R}$. Consider a diagonal operator D as in (3.1) with $16|\tilde{\Omega}|_{a,-2}^{\mathfrak{qT},\mathbb{R}} \leq \gamma$. Consider the sets $\mathcal{C}^{(1)}$, $\mathcal{C}^{(2)}$ defined in (3.4), (3.5) and set

$$\mathcal{O}_+ = \mathcal{O}_{D,K}(\gamma). \quad (3.7)$$

For all $P \in \mathcal{L}_{a,-2}^{\mathfrak{qT},\mathcal{O}}$, there exists $S \in \mathcal{L}_{a,-2}^{\mathfrak{qT},\mathcal{O}_+}$ solving the homological equation

$$-i\dot{S} + [D, S] = \Pi_{0 < |\ell| \leq K} P, \quad (3.8)$$

fulfilling the estimate

$$|S|_{a,-2}^{\mathfrak{qT},\mathcal{O}_+} \leq C(p, \mu, \delta, \mathfrak{c}) a^{-2/\delta} \gamma^{-1} K^{3\tau+1} |P|_{a,-2}^{\mathfrak{qT},\mathcal{O}}. \quad (3.9)$$

If P is self-adjoint, then so is iS ; if P is Gauge covariant, then so is S .

The solution S of the homological equation (3.8) is given, in components, by

$$S_j^{j-\pi(\ell)}(\ell) = \frac{P_j^{j-\pi(\ell)}(\ell)}{\omega \cdot \ell + \Omega_j - \Omega_{j-\pi(\ell)}} \quad \text{if } 0 < |\ell| \leq K, \quad S_j^{j-\pi(\ell)}(\ell) = 0 \quad \text{otherwise.} \quad (3.10)$$

Note that S is well defined for $\omega \in \mathcal{C}_D^{(1)}$ and it is well known that $S \in \mathcal{L}_a(\mathfrak{H}^p)$. Moreover if P is self-adjoint, one verifies that iS is so (just use the characterization (2.5)), same for the Gauge covariance.

Next we show that $S \in \mathcal{L}_{a,-2}^{\mathfrak{qT},\mathcal{O}_+}$, namely it has a decomposition (cfr. (2.16))

$$S = S^{\mathfrak{T}} + S^{(1)} + S^{(2)} \quad \text{with } S^{\mathfrak{T}} \in \mathcal{T}_{a,-2}^{\mathcal{O}_+}, \quad S^{(1)} \in \mathcal{L}_{\frac{\mathfrak{a}}{2};-(1,1)}^{\mathcal{O}_+}, \quad S^{(2)} \in \mathcal{L}_{\frac{\mathfrak{a}}{2};-(2,0)}^{\mathcal{O}_+}.$$

By assumption also P is admissible of order -2 , hence it has the same decomposition. In particular we shall denote the line-Töplitz part of P by

$$[P^{\mathfrak{T}}]_j^{j-\pi(\ell)} = \mathfrak{P}(\ell, v(j), b(j)). \quad (3.11)$$

We then decompose S as follows

$$S_j^{j-\pi(\ell)}(\ell) = \frac{[P^{\mathfrak{T}}]_j^{j-\pi(\ell)}(\ell)}{\omega \cdot \ell + \Omega_j - \Omega_{j-\pi(\ell)}} + \underbrace{\frac{[P^{(1)}]_j^{j-\pi(\ell)}(\ell)}{\omega \cdot \ell + \Omega_j - \Omega_{j-\pi(\ell)}}}_{=:[\tilde{S}^{(1)}]_j^{j-\pi(\ell)}} + \underbrace{\frac{[P^{(2)}]_j^{j-\pi(\ell)}(\ell)}{\omega \cdot \ell + \Omega_j - \Omega_{j-\pi(\ell)}}}_{=:[\tilde{S}^{(2)}]_j^{j-\pi(\ell)}}. \quad (3.12)$$

We further need to decompose the first term in the r.h.s. above, which is the most delicate one because of the divisor. We introduce the following sets

$$\begin{aligned} A_0 &:= \{(\ell, j) \in \mathbb{Z}^d \setminus \{0\} \times \mathbb{Z}^2 : \pi(\ell) = 0, \quad |\ell| < K\} \\ A_1 &:= \{(\ell, j) \in \mathbb{Z}^d \setminus \{0\} \times \mathbb{Z}^2 : \pi(\ell) \neq 0, \quad |\ell| < K, \quad v(j) \parallel \pi(\ell)\} \\ A_2 &:= \{(\ell, j) \in \mathbb{Z}^d \setminus \{0\} \times \mathbb{Z}^2 : \pi(\ell) \neq 0, \quad |\ell| < K, \quad v(j) \not\parallel \pi(\ell), \quad |\ell| < \mathfrak{c} \langle j \rangle^\delta\} \\ A_3 &:= \{(\ell, j) \in \mathbb{Z}^d \setminus \{0\} \times \mathbb{Z}^2 : \pi(\ell) \neq 0, \quad v(j) \not\parallel \pi(\ell), \quad \mathfrak{c} \langle j \rangle^\delta \leq |\ell| < K\} \end{aligned} \quad (3.13)$$

and note that $\{(\ell, j) : 0 < |\ell| < K\} = \cup_{i=0}^3 A_i$. Further denote by $\delta_A := \delta_A(\ell, j) = 1$ if $(\ell, j) \in A$, $\delta_A(\ell, j) = 0$ otherwise. Starting from (3.12) we further decompose

$$S_j^{j-\pi(\ell)}(\ell) = \frac{[P^\mathbb{T}]_j^{j-\pi(\ell)}(\ell)}{\mathfrak{d}(\ell, j)} \delta_{A_0} + \frac{[P^\mathbb{T}]_j^{j-\pi(\ell)}(\ell)}{\mathfrak{d}(\ell, v(j), b(j))} \delta_{A_1} \quad (3.14)$$

$$+ [P^\mathbb{T}]_j^{j-\pi(\ell)}(\ell) \left(\frac{1}{\mathfrak{d}(\ell, j)} - \frac{1}{\mathfrak{d}(\ell, v(j), b(j))} \right) \delta_{A_1} + \frac{[P^\mathbb{T}]_j^{j-\pi(\ell)}(\ell)}{\mathfrak{d}(\ell, j)} \delta_{A_2} + [\tilde{S}^{(1)}]_j^{j-\pi(\ell)} \quad (3.15)$$

$$+ \frac{[P^\mathbb{T}]_j^{j-\pi(\ell)}(\ell)}{\mathfrak{d}(\ell, j)} \delta_{A_3} + [\tilde{S}^{(2)}]_j^{j-\pi(\ell)}. \quad (3.16)$$

In the next lemmata we shall prove the line (3.14) gives a line-Töplitz operator, line (3.15) gives an operator in $\mathcal{L}_{\frac{a}{2};-(1,1)}^{\mathcal{O}_+}$ and line (3.16) one in $\mathcal{L}_{\frac{a}{2};-(2,0)}^{\mathcal{O}_+}$.

Lemma 3.2. *The operator in the r.h.s. of line (3.14), which we denote by $S^\mathbb{T}$, is line-Töplitz and*

$$|S^\mathbb{T}|_{a,-2}^{\mathbb{T}, \mathcal{O}_+} \leq \frac{1}{2\gamma} K^{2\tau+1} |P^\mathbb{T}|_{a,-2}^{\mathbb{T}, \mathcal{O}}. \quad (3.17)$$

Proof. Recalling (3.11), (3.3), one checks that the elements of $S^\mathbb{T}$ are actually given by $(S^\mathbb{T})_j^{j-\pi(\ell)}(\ell) := \mathfrak{S}(\ell, v(j), b(j))$ with

$$\mathfrak{S}(\ell, v, b) := \begin{cases} \frac{\mathfrak{P}(\ell, v, b)}{\mathfrak{d}(\ell, v, b)} & \text{if } \pi(\ell) = 0, \quad \ell \neq 0, \quad |\ell| < K \\ \frac{\mathfrak{P}(\ell, v, b)}{\mathfrak{d}(\ell, v, b)} & \text{if } \pi(\ell) \neq 0, \quad |\ell| < K, \quad v \parallel \pi(\ell) \\ 0 & \text{otherwise} \end{cases}$$

so $S^\mathbb{T}$ is line-Töplitz. To study the Lipschitz variation we remark that, since

$$|\Delta_{\omega, \omega'} \mathfrak{d}(\ell, v, b)| \leq |\ell| + 2|\tilde{\Omega}|_{a,-2}^{\mathbb{q}\mathbb{T}, \mathcal{O}} \leq 2K \quad \forall \omega \neq \omega' \in \mathcal{O}_+,$$

one has

$$\left| \Delta_{\omega, \omega'} \frac{1}{\mathfrak{d}(\ell, v, b)} \right| = \left| \frac{\Delta_{\omega, \omega'} \mathfrak{d}(\ell, v, b)}{\mathfrak{d}(\ell, v, b)(\omega) \mathfrak{d}(\ell, v, b)(\omega')} \right| \leq 2\gamma^{-2} K^{2\tau+1}.$$

It follows that $|\Delta_{\omega, \omega'} S^\mathbb{T}|_{a,-2}^{\mathbb{T}} \leq \gamma^{-1} K^\tau |\Delta_{\omega, \omega'} P^\mathbb{T}|_{a,-2}^{\mathbb{T}} + 2\gamma^{-2} K^{2\tau+1} |P^\mathbb{T}|_{a,-2}^{\mathbb{T}}$. Estimate (3.17) follows using the small divisor estimates in (3.4), (3.5). \square

Lemma 3.3. *The operator in line (3.15), which we denote by $S^{(1)}$, belongs to $\mathcal{L}_{\frac{a}{2};-(1,1)}^{\mathcal{O}_+}$ and there exists $C = C(p, \mu, \delta, \mathfrak{c}) > 0$ such that*

$$|S^{(1)}|_{a/2;-(1,1)}^{\mathcal{O}_+} \leq C \gamma^{-1} K^{3\tau+1} |P|_{a,-2}^{\mathbb{q}\mathbb{T}, \mathcal{O}}. \quad (3.18)$$

Proof. We bound first the operator $\tilde{S}^{(1)}$, defined in (3.12). Using $\omega \in \mathcal{C}_D^{(1)}$, we know that $|(\tilde{S}^{(1)})_j^{j-\pi(\ell)}(\ell)| \leq \gamma^{-1} K^\tau |(P^{(1)})_j^{j-\pi(\ell)}(\ell)|$. Thus, by Remark 2.8, we get $|\tilde{S}^{(1)}|_{a/2;-(1,1)} \leq \gamma^{-1} K^\tau |P^{(1)}|_{a/2;-(1,1)}$. To study the Lipschitz norm we proceed as in the previous Lemma and we conclude that

$$|\tilde{S}^{(1)}|_{a/2;-(1,1)}^{\mathcal{O}_+} \leq 2\gamma^{-1} K^{2\tau+1} |P^{(1)}|_{a/2;-(1,1)}^{\mathcal{O}}. \quad (3.19)$$

Regarding the remaining operator in line (3.15), which we denote by $\widehat{S}^{(1)}$, it has matrix elements given explicitly by

$$(\widehat{S}^{(1)})_j^{j-\pi(\ell)}(\ell) := \begin{cases} \frac{(P^\mathbb{T})_j^{j-\pi(\ell)}(\ell)}{\mathfrak{d}(\ell, j)}, & \text{if } \pi(\ell) \neq 0, \quad |\ell| < \min(K, \mathfrak{c}\langle j \rangle^\delta), \quad v(j) \not\parallel \pi(\ell) \\ (P^\mathbb{T})_j^{j-\pi(\ell)}(\ell) \left(\frac{1}{\mathfrak{d}(\ell, j)} - \frac{1}{\mathfrak{d}(\ell, v(j), b(j))} \right), & \text{if } \pi(\ell) \neq 0, \quad |\ell| < K, \quad v(j) \parallel \pi(\ell) \\ 0 & \text{otherwise.} \end{cases}$$

First line of $\widehat{S}^{(1)}$: We claim that there exists $C = C(\delta, \mu, \gamma)$ such that

$$\sup_{\substack{j \in \mathbb{Z}^2, \ell \in \mathbb{Z}^d: \\ |\ell| < \mathfrak{c}\langle j \rangle^\delta, \pi(\ell) \not\parallel v(j)}} \left| \frac{\langle j \rangle^\mu}{\omega \cdot \ell + \Omega_j - \Omega_{j-\pi(\ell)}} \right|^{\mathcal{O}^+} \leq C\gamma^{-1}K^{2\tau+1}. \quad (3.20)$$

Indeed if $\langle j \rangle \leq J_\delta$ (with J_δ the constant of Lemma 2.12), then the bound follows trivially using also that $\omega \in \mathcal{C}^{(1)}$.

Consider now the case $\langle j \rangle > J_\delta$. Then by (3.3), in order for the denominator to gain the term $\langle j \rangle^\mu$, we need $\pi(\ell) \cdot j$ to be large. This is what we show to happen. In particular we claim that

$$\langle j \rangle \geq J_\delta, \quad |\ell| < \mathfrak{c}\langle j \rangle^\delta, \quad \pi(\ell) \not\parallel v(j) \quad \Rightarrow \quad |j \cdot \pi(\ell)| > 2\langle j \rangle^\mu \quad (3.21)$$

and then (3.20) still holds. To prove (3.21) we distinguish 2 cases.

Case 1: $|b(j)| > 2\langle j \rangle^\mu$. Let $w \in \mathcal{V}$ be the direction parallel to $\pi(\ell)$, then (being w a generator), $|w| \leq |\pi(\ell)| \leq \mathfrak{c}^{-1}|\ell| \leq \langle j \rangle^\delta$. As $v(j)$ is the vector realizing $\min\{|v \cdot j| : v \in \mathcal{V}, |v| \leq |j|^\delta\}$, we have

$$|\pi(\ell) \cdot j| \geq |w \cdot j| \geq |v(j) \cdot j| = |b(j)| > 2\langle j \rangle^\mu$$

and in this case (3.21) follows.

Case 2: $|b(j)| \leq 2\langle j \rangle^\mu$. Then as $|w| \leq \langle j \rangle^\delta$ and $w \neq v(j)$, Lemma 2.12 (i) gives again $|w \cdot j| \geq 2\langle j \rangle^\mu$, proving (3.21) also in this case.

Now, recalling that

$$\mathfrak{d}(\ell, j) = \omega \cdot \ell + 2\pi(\ell) \cdot j - |\pi(\ell)|^2 + \widetilde{\Omega}_j - \widetilde{\Omega}_{j-\pi(\ell)}, \quad (3.22)$$

and using (3.21) we obtain that $|\mathfrak{d}(\ell, j)| \geq 2\langle j \rangle^\mu - (|\omega| \mathfrak{c} + 1)\langle j \rangle^{2\delta} - 1 \geq C\langle j \rangle^\mu$. Hence (3.20) is proved in case of the sup norm. The Lipschitz norm is estimated by the same case analysis.

We deduce that

$$|\widehat{S}^{(1)}\delta_{A_2}|_{a/2; -(1,1)}^{\mathcal{O}^+} \leq C\gamma^{-1}K^{2\tau+1}|P^\mathbb{T}|_{a/2; -(0,2)}^{\mathcal{O}} \leq \left(\frac{a}{2}\right)^{-p-2} C\gamma^{-1}K^{2\tau+1}|P^\mathbb{T}|_{a, -2}^{\mathbb{T}, \mathcal{O}}, \quad (3.23)$$

where in the last inequality we used Lemma 2.17.

Second line of $\widehat{S}^{(1)}$: Again we consider two cases.

Case 1: $|b(j)| > 2\langle j \rangle^\mu$ or $\langle v(j) \rangle > \frac{1}{2}\langle j \rangle^\delta$ or $|\ell| > \mathfrak{c}\langle j \rangle^\delta$. In this case we exploit the decay of $(P^\mathbb{T})_j^{j-\pi(\ell)}$ and get that for $\omega \in \mathcal{O}_{D, K}(\gamma)$

$$\begin{aligned} & |b(j)| \langle j \rangle^\mu \left| (P^\mathbb{T})_j^{j-\pi(\ell)}(\ell) \left(\frac{1}{\mathfrak{d}(\ell, j)} - \frac{1}{\mathfrak{d}(\ell, v(j), b(j))} \right) \right| \\ & \leq |b(j)| \langle j \rangle^\mu \left| (P^\mathbb{T})_j^{j-\pi(\ell)}(\ell) \right| \left(\left| \frac{1}{\mathfrak{d}(\ell, j)} \right| + \left| \frac{1}{\mathfrak{d}(\ell, v(j), b(j))} \right| \right) \\ & \leq C a^{-\frac{\mu}{\delta}} \gamma^{-1} K^\tau e^{-\frac{\mathfrak{a}\mathfrak{c}}{2}|v(j)| - \frac{3\mathfrak{a}}{4}|\ell|} |P^\mathbb{T}|_{a, -2}^{\mathbb{T}, \mathcal{O}}. \end{aligned} \quad (3.24)$$

Case 2: $|b(j)| \leq 2\langle j \rangle^\mu$ and $\langle v(j) \rangle \leq \frac{1}{2}\langle j \rangle^\delta$ and $|\ell| \leq c\langle j \rangle^\delta$. In this case we exploit the decay of $\mathfrak{d}(\ell, j)^{-1} - \mathfrak{d}(\ell, v(j), b(j))^{-1}$. Precisely we note that

$$\mathfrak{d}(\ell, j) := \omega \cdot \ell + |j|^2 - |j - \pi(\ell)|^2 + \tilde{\Omega}_j - \tilde{\Omega}_{j-\pi(\ell)} = \omega \cdot \ell + 2\pi(\ell) \cdot j - |\pi(\ell)|^2 + \tilde{\Omega}_j - \tilde{\Omega}_{j-\pi(\ell)},$$

so decomposing $\tilde{\Omega}$ as in Lemma 2.20, and using that $v(j) \parallel \pi(\ell)$ we arrive at

$$\begin{aligned} \mathfrak{d}(\ell, j) &= \omega \cdot \ell + 2 \frac{|\pi(\ell)|}{|v(j)|} b(j) - |\pi(\ell)|^2 + \mathfrak{a}(v(j), b(j)) - \mathfrak{a}(v(j - \pi(\ell)), b(j - \pi(\ell))) \\ &\quad + r_j^{(1)} - r_{j-\pi(\ell)}^{(1)} + r_j^{(2)} - r_{j-\pi(\ell)}^{(2)}. \end{aligned}$$

Since $|b(j)| \leq 2\langle j \rangle^\mu$, $\langle v(j) \rangle \leq \frac{1}{2}\langle j \rangle^\delta$ and also $|\pi(\ell)| \leq \langle j \rangle^\delta$, Lemma 2.12 (ii) implies that $v(j) = v(j - \pi(\ell))$ and then one has also $b(j - \pi(\ell)) = b(j) - v(j) \cdot \pi(\ell)$. We deduce that

$$\mathfrak{d}(\ell, j) = \mathfrak{d}(\ell, v(j), b(j)) + r_j^{(1)} - r_{j-\pi(\ell)}^{(1)} + r_j^{(2)} - r_{j-\pi(\ell)}^{(2)}$$

and thus we estimate

$$\begin{aligned} |\langle j \rangle^\mu \left(\frac{1}{\mathfrak{d}(\ell, j)} - \frac{1}{\mathfrak{d}(\ell, v(j), b(j))} \right)| &= \left| \frac{\langle j \rangle^\mu (r_j^{(1)} - r_{j-\pi(\ell)}^{(1)} + r_j^{(2)} - r_{j-\pi(\ell)}^{(2)})}{\mathfrak{d}(\ell, j) \mathfrak{d}(\ell, v(j), b(j))} \right| \\ &\leq |\tilde{\Omega}|_{a, -2}^{\mathfrak{qT}, \mathcal{O}_0} \gamma^{-2} K^{2\tau} \leq \gamma^{-1} K^{2\tau} \end{aligned}$$

We reason in the same way for the Lipschitz variation and obtain the bound

$$\sup_{\substack{j \in \mathbb{Z}^2, \ell \in \mathbb{Z}^d: \\ |\ell| < K, \quad \pi(\ell) \parallel v(j)}} \left| \langle j \rangle^\mu \left(\frac{1}{\mathfrak{d}(\ell, j)} - \frac{1}{\mathfrak{d}(\ell, v(j), b(j))} \right) \right|^{\mathcal{O}^+} \leq C \gamma^{-1} K^{3\tau+1}. \quad (3.25)$$

Thus we obtain the same kind of estimate as in (3.24), with $\tau \rightsquigarrow 3\tau + 1$.

In both cases, using Lemma 2.9, we finally get

$$|\widehat{S}^{(1)} \delta_{A_1}|_{a/2; -(1,1)}^{\mathcal{O}^+} \leq C \gamma^{-1} K^{3\tau+1} |P^{\mathfrak{T}}|_{a, -2}^{\mathfrak{T}, \mathcal{O}}. \quad (3.26)$$

Then estimate (3.18) follows from (3.19), (3.23), (3.26). \square

Lemma 3.4. *The operator in line (3.16), which we denote by $S^{(2)}$, belongs to $\mathcal{L}_{\frac{a}{2}; -(2,0)}^{\mathcal{O}^+}$ and there exists $C = C(p, \mu, \delta, \mathfrak{c}) > 0$ such that*

$$|S^{(2)}|_{a/2; -(2,0)}^{\mathcal{O}^+} \leq C \gamma^{-1} K^{2\tau+1} \left(|P^{(2)}|_{a/2; -(2,0)}^{\mathcal{O}} + a^{-2\mu/\delta} |P^{\mathfrak{T}}|_a^{\mathcal{O}} \right) \quad (3.27)$$

Proof. We need to estimate two contributions. The operator $\tilde{S}^{(2)}$, given in (3.12), is estimated similarly as $\tilde{S}^{(1)}$ in the previous lemma, obtaining the estimate

$$|\tilde{S}^{(2)}|_{a/2; -(2,0)}^{\mathcal{O}^+} \leq \gamma^{-1} K^{2\tau+1} |P^{(2)}|_{a/2; -(2,0)}^{\mathcal{O}}. \quad (3.28)$$

Regarding the first operator in (3.16), which we denote by $\widehat{S}^{(2)}$, it has elements only supported in the region $|\ell| \geq c\langle j \rangle^\delta$. Then the small divisor estimate in (3.4) and Lemma A.1 give the bound

$$|\widehat{S}^{(2)}|_{a/2; -(2,0)}^{\mathcal{O}^+} \leq C(p, \mu, \delta, \mathfrak{c}) a^{-2\mu/\delta} |\widehat{S}^{(2)}|_a^{\mathcal{O}^+} \leq \gamma^{-1} C(p, \delta) a^{-2\mu/\delta} K^{2\tau+1} |P^{\mathfrak{T}}|_a^{\mathcal{O}}. \quad \square$$

Proof of Proposition 3.1. Estimate (3.9) follows from Lemmata 3.2, 3.3 and 3.4. \square

4 The KAM reduction scheme

In this section we shall study the linear Schrödinger operator in (1.21) and prove that, provided that V in (1.16) is sufficiently small, then \mathfrak{L} is reducible with sufficient knowledge on the asymptotics of the eigenvalues. As explained in the introduction, the reducibility of Schrödinger operators is by now well-known, the novelties in this result are the fact that the reducing change of variables is quasi-Töplitz of order -1 and the asymptotics of the eigenvalues.

Theorem 4.1 (Reducibility). *Consider a Schrödinger operator as in (1.21) with a travelling wave potential \mathcal{V} as in (1.16). There exist $\epsilon_*, \gamma_* > 0$ such that the following holds. For all $\gamma \in (0, \gamma_*)$ and for any potential satisfying the smallness condition (1.19) there exist: a sequence of Lipschitz functions $\mathbf{R} := [-1, 1]^d \ni \omega \mapsto \Omega_j(\omega)$, a positive measure set $\mathcal{O}_\infty \subset \mathbf{R}$ and a symplectic change of variables G such that*

$$G_* \mathfrak{L} = \text{diag}(\Omega_j), \quad G - \text{Id} \in \mathcal{L}_{\mathbf{a}', -1}^{\text{qT}, \mathcal{O}_\infty}, \quad \text{with } \mathbf{a}' = \mathbf{a}/8. \quad (4.1)$$

Moreover

$$\Omega_j = |j|^2 + \mathbf{a}(v(j), b(j)) + r_j^{(1)} + r_j^{(2)} \quad (4.2)$$

with the bounds

$$\sup_{v \in \mathcal{V}, b \in \mathbb{Z}} |\mathbf{a}(v, b)|^{\mathbf{R}} e^{\mathbf{a}'c|v|} \langle b \rangle^2 + \sup_j |r_j^{(1)}|^{\mathbf{R}} \langle j \rangle^\mu \langle b(j) \rangle + \sup_j |r_j^{(2)}|^{\mathbf{R}} \langle j \rangle^{2\mu} \leq 2 \|V\|_{\mathbf{a}, \mathbf{p}}.$$

Finally if V is Gauge invariant then G is Gauge covariant.

We can be more explicit regarding the set \mathcal{O}_∞ . For $\ell \in \mathbb{Z}^d$, $j \in \mathbb{Z}^2$ and all $v \in \mathcal{V}$ such that $\pi(\ell) \parallel v$ we consider $\mathbf{d}(\ell, j), \mathfrak{d}(\ell, v, b)$ defined as in (3.2) with Ω_j defined in (4.2).

Proposition 4.2. *Under the Hypotheses of Theorem 4.1, there exists $\tau > 0$ such that the set \mathcal{O}_∞ contains the set $\mathcal{O}_{\text{fin}} := \mathcal{O}_0(\gamma) \cap \mathcal{C}^{(1)}(\gamma, \tau) \cap \mathcal{C}^{(2)}(\gamma, \tau)$ where*

$$\mathcal{O}_0 = \mathcal{O}_0(\gamma) := \{\omega \in \mathbf{R} : |\omega \cdot \ell + k| \geq 2\gamma |\ell|^{-(d+1)} \quad \forall (\ell, k) \in \mathbb{Z}^{d+1} : \ell \neq 0\}, \quad (4.3)$$

$$\mathcal{C}^{(1)} = \mathcal{C}^{(1)}(\gamma, \tau) := \{\omega \in \mathbf{R} : |\mathbf{d}(\ell, j)| \geq 2\gamma |\ell|^{-\tau}, \quad \forall \ell \neq 0, \quad j \in \mathbb{Z}^2\}, \quad (4.4)$$

$$\mathcal{C}^{(2)} := \mathcal{C}^{(2)}(\gamma, \tau) := \{\omega \in \mathbf{R} : |\mathfrak{d}(\ell, v, b)| \geq 4\gamma |\ell|^{-\tau}, \quad \forall \ell \neq 0, \quad v \in \mathcal{V} \text{ s.t. } \pi(\ell) \parallel v\}. \quad (4.5)$$

We defer the proof of this proposition to the end of the section.

The first step. In the first step in the reduction scheme we exploit the structure of the Schrödinger operator in (1.21) and perform a change of variables $G_0 = e^{S_0}$ which conjugates \mathfrak{L} to $-\Delta + P_0$, with P_0 quasi-Töplitz of order -2 .

Lemma 4.3. *Fix $a_0 = \mathbf{a}/4$, $\gamma > 0$ and consider a Schrödinger operator as in (1.21) satisfying (1.19). Then there exist positive constants C, q_2, q_4 , depending on $\delta, \mu, \tau, \mathbf{c}, \mathbf{p}$ only, and operators $S_0 \in \mathcal{L}_{\frac{\mathbf{a}}{3}, -1}^{\text{qT}, \mathcal{O}_0}$ and $P_0 \in \mathcal{L}_{a_0, -2}^{\text{qT}, \mathcal{O}_0}$ (with \mathcal{O}_0 defined in (4.3)) satisfying the bounds*

$$|S_0|_{\frac{\mathbf{a}}{3}, -1}^{\text{qT}, \mathcal{O}_0} \leq C \gamma^{-1} \mathbf{a}^{-q_2} \|V\|_{\mathbf{a}, \mathbf{p}}, \quad |P_0|_{a_0, -2}^{\text{qT}, \mathcal{O}_0} \leq C \gamma^{-1} \mathbf{a}^{-q_4} \|V\|_{\mathbf{a}, \mathbf{p}}^2,$$

such that $\mathfrak{L}_0 := (G_0)_* \mathfrak{L} = -\Delta + P_0$.

The proof is performed in several steps. We start by defining S_0 as the solution of the homological equation (see (2.9))

$$i\dot{S}_0 + [\Delta, S_0] = M_V \iff (S_0)_j^{j-\pi(\ell)}(\ell) = \begin{cases} -\frac{V(\ell)}{\omega \cdot \ell + |j|^2 - |j - \pi(\ell)|^2} & \text{if } \pi(\ell) \neq 0 \\ 0 & \text{otherwise} \end{cases} \quad (4.6)$$

(recall that, by (1.17), $V(\ell) = 0$ if $\pi(\ell) = 0$). Note that S_0 is well defined for $\omega \in \mathcal{O}_0$ and

$$(G_0)_*(-\Delta + M_V) = e^{\text{ad}S_0}(-\Delta + M_V) - i \sum_{h=1}^{\infty} \frac{(\text{ad}S_0)^{h-1}}{h!} \dot{S}_0 \quad (4.7)$$

$$= -\Delta + M_V + \sum_{h=1}^{\infty} \frac{(\text{ad}S_0)^{h-1}}{h!} ([-\Delta + M_V, S_0] - i\dot{S}_0) \quad (4.8)$$

$$= -\Delta + [M_V, S_0] + \sum_{h=2}^{\infty} \frac{(\text{ad}S_0)^{h-1}}{h!} ([M_V, S_0] - M_V),$$

so for $\omega \in \mathcal{O}_0$

$$P_0 := [M_V, S_0] + \sum_{h=2}^{\infty} \frac{(\text{ad}S_0)^{h-1}}{h!} ([M_V, S_0] - M_V) = \sum_{h=1}^{\infty} \frac{h(\text{ad}S_0)^h}{(h+1)!} M_V.$$

In Appendix B we prove that there exist positive constants C, q_2, q_3 , depending on δ, μ, τ, c, p only, such that the following holds.

Lemma 4.4. *Let S_0 be the solution of the homological equation (4.6). Then for $2a' < \mathbf{a}$ we have*

$$|S_0|_{a', -1}^{\mathfrak{qT}, \mathcal{O}_0} \leq \frac{C}{\gamma(\mathbf{a} - 2a')^{q_2}} \|V\|_{\mathbf{a}, \mathbf{p}}. \quad (4.9)$$

Moreover the commutator $[S_0, M_V]$ is quasi-Töplitz of order -2 with the bounds

$$|[S_0, M_V]|_{a', -2}^{\mathfrak{qT}, \mathcal{O}_0} \leq C\gamma^{-1}(\mathbf{a} - 2a')^{-q_3} \|V\|_{\mathbf{a}, \mathbf{p}}^2 \quad (4.10)$$

Proof of Lemma 4.3. We need to bound $|(\text{ad}S_0)^h M_V|_{a_0, -2}^{\mathfrak{qT}, \mathcal{O}_0}$ for all $h \geq 1$. If $h = 1$, this is achieved through (4.10) with $a' = a_0$.

If $h = 2$, then by (2.20) of Proposition 2.21 (and recalling that $|\cdot|_{a, -1}^{\mathfrak{qT}, \mathcal{O}_0} \leq |\cdot|_{a, -2}^{\mathfrak{qT}, \mathcal{O}_0}$)

$$|[S_0, [S_0, M_V]]|_{a_0, -2}^{\mathfrak{qT}, \mathcal{O}_0} \leq C\mathbf{a}^{-q_1} |S_0|_{\frac{\mathbf{a}}{3}, -1}^{\mathfrak{qT}, \mathcal{O}_0} |[S_0, M_V]|_{\frac{\mathbf{a}}{3}, -2}^{\mathfrak{qT}, \mathcal{O}_0} \leq C^3 \mathbf{a}^{-q_1 - q_2 - q_3} \gamma^{-2} \|V\|_{\mathbf{a}, \mathbf{p}}^3.$$

Otherwise if $h \geq 3$, then we set $A = (\text{ad}S_0)^{h-2} [S_0, M_V]$ so that, by (2.19) and possibly taking C larger,

$$|A|_{\frac{\mathbf{a}}{3}, -1}^{\mathfrak{qT}, \mathcal{O}_0} \leq (C\mathbf{a}^{-q_0} |S_0|_{\frac{\mathbf{a}}{3}, -1}^{\mathfrak{qT}, \mathcal{O}_0})^{h-2} |[S_0, M_V]|_{\frac{\mathbf{a}}{3}, -2}^{\mathfrak{qT}, \mathcal{O}_0}.$$

Then we apply (2.20), which reads

$$|[S_0, A]|_{a_0, -2}^{\mathfrak{qT}, \mathcal{O}_0} \leq C\mathbf{a}^{-q_1} (C\mathbf{a}^{-q_0} |S_0|_{\frac{\mathbf{a}}{3}, -1}^{\mathfrak{qT}, \mathcal{O}_0})^{h-1} |[S_0, M_V]|_{\frac{\mathbf{a}}{3}, -2}^{\mathfrak{qT}, \mathcal{O}_0} \leq \gamma(C\gamma^{-1}\mathbf{a}^{-\frac{q_4}{2}} \|V\|_{\mathbf{a}, \mathbf{p}})^{h+1}$$

with $q_4 = 2(q_0 + q_1 + q_2 + q_3)$. We deduce that

$$|P_0|_{a_0, -2}^{\mathfrak{qT}, \mathcal{O}_0} \leq \gamma \sum_{h \geq 1} \frac{h(C\gamma^{-1}\mathbf{a}^{-\frac{q_4}{2}} \|V\|_{\mathbf{a}, \mathbf{p}})^{h+1}}{(h+1)!} \leq C'\gamma^{-1}\mathbf{a}^{-q_4} \|V\|_{\mathbf{a}, \mathbf{p}}^2,$$

provided that $C\gamma^{-1}\mathbf{a}^{-q_4} \|V\|_{\mathbf{a}, \mathbf{p}} \leq \frac{1}{2}$. This amounts to fixing $\epsilon_* \leq 2C^{-1}\mathbf{a}^{q_4}$ in (1.19). \square

The iterative scheme. After the first step we have conjugated \mathfrak{L} to an operator of the form $-\Delta + P_0$ where $P_0 \in \mathcal{L}_{a_0, -2}^{\mathfrak{qT}, \mathcal{O}}$ is appropriately small. Now we apply a slight modification of a standard KAM scheme in order to reduce to constant coefficients. As before the main novelty is that through the iteration we are able to control the quasi-Töplitz norm. The KAM reduction scheme is based on the iteration of the following lemma.

Lemma 4.5 (KAM step). *Fix $\gamma > 0$ and $K \in \mathbb{N}$. Consider a Schrödinger operator $D + P$ with D defined in (3.1) and $P \in \mathcal{L}_{a, -2}^{\mathfrak{qT}, \mathcal{O}}$ satisfying*

$$16|\tilde{\Omega}|_{a, -2}^{\mathfrak{qT}, \mathbb{R}} \leq \gamma, \quad 16|P|_{a, -2}^{\mathfrak{qT}, \mathcal{O}} \leq \gamma K^{-3\tau-1}. \quad (4.11)$$

Let \mathcal{O}_+ be the set defined in (3.7). There exists $S \in \mathcal{L}_{a, -2}^{\mathfrak{qT}, \mathcal{O}_+}$ and a time independent operator $\tilde{\Omega}^+ \in \mathcal{L}_{a, -2}^{\mathfrak{qT}, \mathbb{R}}$ such that, setting $G = e^S$, for all $\omega \in \mathcal{O}_+$ one has

$$G_*(D + P) = D_+ + P_+, \quad D_+ = \text{diag}(|j|^2 + \tilde{\Omega}_j^+)_{j \in \mathbb{Z}^2}$$

where $P_+ \in \mathcal{L}_{a', -2}^{\mathfrak{qT}, \mathcal{O}_+}$ for all $0 < a' < a$ with the estimate

$$|P_+|_{a', -2}^{\mathfrak{qT}, \mathcal{O}_+} \leq C(p, \delta) a^{-2/\delta} \gamma^{-1} K^{3\tau+1} (|P|_{a, -2}^{\mathfrak{qT}, \mathcal{O}})^2 + e^{-\frac{a-a'}{2}K} |P|_{a, -2}^{\mathfrak{qT}, \mathcal{O}}, \quad (4.12)$$

and

$$|\tilde{\Omega} - \tilde{\Omega}^+|_{a, -2}^{\mathfrak{qT}, \mathbb{R}} \leq |P|_{a, -2}^{\mathfrak{qT}, \mathcal{O}}. \quad (4.13)$$

Proof. We start by defining S as the solution of the homological equation

$$-i\dot{S} + [D, S] = -\Pi_{0 < |\ell| \leq K} P \iff S_j^{j-\pi(\ell)}(\ell) = \begin{cases} -\frac{P_j^{j-\pi(\ell)}(\ell)}{\mathfrak{d}(\ell, j)} & \text{if } 0 < |\ell| \leq K \\ 0 & \text{otherwise} \end{cases} \quad (4.14)$$

Note that S is well defined for $\omega \in \mathcal{C}^{(1)}$ (defined in (3.4)) and, by Proposition 3.1, one has $S \in \mathcal{L}_{a, -2}^{\mathfrak{qT}, \mathcal{O}_+}$, with the bound (3.9). Then

$$\begin{aligned} G_*(D + P) &= e^{\text{ad}S}(D + P) - i \sum_{h=1}^{\infty} \frac{(\text{ad}S)^{h-1}}{h!} \dot{S} \\ &= (D + P) + \sum_{h=1}^{\infty} \frac{(\text{ad}S)^{h-1}}{h!} ([D + P, S] - i\dot{S}) \\ &= D + \text{diag}(P_j^j(0)) + \Pi_{>K} P + [P, S] + \sum_{h=2}^{\infty} \frac{(\text{ad}S)^{h-1}}{h!} ([P, S] - \Pi_{0 < |\ell| \leq K} P); \end{aligned} \quad (4.15)$$

so for $\omega \in \mathcal{C}^{(1)}$ we put

$$P_+ := \Pi_{>K} P + [P, S] + \sum_{h=2}^{\infty} \frac{(\text{ad}S)^{h-1}}{h!} ([P, S] - \Pi_{0 < |\ell| \leq K} P), \quad D_+ := D + \text{diag}(P_j^j(0))_{j \in \mathbb{Z}^2}.$$

By construction $P = P^{\mathbb{T}} + P^{(1)} + P^{(2)}$ as in (2.16). Now by Lemmas 2.19 and 2.20

$$P_j^j(0) = \mathfrak{P}(0, v(j), b(j)) + (P^{(1)})_j^j(0) + (P^{(2)})_j^j(0).$$

So in order to define D_+ on all of \mathbb{R} we just extend each $\mathfrak{P}(0, v, b), (P^{(i)})_j^j(0)$ by Kirszbraun theorem preserving its weighted Lipschitz norm $|\cdot|^\gamma$. The bound (4.13) follows. The bound (4.12) follows by Corollary 2.22 and Lemma 2.19. \square

Proposition 4.6 (Iteration). *Let \mathfrak{L}_0 be a Schrödinger operator of the form*

$$-\Delta + P_0, \quad -\Delta = \text{diag}(|j|^2), \quad P_0 \in \mathcal{L}_{a_0, -2}^{\mathfrak{qT}, \mathcal{O}_0}, \quad \varepsilon_0 := \gamma^{-1} |P_0|_{a_0, -2}^{\mathfrak{qT}, \mathcal{O}_0} < K_0^{-3\tau-2} \quad (4.16)$$

where K_0 is large enough. Set

$$a_n = a_0 \left(1 - \sum_{k=0}^{n-1} 2^{-k-2}\right), \quad K_n := K_0 4^n, \quad \varepsilon_n = \varepsilon_0 e^{-(\frac{3}{2})^{n+1}}. \quad (4.17)$$

For all $n \geq 1$, there exist

(i)_n A sequence of time-independent operators

$$D_n = -\Delta + \tilde{\Omega}^{(n)}, \quad \text{where } \tilde{\Omega}^{(n)} \in \mathcal{L}_{a_n, -2}^{\mathfrak{qT}, \mathbb{R}} \quad \text{with } |\tilde{\Omega}^{(n)} - \tilde{\Omega}^{(n-1)}|_{a_n, -2}^{\mathfrak{qT}, \mathbb{R}} \leq \gamma \varepsilon_{n-1} \quad (4.18)$$

(we have set $\tilde{\Omega}^{(0)} := 0$).

Following Lemma 2.20 we denote the eigenvalues as

$$\tilde{\Omega}_j^{(n)} = \mathfrak{a}^{(n)}(v(j), b(j)) + r_j^{(n,1)} + r_j^{(n,2)} \quad (4.19)$$

and define the nested sequence of compact sets (recall (3.4), (3.5))

$$\mathcal{O}_n := \mathcal{O}_{n-1} \cap \mathcal{C}_{D_{n-1}, K_{n-1}}^{(1)}(\gamma) \cap \mathcal{C}_{D_{n-1}, K_{n-1}}^{(2)}(\gamma). \quad (4.20)$$

(ii)_n A sequence of operators $P_n \in \mathcal{L}_{a_n, -2}^{\mathfrak{qT}, \mathcal{O}_n}$, $S_n \in \mathcal{L}_{a_{n-1}, -2}^{\mathfrak{qT}, \mathcal{O}_n}$ such that

$$(e^{S_n})_*(D_{n-1} + P_{n-1}) = D_n + P_n, \quad \forall \omega \in \mathcal{O}_n, \quad (4.21)$$

$$\gamma^{-1} |P_n|_{a_n, -2}^{\mathfrak{qT}, \mathcal{O}_n} \leq \varepsilon_n, \quad |S_n|_{a_{n-1}, -2}^{\mathfrak{qT}, \mathcal{O}_n} \leq K_{n-1}^{3\tau+2} \varepsilon_{n-1}. \quad (4.22)$$

Proof. The lemma is proved by standard iterative estimates. \square

Corollary 4.7. *Given a Schrödinger operator \mathfrak{L}_0 satisfying (4.16) there exist a time independent operator $\tilde{\Omega} \in \mathcal{L}_{a_0/2, -2}^{\mathfrak{qT}, \mathbb{R}}$, a Cantor-like set \mathcal{O}_∞ and a symplectic change of variables G with $G - \text{Id} \in \mathcal{L}_{a_0/2, -2}^{\mathfrak{qT}, \mathcal{O}_\infty}$ such that*

$$G_* \mathfrak{L}_0 = -\Delta + \tilde{\Omega}, \quad \forall \omega \in \mathcal{O}_\infty. \quad (4.23)$$

Proof. The proof of this lemma is also standard and follows from the fast convergence of the $\tilde{\Omega}^{(n)}$ in Proposition 4.6. We put $\mathcal{O}_\infty := \bigcap_n \mathcal{O}_n$. \square

Proof of Theorem 4.1. The thesis follows from Lemma 4.3 and Corollary 4.7, except from the fact that \mathcal{O}_∞ has positive measure. In order to show this, we shall prove Proposition 4.2, which gives a cleaner characterization of the set \mathcal{O}_∞ . Then, the measure estimates are deferred to the next section. \square

Proof of Proposition 4.2. We need to show that $\mathcal{O}_{\text{fin}} \subset \bigcap_{n=0}^\infty \mathcal{O}_n$, i.e. we have to verify conditions (3.4)-(3.5) for $\omega \in \mathcal{O}_{\text{fin}}$. Proving (3.4) is standard (see for instance [1]). Regarding (3.5), by (4.5) one has, for $\omega \in \mathcal{O}_{\text{fin}}$,

$$\begin{aligned} & \left| \omega \cdot \ell + 2 \frac{|\pi(\ell)|}{|v|} b - |\pi(\ell)|^2 + \mathfrak{a}^{(n)}(v, b) - \mathfrak{a}^{(n)}(v, b - v \cdot \pi(\ell)) \right| \geq \\ & 4\gamma |\ell|^{-\tau} - |\mathfrak{a}^{(n)}(v, b) - \mathfrak{a}(v, b)| - |\mathfrak{a}^{(n)}(v, b - v \cdot \pi(\ell)) - \mathfrak{a}(v, b - v \cdot \pi(\ell))|. \end{aligned}$$

The estimate follows by recalling that $|\ell| < K_n$, using (2.18) to bound $|\mathfrak{a}^{(n)}(v, b) - \mathfrak{a}(v, b)|$ in terms of $|\tilde{\Omega}^{(n)} - \tilde{\Omega}|_{a_n, -2}^{\mathfrak{qT}, \mathbb{R}}$ and then exploiting the bounds (4.18) (see e.g. the proof of Lemma 7.6 of [35] for details). \square

5 Measure estimates

The purpose of this section is to prove that the set of frequencies ω for which the non-resonance conditions (1.29) hold up to order N has positive measure.

Recalling the notations of Theorem 4.1 and denoting

$$\Omega_j = |j|^2 + \mathbf{a}(v(j), b(j)) + R(j), \quad R(j) = r_j^{(1)} + r_j^{(2)},$$

we have

$$\sup_{v,b} e^{\mathbf{a}'c|v|} \langle b \rangle^2 |\mathbf{a}(v, b)|^{\mathbf{R}} + \sup_j |R(j)|^{\mathbf{R}} \langle j \rangle^\mu \leq 2\epsilon\gamma. \quad (5.1)$$

Denote by

$$\mathbf{v} := (v_h)_{h=1,\dots,N} \in \mathcal{V}^N \quad \mathbf{b} := (b_h)_{h=1,\dots,N} \in \mathbb{Z}^N, \quad \boldsymbol{\eta} := (\eta_{ab})_{\substack{a=1,2 \\ b=1,\dots,N}} \in \{-1, 0, 1\}^{2N}$$

and for $(\ell, K, \mathbf{v}, \mathbf{b}, \boldsymbol{\eta}) \in \mathfrak{A}_N := \mathbb{Z}^d \times \mathbb{Z} \times \mathcal{V}^N \times \mathbb{Z}^N \times \{-1, 0, 1\}^{2N}$, consider an expression of the form

$$\mathbf{d}(\ell, K, \mathbf{v}, \mathbf{b}, \boldsymbol{\eta}) := \omega \cdot \ell + K + \sum_{h=1}^N \eta_{1h} \mathbf{a}(v_h, b_h) + \sum_{k=1}^N \eta_{2k} R(j_k) \quad (5.2)$$

We shall prove the following result:

Lemma 5.1. *Given $N \in \mathbb{N}$, $N \geq 2$, there exist $\gamma_*, \epsilon_*, \tau_N > 0$ such that for any $0 < \gamma < \gamma_*$ and $\epsilon < \epsilon_*$ the set*

$$\mathcal{C}_*(N, \gamma, \tau_N) := \mathcal{C}_* := \left\{ \omega \in \mathbb{R} : |\mathbf{d}(\ell, K, \mathbf{v}, \mathbf{b}, \boldsymbol{\eta})| \geq \frac{\gamma}{\langle \ell \rangle^{\tau_N}}, \quad \forall (\ell, K, \mathbf{v}, \mathbf{b}, \boldsymbol{\eta}) \in \mathfrak{A}_N : (\ell, K) \neq (0, 0) \right\} \quad (5.3)$$

has positive measure. More precisely there exists a constant $c_* > 0$ such that

$$\text{meas}(\mathbb{R} \setminus \mathcal{C}_*) \leq c_* \gamma.$$

Proof. We first notice that for all $L > 0$, we can ensure that

$$|(5.2)| \geq \frac{\gamma}{\langle \ell \rangle^{d+1}}, \quad \forall (\ell, K) \neq (0, 0), \quad |\ell| \leq L$$

just by taking $\epsilon_* = \epsilon_*(L)$ small enough. So we choose $L > 0$ (to be fixed later). Next we note that, for $\ell \neq 0$, if $|K| > 2|\ell|$ then (for ϵ_* small enough) one has $|\mathbf{d}(\ell, K, \mathbf{v}, \mathbf{b}, \boldsymbol{\eta})| \geq \frac{1}{2} > \gamma$.

So from now on we restrict to $|\ell| \geq L$ and $|K| \leq 2|\ell|$.

We prove the claim by finite induction on the number of η_{r_1, r_2} different from 0. More precisely for every $0 \leq n \leq 2N$ we shall show that for γ small enough, there exist a sequence of increasing τ_n and a sequence of nested sets $\mathcal{C}^n = \mathcal{C}^n(\gamma, \tau_n)$ such that, provided

$$|\eta_{11}| + \dots + |\eta_{23}| = n,$$

then for any $\omega \in \mathcal{C}^n$ one has

$$|(5.2)| \geq \frac{\gamma}{\langle \ell \rangle^{\tau_n}}, \quad \forall (\ell, K) \neq (0, 0) \quad (5.4)$$

and moreover

$$\text{meas}(\mathbb{R} \setminus \mathcal{C}^0) \leq C\gamma, \quad \text{meas}(\mathcal{C}^n \setminus \mathcal{C}^{n+1}) \leq C\gamma \quad (5.5)$$

for some $C > 0$. Then the lemma follows by taking $\mathcal{C}_* := \mathcal{C}^{2N}$ and $c_* := (2N + 1)C$. Then γ_* is fixed so that $(2N + 1)C\gamma_* < \text{meas}(\mathbb{R})$, and the measure of \mathcal{C}_* results positive.

Case $n = 0$. In this case (5.2) reduces to $\omega \cdot \ell + K$, so the set \mathcal{C}^0 coincides with the set \mathcal{O}_0 defined in (4.3), with $\tau_0 := d + 1$. It is well known that $\text{meas}(\mathbb{R} \setminus \mathcal{O}_0) \leq C\gamma$ for some $C > 0$.

Case $n \rightsquigarrow n+1$. Assume that (5.4) holds for any possible choice of (η_{ab}) s.t. $|\eta_{11}| + \dots + |\eta_{23}| = n$, $1 \leq n \leq 2N - 1$, for some $(\tau_a)_{a=1, \dots, n}$. We can assume that $|\ell| \geq L$ and $|\eta_{11}| + \dots + |\eta_{23}| = n + 1$. We further divide two subcases.

Case I. At least one of the following conditions holds:

I.a There exists $h = 1, \dots, N$ such that $\eta_{1h} \neq 0$ and either $\mathbf{ca}'|v_h| \geq |\ell|$ or $|b_h| \geq |\ell|^{\tau_n/2}$.

I.b There exists $h = 1, \dots, N$ such that $\eta_{2h} \neq 0$ and $\langle j_h \rangle \geq |\ell|^{\tau_n/\mu}$.

In Case I.a, assume w.l.o.g. that $\eta_{1N} \neq 0$ then

$$\begin{aligned} |(5.2)| &\geq |\omega \cdot \ell + K + \sum_{a=1}^{N-1} \eta_{1a} \mathbf{a}(v_a, b_a) + \sum_{a=1}^N \eta_{2a} R(j_a)| - |\mathbf{a}(v_N, b_N)| \\ &\geq \frac{\gamma}{\langle \ell \rangle^{\tau_n}} - 2\gamma \epsilon \langle b_N \rangle^{-2} e^{-\mathbf{ca}'|v_N|} \geq \frac{\gamma}{\langle \ell \rangle^{\tau_n+1}} \end{aligned}$$

which is true provided ϵ_* is small enough and L sufficiently large. In Case I.b, assume w.l.o.g. that $\eta_{2N} \neq 0$ then

$$\begin{aligned} |(5.2)| &\geq |\omega \cdot \ell + K + \sum_{a=1}^N \eta_{1a} \mathbf{a}(v_a, b_a) + \sum_{a=1}^{N-1} \eta_{2a} R(j_a)| - |R(j_N)| \\ &\geq \frac{\gamma}{\langle \ell \rangle^{\tau_n}} - \gamma \epsilon \frac{1}{\langle j_N \rangle^\mu} \geq \frac{\gamma}{\langle \ell \rangle^{\tau_n+1}} \end{aligned}$$

Case II. For all $k = 1, 2$ and $h = 1, \dots, N$ such that $\eta_{kh} \neq 0$ one has $\mathbf{ca}'|v_h| < |\ell|$ and $|b_h| < |\ell|^{\tau_n/2}$ if $k = 1$ and one has $\langle j_h \rangle < |\ell|^{\tau_n/\mu}$ if $k = 2$. Now for $|\ell| \geq L$, $|K| \leq 2|\ell|$ and $\mathbf{v}, \mathbf{b}, \boldsymbol{\eta}$ as in Case II we define

$$\mathcal{R}_{\ell, K, \mathbf{v}, \mathbf{b}, \boldsymbol{\eta}}(\tau_{n+1}, \gamma) := \{\omega \in \mathbb{R} : |\mathbf{d}(\ell, K, \mathbf{v}, \mathbf{b}, \boldsymbol{\eta})| \leq \gamma |\ell|^{-\tau_{n+1}}\}.$$

Now (recall $|\ell| \geq L$ and the Lipschitz estimates on $\mathbf{a}(v, b)$ and $R(j)$)

$$\left| \frac{d}{d|\omega|} \mathbf{d}(\ell, K, \mathbf{v}, \mathbf{b}, \boldsymbol{\eta}) \right| \geq |\ell| - 2N\epsilon \geq 1$$

so $\text{meas}(\mathcal{R}_{\ell, K, \mathbf{v}, \mathbf{b}, \boldsymbol{\eta}}) \leq 2\gamma |\ell|^{-\tau_{n+1}}$ and

$$\text{meas} \left(\bigcup_{\substack{|\ell| \geq L, |K| \leq 2|\ell| \\ \mathbf{v}, \mathbf{b}, \boldsymbol{\eta} \in \text{Case II}}} \mathcal{R}_{\ell, K, \mathbf{v}, \mathbf{b}, \boldsymbol{\eta}} \right) \leq \sum_{|\ell| \geq L} \sum_{\substack{|K| \leq 2|\ell|, |v_h| \leq |\ell| \\ |b_h| \leq |\ell|^{\tau_n/2}, \langle j_h \rangle \leq |\ell|^{\tau_n/\mu}}} \frac{\gamma}{|\ell|^{\tau_{n+1}}} \leq C\gamma \sum_{|\ell| \geq L} |\ell|^{2N+1 + (\frac{1}{2} + \frac{2}{\mu})N\tau_n - \tau_{n+1}}$$

which is finite provided

$$\tau_{n+1} > \left(\frac{5}{2} + \frac{2}{\mu} \right) N\tau_n + d + 1.$$

Then we put

$$\mathcal{C}^{n+1} := \mathcal{C}^n \setminus \bigcup_{\substack{|\ell| \geq L, \\ |K| \leq 2|\ell| \\ \mathbf{v}, \mathbf{b}, \boldsymbol{\eta} \in \text{Case II}}} \mathcal{R}_{\ell, K, \mathbf{v}, \mathbf{b}, \boldsymbol{\eta}}$$

and clearly it fulfills (5.5). This concludes the proof of the inductive step. \square

We are now ready to discuss the measure of the non-resonant ω . Let us start by discussing the set \mathcal{O}_{fin} of Proposition 4.2.

Proposition 5.2. *There exist $\gamma_*, \epsilon_*, \tau > 0$ such that for any $0 < \gamma < \gamma_*$ and $\epsilon < \epsilon_*$, the set \mathcal{O}_{fin} of Proposition 4.2 has positive measure, in particular*

$$\text{meas}(\mathbb{R} \setminus \mathcal{O}_{\text{fin}}) \leq C\gamma.$$

Proof. We show that the set \mathcal{C}_* defined in (5.3) with $N = 2$ is contained in \mathcal{O}_{fin} . Indeed $\mathcal{C}^{(1)}$ defined in (4.4) contains \mathcal{C}_* , just taking $j_1 = j$, $j_2 = j - \pi(\ell)$, $(v_i, b_i) = (v(j_i), b(j_i))$ for $i = 1, 2$, $K = |j_1|^2 - |j_2|^2$, $\eta_{11} = \eta_{21} = 1$, $\eta_{12} = \eta_{22} = -1$.

Also $\mathcal{C}^{(2)}$ defined in (4.5) contains \mathcal{C}_* , setting $v_1 = v_2 = v$, $b_1 = b$, $b_2 = b - v \cdot \pi(\ell)$, $K = 2 \frac{|\pi(\ell)|}{|v|} b - |\pi(\ell)|^2$ (note that $\frac{|\pi(\ell)|}{|v|} b \in \mathbb{Z}$ as $\pi(\ell) \parallel v$) and taking $\eta_{11} = -\eta_{12}$, $\eta_{21} = \eta_{22} = 0$. \square

We finally prove Theorem 1.8:

Proof of Theorem 1.8. Reducibility is a consequence of Theorem 4.1. It only remains to prove that $\forall N \in \mathbb{N}$ Melnikov conditions of order N are satisfied, namely that $\mathcal{C}_* \subseteq \mathcal{G}_N$, with \mathcal{G}_N as in (1.28) and \mathcal{C}_* as in (5.3). First we observe that, since L as in (1.28) satisfies $|L| \leq N$, it has at most N non vanishing components L_{j_1}, \dots, L_{j_N} . Then it is sufficient to set

$$\begin{aligned} \mathbf{v} &= \underbrace{(v(j_1), \dots, v(j_1))}_{|L_{j_1}| \text{ times}}, \dots, \underbrace{(v(j_N), \dots, v(j_N))}_{|L_{j_N}| \text{ times}}, \quad \mathbf{b} = \underbrace{(b(j_1), \dots, b(j_1))}_{|L_{j_1}| \text{ times}}, \dots, \underbrace{(b(j_N), \dots, b(j_N))}_{|L_{j_N}| \text{ times}}, \\ \boldsymbol{\eta} &= \underbrace{(\text{sign}(L_{j_1}), \dots, \text{sign}(L_{j_1}))}_{|L_{j_1}| \text{ times}}, \dots, \underbrace{(\text{sign}(L_{j_N}), \dots, \text{sign}(L_{j_N}))}_{|L_{j_N}| \text{ times}}, \quad K = \sum_{j \in \mathbb{Z}^2} |j|^2 L_j. \end{aligned} \quad (5.6)$$

Then the result follows by Lemma 5.1, observing that the set \mathcal{G}_N is defined in such a way that, with K as in (5.6), one always has $(\ell, K) \neq (0, 0)$. \square

6 Nonlinear stability results

In order to prove stability we apply a classical Birkhoff normal form procedure, see for instance Theorem 8.1 of [35]. We start by recalling some standard notions on time dependent Hamiltonians.

One works in the *extended phase space* $(\varphi, \mathcal{Y}, u) \in \mathbb{T}^d \times \mathbb{R}^d \times \mathfrak{H}^p$ with the symplectic form $d\mathcal{Y} \wedge d\varphi + idu \wedge d\bar{u}$.

In this context we shall consider *real on real* analytic functions on the complex domain

$$D(a, r) := \mathbb{T}_a^d \times D(r), \quad D(r) := \left\{ \mathcal{Y} \in \mathbb{C}^d : |\mathcal{Y}|_1 := \sum_{i=1}^d |\mathcal{Y}_i| < r^2, \quad u \in \mathfrak{H}^p : \|u\|_p < r \right\},$$

with \mathbb{T}_a^d defined in (2.4). Given a real analytic function $F(\varphi, \mathcal{Y}, u)$ we consider its Taylor-Fourier series

$$F(\varphi, \mathcal{Y}, u) = \sum_{\substack{\alpha, \beta \in \mathbb{N}^{2^2} \\ \ell \in \mathbb{Z}^d, l \in \mathbb{N}^d}} F_{\alpha, \beta, l, \ell} e^{i\ell \cdot \varphi} \mathcal{Y}^l u^\alpha \bar{u}^\beta, \quad F_{\alpha, \beta, l, \ell} = \bar{F}_{\beta, \alpha, l, -\ell}, \quad u^\alpha \bar{u}^\beta = \prod_{j \in \mathbb{Z}^2} u_j^{\alpha_j} \bar{u}_j^{\beta_j}$$

Correspondingly we expand vector fields in Taylor Fourier series (again well defined and pointwise absolutely convergent):

$$X^{(w)}(\varphi, \mathcal{Y}, u) = \sum_{\alpha, \beta \in \mathbb{N}^{2^2}, \ell \in \mathbb{Z}^d, l \in \mathbb{N}^d} X_{\alpha, \beta, l, \ell}^{(w)} e^{i\ell \cdot \varphi} \mathcal{Y}^l u^\alpha \bar{u}^\beta,$$

where w denotes the components φ_i, \mathcal{Y}_i or u_j, \bar{u}_j . To a vector field we associate its *majorant*

$$\underline{X}_a^{(w)}[\mathcal{Y}, u] := \sum_{\ell \in \mathbb{Z}^d, l \in \mathbb{N}^d, \alpha, \beta \in \mathbb{N}^{2^2}} |X_{\alpha, \beta, l, \ell}^{(w)}| e^{a|\ell|} \mathcal{Y}^l u^\alpha \bar{u}^\beta. \quad (6.1)$$

Then we have the following

Definition 6.1. A vector field $X : D(a, r) \rightarrow \mathbb{C}^d \times \mathbb{C}^d \times \mathfrak{H}^p$ will be said to be majorant analytic in $D(a, r)$ if \underline{X}_a defines an analytic vector field $D(r) \rightarrow \mathbb{C}^d \times \mathbb{C}^d \times \mathfrak{H}^p$.

Since Hamiltonian functions are defined modulo constants, we give the following definition of regular Hamiltonian and its norm:

Definition 6.2. A real valued Hamiltonian H will be said to be regular in $D(a, r)$ if its Hamiltonian vector field X_H is majorant analytic in $D(a, r)$. We define its norm by

$$|H|_{a, r} := \sup_{(\mathcal{Y}, u) \in D(r)} \|(X_H)_a(\mathcal{Y}, u)\|_r, \quad \|X\|_r := |X^{(\varphi)}|_\infty + \frac{|X^{(\mathcal{Y})}|_1}{r^2} + \frac{\|X^{(u)}\|}{r} + \frac{\|X^{(\bar{u})}\|}{r}. \quad (6.2)$$

Note that the norm $|F|_{a, r}$ controls the norm $\|X_F\|_r$ in $D(a, r)$.

If $F(\omega; \varphi, \mathcal{Y}, u) \equiv F(\omega)$ depends on the frequency $\omega \in \mathcal{O}$, where \mathcal{O} is some compact set, we define the *weighted Lipschitz* norm:

$$|F|_{a, r}^{\mathcal{O}} := \sup_{\omega \in \mathcal{O}} |F(\omega, \cdot)|_{a, r} + \gamma \sup_{\omega_1 \neq \omega_2 \in \mathcal{O}} \frac{|F(\omega_1, \cdot) - F(\omega_2, \cdot)|_{a, r}}{|\omega_1 - \omega_2|}. \quad (6.3)$$

Definition 6.3. We say that a regular Hamiltonian H is Gauge and Momentum preserving if

$$\pi(\ell) + \sum_{j \in \mathbb{Z}^2} j(\alpha_j - \beta_j) \neq 0 \quad \text{or} \quad \sum_{i=1}^d \ell_i + \sum_{j \in \mathbb{Z}^2} (\alpha_j - \beta_j) \neq 0 \quad \Rightarrow \quad H_{\alpha, \beta, l, \ell} = 0$$

By convention we define the scaling degree of a monomial $e^{i\ell \cdot \varphi} \mathcal{Y}^l u^\alpha \bar{u}^\beta$ as $\deg(l, \alpha, \beta) := 2|l| + |\alpha| + |\beta| - 2$, and define the projection on the homogeneous components of scaling degree d as

$$H^{(d)} := \sum_{\substack{\ell \in \mathbb{Z}^d, l \in \mathbb{N}^d, \alpha, \beta \in \mathbb{N}^{2^2} \\ 2|l| + |\alpha| + |\beta| = d+2}} H_{\alpha, \beta, l, \ell} e^{i\ell \cdot \varphi} \mathcal{Y}^l u^\alpha \bar{u}^\beta,$$

similarly for $H^{(\leq d)}$ and $H^{(\geq d)}$. One easily verifies that $\{H^{(d_1)}, H^{(d_2)}\}$ has scaling degree $d_1 + d_2$.

Proof of Theorem 1.4. A direct computation ensures that the NLS equation (1.15) is Hamiltonian

$$H = \omega \cdot \mathcal{Y} + \sum_j |j|^2 |u_j|^2 + (u, V(\varphi)u)_{\ell_2(\mathbb{Z}^2)} + P(\varphi, u), \quad P = \int_{\mathbb{T}^2} P(\varphi + \mathbf{K}x, u(x), \bar{u}(x)) dx. \quad (6.4)$$

By assumptions P is a Gauge and Momentum preserving regular Hamiltonian of scaling ≥ 1 , with norm

$$|P|_{a,r} \leq Cr, \quad \forall r \leq r_*.$$

We now fix $0 < \gamma < \gamma_*$ and assume that $\|V\|_{\mathfrak{a},p} \gamma^{-1} < \epsilon_*$ so that we may apply Theorem 4.1. The time dependent symplectic change of variables G defined in Theorem 4.1 gives rise to the symplectic transformation \mathcal{G} defined in (2.8) on the extended phase space. By definition, \mathcal{G} preserves the scaling degree and maps $D(\mathfrak{a}/8, r) \rightarrow D(\mathfrak{a}/8, \rho r)$ for all $r > 0$, where $\rho - 1 \lesssim \|V\|_{\mathfrak{a},p} \gamma^{-1}$. Thus for all regular F of scaling ≥ 1 one has that $F \circ \mathcal{G}$ is still regular, of scaling ≥ 1 , and satisfies

$$|F \circ \mathcal{G}|_{a,\rho r} \leq 2|F|_{a,r}, \quad \forall a \leq \mathfrak{a}/8.$$

Finally, since V is Gauge invariant then \mathcal{G} is Gauge covariant. By construction \mathcal{G} reduces the Hamiltonian H to the diagonal form

$$H_1 = H \circ \mathcal{G} = \omega \cdot \mathcal{Y} + \sum_j \Omega_j |u_j|^2 + \mathcal{K}(\varphi, u), \quad |\mathcal{K}|_{\mathfrak{a}/8,r} \leq Cr, \quad \forall r \leq r_*,$$

where \mathcal{K} is a Gauge and Momentum preserving regular Hamiltonian of scaling ≥ 1 . We note that, thanks to Theorem 1.8 with $N = 3$, we can impose third order Melnikov conditions of the formula (1.29) for any $(\ell, L) \in \mathcal{G}_3$. Since \mathcal{K} is a regular Gauge and momentum preserving Hamiltonian, we can eliminate all of its monomials of scaling degree 1. Precisely, as H_1 fits the hypotheses of Theorem 8.1 of [35] (with $\varepsilon = 1$), there exists a symplectic change of variables which cancels the terms of scaling degree one in \mathcal{K} and conjugates H_1 to

$$H_2 = \omega \cdot \mathcal{Y} + \sum_j \Omega_j |u_j|^2 + \tilde{\mathcal{K}}(\varphi, u), \quad |\tilde{\mathcal{K}}|_{\mathfrak{a}/8,r} \leq Cr^2, \quad \forall r \leq r_*,$$

where $\tilde{\mathcal{K}}$ has scaling ≥ 2 . Then the stability times in Theorem 1.4 follow by a standard contraction mapping Lemma argument, see for instance Corollary 5.1 of [9].

A Technical lemmas

Lemma A.1 (Bony). *Given $M \in \mathcal{L}_a(\mathfrak{H}^p)$ and setting*

$$(M^{(B)})_j^{j'}(\ell) := \begin{cases} 0 & \text{if } |\ell| > \mathfrak{c}\langle j \rangle^\delta \\ M_j^{j'}(\ell) & \text{otherwise} \end{cases}, \quad M^{(R)} = M - M^{(B)} \quad (\text{A.1})$$

we have that $M^{(R)} \in \mathcal{L}_{a',-n}$ for any $n > 0$ and any $0 \leq a' < a$ with the quantitative estimate

$$|M^{(R)}|_{a',-n} \lesssim_{p,\delta} \frac{|M|_a}{(a - a')^{\mu n/\delta}} \quad (\text{A.2})$$

Proof. Exploiting the definition (2.11) we have

$$\begin{aligned} (\underline{M}_{a',-n}^{(R)})_j^{j'} &:= \sum_{\substack{\ell: j-j'=\pi(\ell) \\ |\ell|>c\langle j \rangle^\delta}} e^{a'|\ell|} |M_j^{j'}(\ell)| \langle j \rangle^{\mu n} \leq c^{-\mu n/\delta} \sum_{\ell: j-j'=\pi(\ell)} e^{a|\ell|} e^{-(a-a')|\ell|} |\ell|^{n\mu/\delta} |M_j^{j'}(\ell)| \\ &\leq C(\mu n/\delta, a-a') (\underline{M}_a)_j^{j'}, \end{aligned}$$

with $C(p, \sigma)$ defined as in (2.12) for $p, \sigma > 0$. \square

Proof of Lemma 2.14. To prove (2.13), we just use the fact that one has

$$\begin{aligned} (\underline{MN}_{a,-\mathbf{N}_1})_j^{j'} &:= \sum_{\ell_1, \ell_2: j-j'=\pi(\ell_1+\ell_2)} e^{a|\ell_1+\ell_2|} |M_j^{j-\pi(\ell_1)}(\ell_1)| \langle j \rangle^{\mu n_1} |b(j)|^{m_1} |N_{j-\pi(\ell_1)}^{j'}(\ell_2)| \\ &\leq \sum_{j_1} (\underline{M}_{a,-\mathbf{N}_1})_j^{j_1} (\underline{N}_{a,\bar{0}})_{j_1}^{j'}. \end{aligned}$$

For the second statement we divide $M = M^{(B)} + M^{(R)}$ as in (A.1), then

$$\begin{aligned} (\underline{M}^{(B)} N_{a;-(n_1+n_2,0)})_j^{j'} &:= \sum_{\substack{\ell_1, \ell_2: j-j'=\pi(\ell_1+\ell_2) \\ |\ell_1|<c\langle j \rangle^\delta}} e^{a|\ell_1+\ell_2|} |M_j^{j-\pi(\ell_1)}(\ell_1)| \langle j \rangle^{\mu(n_1+n_2)} |N_{j-\pi(\ell_1)}^{j'}(\ell_2)| \\ &\leq \sum_{\substack{\ell_1, \ell_2: j-j'=\pi(\ell_1+\ell_2) \\ |\ell_1|<c\langle j \rangle^\delta}} \left(e^{a|\ell_1|} |M_j^{j-\pi(\ell_1)}(\ell_1)| \langle j \rangle^{\mu n_1} e^{a|\ell_2|} \langle j - \pi(\ell_1) \rangle^{\mu n_2} |N_{j-\pi(\ell_1)}^{j'}(\ell_2)| \right) \left(\frac{\langle j \rangle}{\langle j - \pi(\ell_1) \rangle} \right)^{\mu n_2} \\ &\leq 2^{\mu n_2} \sum_{j_1} (\underline{M}^{(B)})_{a;-\mathbf{N}_1}^{j_1} (\underline{N}_{a;-\mathbf{N}_2})_{j_1}^{j'}. \end{aligned}$$

Moreover by (2.13) with $M \rightsquigarrow M^{(R)}$ and estimate (A.2) we have

$$|M^{(R)} N|_{a';-(n_1+n_2,0)} \leq |M^{(R)}|_{a';-(n_1+n_2,0)} |N|_{a';\bar{0}} \lesssim \frac{|M|_a |N|_{a'}}{(a-a')^{\frac{\mu(n_1+n_2)}{\delta}}}.$$

\square

B Proof of Proposition 2.21 and Lemma 4.4

Before proving Proposition 2.21, we give the following auxiliary result:

Lemma B.1. *Given $n_1, m_1, m_2 \in \mathbb{N}$, let $\mathbf{N}_1 = (n_1, m_1)$, $R \in \mathcal{L}_{\frac{a}{2};-\mathbf{N}_1}$ and $T \in \mathcal{T}_{a,-m_2}$. Then $RT \in \mathcal{L}_{\frac{a}{2};-\mathbf{N}_1}$, and $TR \in \mathcal{L}_{\frac{a}{2};-(n_1, m_2)}$, with the bounds*

$$|RT|_{\frac{a}{2};-\mathbf{N}_1} \lesssim a^{-p-2} |R|_{\frac{a}{2};-\mathbf{N}_1} |T|_{a,0}^{\mathbb{T}}, \quad (\text{B.1})$$

$$|TR|_{\frac{a}{2};-(n_1, m_2)} \lesssim_{\mu, n_1} a^{-\mu n_1} |R|_{\frac{a}{2};-(n_1, 0)} |T|_{a,-m_2}^{\mathbb{T}}. \quad (\text{B.2})$$

Proof. Bound (B.1) follows directly from Lemma 2.14 and Lemma 2.17. Analogously, to prove

(B.2) one observes that

$$\begin{aligned}
(\underline{TR}_{\frac{a}{2}; -(n_1, m_2)}^j)^{j'} &:= \sum_{\substack{\ell_1, \ell_2: \\ j-j'=\pi(\ell_1+\ell_2)}} e^{\frac{a}{2}|\ell_1+\ell_2|} |T_j^{j-\pi(\ell_1)}(\ell_1)| |R_{j-\pi(\ell_1)}^{j'}(\ell_2)| \langle j \rangle^{\mu n_1} |b(j)|^{m_2} \\
&\lesssim_{n_1, \mu} a^{-\mu n_1} \sum_{\substack{\ell_1, \ell_2: \\ j-j'=\pi(\ell_1+\ell_2)}} e^{\frac{3}{4}a|\ell_1|} |T_j^{j-\pi(\ell_1)}(\ell_1)| |b(j)|^{m_2} e^{\frac{a}{2}|\ell_2|} \langle j-\pi(\ell_1) \rangle^{\mu n_1} |R_{j-\pi(\ell_1)}^{j'}(\ell_2)| \\
&\lesssim_{\mu, n_1} a^{-\mu n_1} \sum_{j_1} (\underline{T}_{\frac{3}{4}a, 0, -m_2})_{j_1}^{j_1} (\underline{R}_{\frac{a}{2}, -n_1, 0})_{j_1}^{j_1},
\end{aligned}$$

and the result follows by Lemma 2.14, by Remark 2.8, and by Lemma 2.17. \square

Proof of Proposition 2.21. We start by proving estimate (2.19).

As $M_\iota \in \mathcal{A}_{a, -m_\iota}$, $\iota = 1, 2$, there exist operators $M_\iota^T, M_\iota^{(1)}, \dots, M_\iota^{(m_\iota)}$ such that $M_\iota = M_\iota^T + \sum_{i=1}^{m_\iota} M_\iota^{(i)}$ and

$$|M_\iota^T|_{a, -m_\iota}^T + \sum_{i=1}^{m_\iota} |M_\iota^{(i)}|_{\frac{a}{2}; -(i, m_\iota - i)} \leq 2|M_\iota|_{a, -m_\iota}^{\text{qT}}. \quad (\text{B.3})$$

To each M_ι^T we may associate the symbol $\mathfrak{T}_\iota(\ell, v, b)$. First of all we define the line-Töplitz part of $M_1 M_2$ by setting $(M_1 M_2)^T := T$ with

$$(T)_j^{j'}(\ell) := \mathfrak{T}(\ell, v(j), b(j)), \quad \mathfrak{T}(\ell, v, b) := \sum_{\ell_1 + \ell_2 = \ell} \mathfrak{T}_1(\ell_1, v, b) \mathfrak{T}_2(\ell_2, v, b - v \cdot \pi(\ell_1)).$$

We estimate its $|T|_{a, -m}^T$ norm, given by (2.15), with $m = \min(m_1, m_2)$:

$$\begin{aligned}
|T|_{a, -m}^T &\leq \sup_{v \in \mathcal{V}, b \in \mathbb{Z}} \sum_{\ell \in \mathbb{Z}^d} e^{a(c|v|+|\ell|)} |\mathfrak{T}(\ell, v, b)| \langle b \rangle^m \\
&\leq \sup_{v \in \mathcal{V}, b \in \mathbb{Z}} \sum_{\ell_1 \in \mathbb{Z}^d} e^{a|\ell_1|} |\mathfrak{T}_1(\ell_1, v, b)| \langle b \rangle^{m_1} \sup_{v' \in \mathcal{V}, b' \in \mathbb{Z}} \sum_{\ell_2 \in \mathbb{Z}^d} e^{a(c|v'|+|\ell_2|)} |\mathfrak{T}_2(\ell_2, v', b')| \leq |M_1^T|_{a, -m_1}^T |M_2^T|_{a, 0}^T.
\end{aligned}$$

Then we set

$$R := M_1 M_2 - T = \sum_{i=1}^{m_1} M_1^{(i)} M_2^T + \sum_{i=1}^{m_2} M_1^T M_2^{(i)} + \sum_{i_1=1}^{m_1} \sum_{i_2=1}^{m_2} M_1^{(i_1)} M_2^{(i_2)} + (M_1^T M_2^T - T). \quad (\text{B.4})$$

We start with estimating the first summand. To each term in the sum we apply Lemma B.1 with $R = M_1^{(i)} \in \mathcal{L}_{\frac{a}{2}; -(i, m_1 - i)}$ and $T = M_2^T \in \mathcal{T}_{a, -m_2}$, deducing that

$$|M_1^{(i)} M_2^T|_{\frac{a}{2}; -(i, m_1 - i)} \lesssim a^{-p-2} |M_1^{(i)}|_{\frac{a}{2}; -(i, m_1 - i)} |M_2^T|_{a, \bar{0}}^T \stackrel{(\text{B.3})}{\lesssim} a^{-p-2} |M_1|_{a, -m_1}^{\text{qT}} |M_2|_{a, -m_2}^{\text{qT}}.$$

The second summand in (B.4) is estimated analogously using Lemma B.1 with $R = M_2^{(i)} \in \mathcal{L}_{\frac{a}{2}; -(i, m_2 - i)}$ and $T = M_1^T \in \mathcal{T}_{a, -m_1}$, getting

$$|M_1^T M_2^{(i)}|_{\frac{a}{2}; -(i, m_1)} \lesssim_{j, m_1} a^{-\mu i} |M_2^{(i)}|_{\frac{a}{2}; -(i, 0)} |M_1^T|_{a, -m_1}^T \stackrel{(\text{B.3})}{\lesssim_{m_1, m_2}} a^{-\mu m_2} |M_1|_{a, -m_1}^{\text{qT}} |M_2|_{a, -m_2}^{\text{qT}}.$$

Furthermore, by Lemma 2.14 one has that $\forall i_1 = 1, \dots, m_1$ and $\forall i_2 = 1, \dots, m_2$

$$|M_1^{(i_1)} M_2^{(i_2)}|_{\frac{a}{2}; -(i_1, m_1 - i_1)} \leq |M_1^{(i_1)}|_{\frac{a}{2}; -(i_1, m_1 - i_1)} |M_2^{(i_2)}|_{\frac{a}{2}; (0, 0)} \stackrel{(\text{B.3})}{\leq} 4 |M_1|_{a, -m_1}^{\text{qT}} |M_2|_{a, -m_2}^{\text{qT}}.$$

Remark that, since $m = \min\{m_1, m_2\}$, in particular $\forall i_1 = 1, \dots, m_1, \forall i_2 = 1, \dots, m_2$

$$M_1^{(i_1)} M_2^T \in \mathcal{L}_{\frac{a}{2}; -(i_1, m-i_1)}, \quad M_1^T M_2^{(i_2)} \in \mathcal{L}_{\frac{a}{2}; -(i_2, m)}, \quad M_1^{(i_1)} M_2^{(i_2)} \in \mathcal{L}_{\frac{a}{2}; -(i_1, m-i_1)}.$$

Thus it only remains to estimate the last summand in (B.4). In particular, we are going to prove that $M_1^T M_2^T - T \in \mathcal{L}_{\frac{a}{2}; -(m, 0)}$.

Recalling that $j - j' = \pi(\ell)$, we have:

$$(M_1^T M_2^T - T)_j^{j'}(\ell) = \sum_{\substack{\ell_1 + \ell_2 = \ell \\ j_1 = j - \pi(\ell_1)}} \mathfrak{F}_1(\ell_1, v(j), b(j)) \left(\mathfrak{F}_2(\ell_2, v(j_1), b(j_1)) - \mathfrak{F}_2(\ell_2, v(j), b(j) - v(j) \cdot \pi(\ell_1)) \right)$$

Now if $v(j) = v(j_1)$ then $b(j_1) := v(j_1) \cdot j_1 = v(j) \cdot (j - \pi(\ell_1)) = b(j) - v(j) \cdot \pi(\ell_1)$, and thus the term in the parenthesis above disappears. Consequently we can restrict the sum to the indexes j_1 fulfilling $v(j_1) \neq v(j)$.

For these remaining terms, using that $|j - j'| = |\pi(\ell)| \leq c^{-1}|\ell|$, we set $m' \leq m_1 + m_2$ and estimate

$$\begin{aligned} \left(\overline{M_1^T M_2^T - T}_{\frac{a}{2}; -(m', 0)} \right)_j^{j'} &\leq e^{-c\frac{a}{4}|j-j'|} \sum_{\substack{\ell_1, \ell_2 \\ j_1 = j - \pi(\ell_1), \\ v(j) \neq v(j_1)}} e^{\frac{3}{4}a|\ell_1 + \ell_2|} |\mathfrak{F}_1(\ell_1, v(j), b(j))| |\mathfrak{F}_2(\ell_2, v(j_1), b(j_1))| \langle j \rangle^{\mu m'} \\ &+ e^{-c\frac{a}{4}|j-j'|} \sum_{\substack{\ell_1, \ell_2 \\ j_1 = j - \pi(\ell_1), \\ v(j) \neq v(j_1)}} e^{\frac{3}{4}a|\ell_1 + \ell_2|} |\mathfrak{F}_1(\ell_1, v(j), b(j))| |\mathfrak{F}_2(\ell_2, v(j), b(j) - v(j) \cdot \pi(\ell_1))| \langle j \rangle^{\mu m'}. \end{aligned} \tag{B.5}$$

If we show that the two sums above are uniformly bounded, then Lemma 2.9 will give us the needed boundedness. Let us start with the first sum. We distinguish two cases.

Case 1: $\langle j \rangle \leq J_\delta$ (given by Lemma 2.12). Then trivially

$$\sum_{\substack{\ell_1, \ell_2 \\ j_1 = j - \pi(\ell_1), \\ v(j) \neq v(j_1)}} e^{\frac{3}{4}a|\ell_1 + \ell_2|} |\mathfrak{F}_1(\ell_1, v(j), b(j))| |\mathfrak{F}_2(\ell_2, v(j_1), b(j_1))| \langle j \rangle^{\mu m'} \lesssim_\delta |M_1^T|_{a,0}^T |M_2^T|_{a,0}^T.$$

Case 2: $\langle j \rangle > J_\delta$. We further distinguish 4 subcases:

(A) $|\ell_1| > c\langle j \rangle^\delta$; (B) $|\ell_1| \leq c\langle j \rangle^\delta$ and $|v(j_1)| > \frac{1}{2}\langle j \rangle^\delta$; (C) $|\ell_1| \leq c\langle j \rangle^\delta$, $|v(j_1)| \leq \frac{1}{2}\langle j \rangle^\delta$ and $|b(j)| > \langle j \rangle^\mu$; (D) $|\ell_1| \leq c\langle j \rangle^\delta$, $|v(j_1)| \leq \frac{1}{2}\langle j \rangle^\delta$ and $|b(j)| \leq \langle j \rangle^\mu$.

In case (A) we have

$$\begin{aligned} &\sum_{\substack{\ell_1, \ell_2 \\ j_1 = j - \pi(\ell_1), \\ v(j) \neq v(j_1)}}^{(A)} e^{\frac{3}{4}a|\ell_1 + \ell_2|} |\mathfrak{F}_1(\ell_1, v(j), b(j))| |\mathfrak{F}_2(\ell_2, v(j_1), b(j_1))| \langle j \rangle^{\mu m'} \\ &\lesssim_{c, \mu} \sum_{\ell_1} e^{\frac{3}{4}a|\ell_1|} |\ell_1|^{\frac{\mu m'}{\delta}} |\mathfrak{F}_1(\ell_1, v(j), b(j))| |M_2^T|_{a,0}^T \lesssim_{c, \mu, \delta} a^{-\frac{\mu m'}{\delta}} |M_1^T|_{a,0}^T |M_2^T|_{a,0}^T. \end{aligned}$$

In case (B) we get

$$\begin{aligned}
& \sum_{\substack{\ell_1, \ell_2 \\ j_1 = j - \pi(\ell_1), \\ v(j) \neq v(j_1)}}^{(B)} e^{\frac{3}{4}a|\ell_1 + \ell_2|} |\mathfrak{F}_1(\ell_1, v(j), b(j))| |\mathfrak{F}_2(\ell_2, v(j_1), b(j_1))| \langle j \rangle^{\mu m'} \\
& \lesssim_{\mu, \delta} |M_1^{\mathbb{T}}|_{a,0}^{\mathbb{T}} \sum_{\ell_2} e^{\frac{3}{4}a|\ell_2|} |\mathfrak{F}_2(\ell_2, v(j_1), b(j_1))| |v(j_1)|^{\frac{\mu m'}{\delta}} \lesssim_{\mu, \delta, c} a^{-\frac{\mu m'}{\delta}} |M_1^{\mathbb{T}}|_{a,0}^{\mathbb{T}} |M_2^{\mathbb{T}}|_{a,0}^{\mathbb{T}}.
\end{aligned}$$

In case (C) we start by remarking that $|b(j_1)| > \frac{1}{2}\langle j \rangle^\mu$ provided J_δ is large enough. Indeed assume by contradiction that $|b(j_1)| \leq \frac{1}{2}\langle j \rangle^\mu$. Then, by the definition of $b(j)$, we have that

$$|b(j)| \leq |v(j_1) \cdot j| \leq |v(j_1) \cdot j_1| + |v(j_1) \cdot \pi(\ell_1)| \leq |b(j_1)| + \langle j \rangle^{2\delta} \leq \frac{1}{2}\langle j \rangle^\mu + \langle j \rangle^{2\delta} \leq \langle j \rangle^\mu,$$

where in the last inequality we have used $\langle j \rangle > 2^{\frac{1}{1-4\delta}}$, contradicting the hypotheses. Now, as $\mu \leq 1$, we bound

$$\begin{aligned}
& \sum_{\substack{\ell_1, \ell_2 \\ j_1 = j - \pi(\ell_1), \\ v(j) \neq v(j_1)}}^{(C)} e^{\frac{3}{4}a|\ell_1 + \ell_2|} |\mathfrak{F}_1(\ell_1, v(j), b(j))| |\mathfrak{F}_2(\ell_2, v(j_1), b(j_1))| \langle j \rangle^{m_1 \mu} \langle j \rangle^{m_2 \mu} \\
& \lesssim \sum_{\ell_1} e^{\frac{3}{4}a|\ell_1|} |\mathfrak{F}_1(\ell_1, v(j), b(j))| \langle b(j) \rangle^{m_1} \sum_{\ell_2} e^{\frac{3}{2}|\ell_2|} |\mathfrak{F}_2(\ell_2, v(j_1), b(j_1))| \langle b(j_1) \rangle^{m_2} \\
& \lesssim_{\mu, \delta} a^{-\frac{\mu(m_1 + m_2)}{\delta}} |M_2^{\mathbb{T}}|_{a, -m_1}^{\mathbb{T}} |M_1^{\mathbb{T}}|_{a, -m_2}^{\mathbb{T}}.
\end{aligned}$$

Concerning (D), we claim that in this case $v(j) = v(j_1)$, so there is no contribution to the sum. Indeed if $v(j_1) \neq v(j)$, Lemma 2.12 (ii) (which we can apply as $|j - j_1| \leq \langle j \rangle^\delta$) would imply $|v(j_1)| > \frac{1}{2}\langle j \rangle^\delta$, getting a contradiction. This concludes the proof of (2.19).

Proof of (2.20): Let $M_i^{\mathbb{T}} \in \mathcal{T}_{a,-1}$ and $M_i^{(1)} \in \mathcal{L}_{a;-(1,0)}$ be quasi-Töplitz decompositions of the M_i .

$$|M_i^{\mathbb{T}}|_{a,-1}^{\mathbb{T}} + |M_i^{(1)}|_{a;-(1,0)} \leq (1 + \varepsilon) |M_i|_{a,-1}^{\mathbb{qT}}.$$

Arguing as in the proof of Proposition 2.21, to each $M_i^{\mathbb{T}}$ we associate the symbol $\mathfrak{F}_i(\ell, v, b)$. We set

$$\mathfrak{F}(\ell, v, b) := \sum_{\ell_1 + \ell_2 = \ell} \mathfrak{F}_1(\ell_1, v, b) \mathfrak{F}_2(\ell_2, v, b - v \cdot \pi(\ell_1)), \quad (T)_j^{j'}(\ell) := \mathfrak{F}(\ell, v(j), b(j)),$$

and we define the line-Toplitz part of $M_1 M_2$ as $(M_1 M_2)^{\mathbb{T}} = T$. Then T is by definition line-Töplitz, moreover to estimate $|T|_{a', -2}^{\mathbb{T}} \forall a' < a$ we only need to control:

$$\begin{aligned}
& \sup_{v \in \mathcal{V}, b \in \mathbb{Z}} \sum_{\ell \in \mathbb{Z}^d} e^{a'(c|v| + |\ell|)} \left| \sum_{\ell_1 + \ell_2 = \ell} \mathfrak{F}_1(\ell_1, v, b) \mathfrak{F}_2(\ell_2, v, b - v \cdot \pi(\ell_1)) \right| \langle b \rangle^2 \\
& \lesssim \sup_{v \in \mathcal{V}, b \in \mathbb{Z}} \sum_{\ell_1 \in \mathbb{Z}^d} e^{a'(c|v| + |\ell_1|)} |\mathfrak{F}_1(\ell_1, v, b)| \langle b \rangle \langle v \cdot \pi(\ell_1) \rangle \sup_{v' \in \mathcal{V}, b' \in \mathbb{Z}} \sum_{\ell_2 \in \mathbb{Z}^d} e^{a'(c|v'| + |\ell_2|)} |\mathfrak{F}_2(\ell_2, v', b')| \langle b' \rangle \\
& \lesssim \sup_{v \in \mathcal{V}, b \in \mathbb{Z}} \sum_{\ell_1 \in \mathbb{Z}^d} e^{a(|\ell_1| + c|v|)} |\mathfrak{F}_1(\ell_1, v, b)| \langle b \rangle e^{-(a-a')c|v| - (a-a')|\ell_1|} \langle v \rangle \langle \ell_1 \rangle |M_2^{\mathbb{T}}|_{a', -1}^{\mathbb{T}} \\
& \lesssim_c (a - a')^{-2} |M_1^{\mathbb{T}}|_{a, -1}^{\mathbb{T}} |M_2^{\mathbb{T}}|_{a', -1}^{\mathbb{T}}.
\end{aligned}$$

We are left with estimating

$$M_1 M_2 - T = M_1^T M_2^{(1)} + M_1^{(1)} M_2^T + M_1^{(1)} M_2^{(1)} + (M_1^T M_2^T - T). \quad (\text{B.6})$$

We will actually prove the following:

$$M_1^T M_2^{(1)} \in \mathcal{L}_{\frac{a'}{2}; -(1,1)}, \quad M_1^{(1)} M_2^T \in \mathcal{L}_{\frac{a'}{2}; -(1,1)} + \mathcal{L}_{\frac{a'}{2}; -(2,0)}, \quad M_1^{(1)} M_2^{(1)} + (M_1^T M_2^T - T) \in \mathcal{L}_{\frac{a'}{2}; -(2,0)}.$$

The estimate of the last two terms in (B.6) is straightforward: by Formula (2.14) in Lemma 2.14 one has $M_1^{(1)} M_2^{(1)} \in \mathcal{L}_{\frac{a'}{2}; -(2,0)}$, with

$$|M_1^{(1)} M_2^{(1)}|_{\frac{a'}{2}; -(2,0)} \lesssim_{\mu, \delta} (a - a')^{-\frac{2\mu}{\delta}} |M_1^{(1)}|_{\frac{a'}{2}; -(1,0)} |M_2^{(1)}|_{\frac{a'}{2}; -(1,0)},$$

while arguing as in the proof of Proposition 2.21, formula (B.4) with $m' = 2$, one obtains that there exists $\gamma = \gamma(\mu, \delta, p) > 0$ such that $T_1 T_2 - T \in \mathcal{L}_{\frac{a'}{2}; -(2,0)}$, with

$$|M_1^T M_2^T - T|_{\frac{a'}{2}; -(2,0)} \lesssim_{\mu, \delta, c, p} a^{-\gamma} |M_1^T|_{a, -1}^T |M_2^T|_{a, -1}^T.$$

Concerning $M_1^T M_2^{(1)}$, one has:

$$\begin{aligned} \left(\underline{M_1^T M_2^{(1)}}_{\frac{a'}{2}; -(1,1)} \right)_j^{j'} &\lesssim_{\mu} \sum_{\substack{\ell_1, \ell_2: \\ j-j'=\pi(\ell_1+\ell_2)}} e^{\frac{a'}{2}|\ell_1+\ell_2|} |(M_1^T)_j^{j-\pi(\ell_1)}(\ell_1)| \langle j-\pi(\ell_1) \rangle^{\mu} \langle \pi(\ell_1) \rangle^{\mu} \langle b(j) \rangle | (M_2^{(1)})_{j-\pi(\ell_1)}^{j'}(\ell_2) | \\ &\lesssim_{\mu} a^{-\mu} \sum_{j_1} \left(\underline{M_1^T} \right)_{\frac{3}{4}a; -(0,1)}^{j_1} \left(\underline{M_2^{(1)}} \right)_{\frac{a'}{2}; -(1,0)}^{j_1}, \end{aligned}$$

which implies

$$|M_1^T M_2^{(1)}|_{\frac{a'}{2}; -(1,1)} \lesssim_{\mu} a^{-\mu} |M_1^T|_{a, -1}^T |M_2^{(1)}|_{\frac{a'}{2}; -(1,0)},$$

by Lemma 2.14 and by Lemma 2.17.

In order to conclude the proof, we now estimate $M_1^{(1)} M_2^T$. Recalling that one has

$$(M_1^{(1)} M_2^T)_j^{j'}(\ell) = \sum_{\ell_1, \ell_2: \ell_1 + \ell_2 = \ell} (M_1^{(1)})_j^{j-\pi(\ell_1)}(\ell_1) (M_2^T)_{j-\pi(\ell_1)}^{j'}(\ell_2), \quad (\text{B.7})$$

with $(M_1^{(1)} M_2^T)_j^{j'} \neq 0$ only if $j - j' = \pi(\ell)$, we decompose $M_1^{(1)} M_2^T$ as follows:

$$M_1^{(1)} M_2^T = (M_1^{(1)} M_2^T)^{(A)} + (M_1^{(1)} M_2^T)^{(B)} + (M_1^{(1)} M_2^T)^{(C)} + (M_1^{(1)} M_2^T)^{(D)} + (M_1^{(1)} M_2^T)^{(E)}, \quad (\text{B.8})$$

where the summands in (B.8) respectively corresponds to the cases:

- (A) $\langle j - \pi(\ell_1) \rangle > J_{\delta}$, with J_{δ} given by Lemma 2.12, and $|\ell_1| > c \langle j \rangle^{\delta}$;
- (B) $\langle j - \pi(\ell_1) \rangle > J_{\delta}$, $|\ell_1| \leq c \langle j \rangle^{\delta}$ and $|v(j - \pi(\ell_1))| > \frac{1}{2} \langle j - \pi(\ell_1) \rangle^{\delta}$;
- (C) $\langle j - \pi(\ell_1) \rangle > J_{\delta}$, $|\ell_1| \leq c \langle j \rangle^{\delta}$, $|v(j - \pi(\ell_1))| \leq \frac{1}{2} \langle j - \pi(\ell_1) \rangle^{\delta}$, and $|b(j - \pi(\ell_1))| > \langle j - \pi(\ell_1) \rangle^{\mu}$;
- (D) $\langle j - \pi(\ell_1) \rangle > J_{\delta}$, $|\ell_1| \leq c \langle j \rangle^{\delta}$, $|v(j - \pi(\ell_1))| \leq \frac{1}{2} \langle j - \pi(\ell_1) \rangle^{\delta}$ and $|b(j - \pi(\ell_1))| \leq \langle j - \pi(\ell_1) \rangle^{\mu}$;
- (E) $\langle j - \pi(\ell_1) \rangle \leq J_{\delta}$.

Concerning $(M_1^{(1)} M_2^T)^{(A)}$, using $|\ell_1| > \mathbf{c}\langle j \rangle^\delta$ one obtains

$$\left(\underline{(M_1^{(1)} M_2^T)}_{\frac{a'}{2}; -(2,0)} \right)_j^{j'} \lesssim_{\mathbf{c}, \mu, \delta} (a - a')^{-\frac{2\mu}{\delta}} \sum_{j_1} \left(\underline{(M_1^{(1)})}_{\frac{a}{2}, 0, 0} \right)_j^{j_1} \left(\underline{(M_2^T)}_{\frac{a}{2}, 0, 0} \right)_{j_1}^{j'}.$$

For $(M_1^{(1)} M_2^T)^{(B)}$, we define $\theta := \mathbf{c}^{-1}(a - a')/2$ and we observe that

$$\langle j \rangle^\delta \lesssim_\delta \langle j - \pi(\ell_1) \rangle^\delta + \langle \pi(\ell_1) \rangle^\delta \lesssim_\delta \langle j - \pi(\ell_1) \rangle^\delta \lesssim_\delta |v(j - \pi(\ell_1))|,$$

from which one obtains

$$\left(\underline{(M_1^{(1)} M_2^T)}_{\frac{a'}{2}; -(2,0)} \right)_j^{j'} \lesssim_{\mathbf{c}, \mu, \delta} a^{-\frac{2\mu}{\delta}} e^{-\theta|j-j'|} |M_1^{(1)}|_{\frac{a}{2}; (0,0)} |M_2^T|_{\frac{a}{2}, 0}.$$

Then the estimate on $(M_1^{(1)} M_2^T)^{(B)}$ follows by Lemma 2.9. The estimate of $(M_1^{(1)} M_2^T)^{(C)}$ simply follows observing that, since

$$\langle j \rangle^\mu \lesssim_\mu \langle j - \pi(\ell_1) \rangle^\mu + \langle \pi(\ell_1) \rangle^\mu \lesssim_\mu \langle j - \pi(\ell_1) \rangle^\mu \lesssim_\mu |b(j - \pi(\ell_1))|,$$

one has

$$\begin{aligned} \left(\underline{(M_1^{(1)} M_2^T)}_{\frac{a'}{2}; -(2,0)} \right)_j^{j'} &\leq \sum_{\substack{\ell_1, \ell_2: j-j'=\pi(\ell_1+\ell_2), \\ |b(j-\pi(\ell_1))| \geq \langle j \rangle^\mu}} e^{a'|\ell_1+\ell_2|} \langle j \rangle^{2\mu} |(M_1^{(1)})_j^{j-\pi(\ell_1)}(\ell_1) (M_2^T)_{j-\pi(\ell_1)}^{j'}(\ell_2)| \\ &\leq \sum_{\ell_1, \ell_2: j-j'=\pi(\ell_1+\ell_2)} e^{\frac{a'}{2}|\ell_1+\ell_2|} \langle j \rangle^\mu \langle b(j - \pi(\ell_1)) \rangle |(M_1^{(1)})_j^{j-\pi(\ell_1)}(\ell_1) (M_2^T)_{j-\pi(\ell_1)}^{j'}(\ell_2)| \\ &\leq \sum_{j_1} \left(\underline{(M_1^{(1)})}_{\frac{a'}{2}, -(1,0)} \right)_j^{j_1} \left(\underline{(M_2^T)}_{\frac{a'}{2}, -(0,1)} \right)_j^{j_1}, \end{aligned}$$

and applying Lemma 2.14 and Lemma 2.17. In order to estimate $(M_1^{(1)} M_2^T)^{(D)}$, we start with recalling that, by Item 2 of Lemma 2.12 with j replaced by $j - \pi(\ell_1)$ and h replaced by j , one has $v(j - \pi(\ell_1)) = v(j)$. Then one has

$$\begin{aligned} |b(j)| &= |j \cdot v(j - \pi(\ell_1))| \leq |(j - \pi(\ell_1)) \cdot v(j - \pi(\ell_1))| + |\pi(\ell_1)| |v(j - \pi(\ell_1))| \\ &\leq \langle b(j - \pi(\ell_1)) \rangle + \mathbf{c}^{-1} |\ell_1| |v(j - \pi(\ell_1))| \\ &\lesssim_{\mathbf{c}} \langle b(j - \pi(\ell_1)) \rangle + \langle \ell_1 \rangle \langle v(j - \pi(\ell_1)) \rangle. \end{aligned}$$

This in turn, setting $\sigma := (a - a')/2$ and $\theta := \mathbf{c}^{-1}\sigma$, enables to deduce

$$\begin{aligned} &\left(\underline{(M_1^{(1)} M_2^T)}_{\frac{a'}{2}; -(1,1)} \right)_j^{j'} \\ &\lesssim_{\mathbf{c}} \sum_{\substack{\ell_1, \ell_2: \\ j-j'=\pi(\ell_1+\ell_2)}} e^{\frac{a'}{2}|\ell_1+\ell_2|} |(M_1^{(1)})_j^{j-\pi(\ell_1)}(\ell_1)| \langle j \rangle^\mu \langle b(j - \pi(\ell_1)) \rangle |\ell_1| |v(j - \pi(\ell_1))| |(M_2^T)_{j-\pi(\ell_1)}^{j'}(\ell_2)| \\ &\lesssim_{\mathbf{c}} (a\sigma)^{-1} e^{-\frac{\theta}{2}|j-j'|} |M_1^{(1)}|_{\frac{a'}{2}; -(1,0)} |M_2^T|_{a, -1}. \end{aligned}$$

Finally, using that $\langle j \rangle^{2\mu} \lesssim_\mu \langle j - \pi(\ell_1) \rangle^{2\mu} + \langle \pi(\ell_1) \rangle^{2\mu} \lesssim_\mu \mu, \delta, \mathbf{c} J_\delta^{2\mu} |\ell_1|^{2\mu}$, one obtains

$$\left(\underline{(M_1^{(1)} M_2^T)}_{\frac{a'}{2}; -(2,0)} \right)_j^{j'} \lesssim_{\delta, \mathbf{c}, \mu} (a - a')^{-2\mu} \sum_{j_1} \left(\underline{(M_1^{(1)})}_{\frac{a}{2}; (0,0)} \right)_j^{j_1} \left(\underline{(M_2^T)}_{\frac{a'}{2}; (0,0)} \right)_{j_1}^{j'}.$$

□

Proof of Lemma 4.4. We set (recall that $V(\ell) = 0$ if $\pi(\ell) = 0$) $(S_0^T)^{j'}(\ell) = \mathfrak{S}(\ell, v(j), b(j))$ with

$$\mathfrak{S}(\ell, v, b) = \begin{cases} \frac{-V(\ell)}{\omega \cdot \ell + 2 \frac{|\pi(\ell)|}{|v|} b - |\pi(\ell)|^2}, & \text{if } \pi(\ell) \neq 0, \quad v \parallel \pi(\ell) \\ 0 & \text{otherwise} \end{cases}. \quad (\text{B.9})$$

One has

$$\begin{aligned} |S_0^T|_{a', -1}^T &\leq \sup_{v \in \mathcal{V}, b \in \mathbb{Z}} \sum_{\ell \in \mathbb{Z}^d} e^{a'c|v|+a'|\ell|} |\mathfrak{S}(\ell, v, b)| \langle b \rangle^1 \\ &= \sup_{v \in \mathcal{V}, b \in \mathbb{Z}} \sum_{\ell \in \mathbb{Z}^d} e^{a'c|v|+a'|\ell|} \frac{|V(\ell)|}{|\omega \cdot \ell + 2 \frac{|\pi(\ell)|}{|v|} b - |\pi(\ell)|^2|} \langle b \rangle^1. \end{aligned}$$

We estimate $*$ = $\frac{\langle b \rangle}{|\omega \cdot \ell - |\pi(\ell)|^2 + 2 \frac{|\pi(\ell)|}{|v|} b|}$ by distinguishing two cases. If $|b| > |\omega \cdot \ell - |\pi(\ell)|^2|$ then (recalling that $\frac{|\pi(\ell)|}{|v|}$ is a non-zero integer) one has $*$ < 2 . Otherwise if $|b| \leq |\omega \cdot \ell - |\pi(\ell)|^2|$ then, by (4.3) and recalling that $|\pi(\ell)| \leq c^{-1}|\ell|$,

$$* \leq (|\omega \cdot \ell - |\pi(\ell)|^2| + 1) |\ell|^\tau \gamma^{-1} \lesssim \gamma^{-1} |\ell|^{\tau+2}.$$

In conclusion,

$$\begin{aligned} |S_0^T|_{a', -1}^T &\lesssim \gamma^{-1} \sum_{\ell: j-j'=\pi(\ell) \neq 0} e^{a'c|\pi(\ell)|+a'|\ell|} |V(\ell)| |\ell|^{(\tau+2)} \\ &\lesssim \gamma^{-1} \sum_{\ell: j-j'=\pi(\ell) \neq 0} e^{2a'|\ell|} |V(\ell)| |\ell|^{(\tau+2)} \lesssim_{c, \gamma} (\mathbf{a} - 2a')^{-(\tau+2)} \|V\|_{\mathbf{a}, \mathbf{p}}. \end{aligned} \quad (\text{B.10})$$

Recalling that, if $\pi(\ell) \parallel v(j)$, $j \cdot \pi(\ell) = \frac{|\pi(\ell)|}{|v(j)|} b(j)$, we have

$$(S_0 - S_0^T)^{j'}(\ell) = \begin{cases} \frac{-V(\ell)}{\omega \cdot \ell + 2j \cdot \pi(\ell) - |\pi(\ell)|^2}, & j' = j - \pi(\ell), \quad \pi(\ell) \neq 0, \quad \pi(\ell) \not\parallel v(j), \\ 0 & \text{otherwise.} \end{cases}$$

Accordingly, we divide the indexes j, ℓ which contribute to $S_0 - S_0^T$ following the two conditions: i) One either has $|\ell| \geq c\langle j \rangle^\delta$ or $\langle j \rangle \leq j_{\min} := 2^{\frac{1}{2\delta}}$; ii) One has $|\ell| < c\langle j \rangle^\delta$, $\langle j \rangle > j_{\min}$ and $\pi(\ell) \not\parallel v(j)$. We decompose accordingly $S_0 - S_0^T = R + P$.

First, by applying Lemma A.1 to $S_0 - S_0^T$ and setting $R_1 = (S_0 - S_0^T)^{(R)}$, we deduce

$$|R_1|_{\frac{a'}{2}; -(2,0)} \lesssim_\delta \mathbf{a}^{-\frac{2\mu}{\delta}} \|V\|_{\mathbf{a}, \mathbf{p}}.$$

To deal with the case $|j| \leq j_{\min}$, we set $(R_2)^{j'}(\ell) := (S_0)^{j'}(\ell)$ if $j \leq j_{\min}$. Since if $|j| \leq j_{\min}$ we have

$$\frac{\langle j \rangle^{2\mu}}{|\omega \cdot \ell + 2j \cdot \pi(\ell) - |\pi(\ell)|^2|} \leq \gamma^{-1} \langle j_{\min} \rangle^{2\mu} |\ell|^\tau, \quad (\text{B.11})$$

defining $R := R_1 + R_2$, we get that

$$|R|_{\frac{a'}{2}; -(2,0)} \lesssim_{\delta, \tau} \mathbf{a}^{-\max\{\tau, \frac{2\mu}{\delta}\}} \|V\|_{\mathbf{a}, \mathbf{p}}. \quad (\text{B.12})$$

To deal with case ii), we define

$$P_j^{j'}(\ell) := \begin{cases} \frac{-V(\ell)}{\omega \cdot \ell + 2j \cdot \pi(\ell) - |\pi(\ell)|^2} & \text{if } j > j_{\min}, \quad |\ell| < c\langle j \rangle^\delta \quad \text{and } \pi(\ell) \not\parallel v(j), \\ 0 & \text{otherwise} \end{cases}.$$

Now we claim that if $\langle j \rangle \geq 2^{\frac{1}{2\delta}}$, $|\ell| < c\langle j \rangle^\delta$ and $\pi(\ell) \not\parallel v(j)$, then $|j \cdot \pi(\ell)| > \langle j \rangle^\mu$. Indeed, $|\ell| < c\langle j \rangle^\delta$ implies that $|\pi(\ell)| < \langle j \rangle^\delta$. Fixing $w \in \mathcal{V}$ to be the direction parallel to $\pi(\ell)$ one has $|w| < \langle j \rangle^\delta$ hence

$$|\pi(\ell) \cdot j| \geq |w \cdot j| \geq |b(j)| := \min_{v \in \mathcal{V}: |v| < \langle j \rangle^\delta} |j \cdot v|$$

Then, if $|b(j)| > \langle j \rangle^\mu$ our claim follows. Otherwise, we prove the claim by contradiction. Recall that $v(j)$ attains the minimum above and by ii) $v(j) \not\parallel w$. If $|w \cdot j| \leq \langle j \rangle^\mu$ then, by Lemma 2.11 $\langle j \rangle < \sqrt{2}\langle j \rangle^{1-\delta}$, which contradicts $\langle j \rangle \geq 2^{\frac{1}{2\delta}}$.

As a consequence if $\langle j \rangle \geq 2^{\frac{1}{2\delta}}$, $|\ell| < c\langle j \rangle^\delta$ and $\pi(\ell) \not\parallel v(j)$, then

$$\frac{\langle j \rangle^\mu}{|\omega \cdot \ell + 2j \cdot \pi(\ell) - |\pi(\ell)|^2|} \leq \frac{\langle j \rangle^\mu}{2|j \cdot \pi(\ell)| - |\omega \cdot \ell - |\pi(\ell)|^2|} \leq \frac{\langle j \rangle^\mu}{2\langle j \rangle^\mu - (|\omega| + c^{-2})c^2\langle j \rangle^{2\delta}} \leq 1$$

provided that $\langle j \rangle \geq j_{\min}$. Therefore we have

$$|P|_{\frac{\alpha'}{2}; -(1,0)} \lesssim \|V\|_{\mathbf{a},\mathbf{p}}. \quad (\text{B.13})$$

Recalling that $S_0 = S_0^\Gamma + P + R$, estimates (B.10), (B.12), (B.13) then imply that S is quasi-Töplitz of order -1 . We now prove estimate (4.10). We start with explicitly computing $[S_0, M_V]$. Setting $h = j - \pi(\ell)$, one has

$$\begin{aligned} ([S_0, M_V]_j^h)(\ell) &= \sum_{\ell_1 + \ell_2 = \ell} V(\ell_1) ((S_0)_j^{j - \pi(\ell_2)}(\ell_2) - (S_0)_{j - \pi(\ell_1)}^{j - \pi(\ell)}(\ell_2)) \\ &= \sum_{\ell_1 + \ell_2 = \ell} V(\ell_1) V(\ell_2) \left(\frac{1}{\omega \cdot \ell_2 + 2(j - \pi(\ell_1)) \cdot \pi(\ell_2) - |\pi(\ell_2)|^2} - \frac{1}{\omega \cdot \ell_2 + 2j \cdot \pi(\ell_2) - |\pi(\ell_2)|^2} \right) \\ &= \sum_{\ell_1 + \ell_2 = \ell} \frac{-2(\pi(\ell_1) \cdot \pi(\ell_2)) V(\ell_1) V(\ell_2)}{(\omega \cdot \ell_2 + 2j \cdot \pi(\ell_2) - |\pi(\ell_2)|^2)(\omega \cdot \ell_2 + 2(j - \pi(\ell_1)) \cdot \pi(\ell_2) - |\pi(\ell_2)|^2)}. \end{aligned}$$

We then split the above sum into two terms, according to whether $\pi(\ell_2) \parallel v(j)$ or not. In the first case, we set

$$\begin{aligned} ([S_0, M_V]_j^T)^{j - \pi(\ell)}(\ell) &:= \sum_{\substack{\ell_1 + \ell_2 = \ell \\ \pi(\ell_2) \parallel v(j)}} \frac{-(2\pi(\ell_1) \cdot \pi(\ell_2)) V(\ell_1) V(\ell_2)}{(\omega \cdot \ell_2 + 2j \cdot \pi(\ell_2) - |\pi(\ell_2)|^2)(\omega \cdot \ell_2 + 2(j - \pi(\ell_1)) \cdot \pi(\ell_2) - |\pi(\ell_2)|^2)} \\ &= \sum_{\substack{\ell_1 + \ell_2 = \ell \\ \pi(\ell_2) \parallel v(j)}} \frac{-(2\pi(\ell_1) \cdot \pi(\ell_2)) V(\ell_1) V(\ell_2)}{(\omega \cdot \ell_2 + 2b(j) \frac{|\pi(\ell_2)|}{|v(j)|} - |\pi(\ell_2)|^2)(\omega \cdot \ell_2 + 2b(j) \frac{|\pi(\ell_2)|}{|v(j)|} - 2\pi(\ell_1) \cdot \pi(\ell_2) - |\pi(\ell_2)|^2)}, \end{aligned} \quad (\text{B.14})$$

and we are going to prove that its symbol

$$\mathfrak{I}(\ell, b, v) := \sum_{\substack{\ell_1 + \ell_2 = \ell \\ \pi(\ell_2) \parallel v}} \frac{-(2\pi(\ell_1) \cdot \pi(\ell_2)) V(\ell_1) V(\ell_2)}{(\omega \cdot \ell_2 + 2b(j) \frac{|\pi(\ell_2)|}{|v|} - |\pi(\ell_2)|^2)(\omega \cdot \ell_2 + 2b \frac{|\pi(\ell_2)|}{|v|} - 2\pi(\ell_1) \cdot \pi(\ell_2) - |\pi(\ell_2)|^2)}$$

is of order -2 . In the second case, we set

$$R_j^h(\ell) := \sum_{\substack{\ell_1 + \ell_2 = \ell \\ \pi(\ell_2) \parallel v(j)}} \frac{-(2\pi(\ell_1) \cdot \pi(\ell_2))V(\ell_1)V(\ell_2)}{(\omega \cdot \ell_2 + 2j \cdot \pi(\ell_2) - |\pi(\ell_2)|^2)(\omega \cdot \ell_2 + 2(j - \pi(\ell_1)) \cdot \pi(\ell_2) - |\pi(\ell_2)|^2)}, \quad (\text{B.15})$$

with $h = j - \pi(\ell)$, and we shall prove that $R \in \mathcal{L}_{\frac{a'}{2}; -(2,0)}$.

We start with exhibiting estimates on $[[S_0, M_V]^T]_{a', -2}^T$. Let

$$f^\epsilon(\ell_1, \ell_2) := \omega \cdot \ell_2 - |\pi(\ell_2)|^2 - 2\epsilon\pi(\ell_1) \cdot \pi(\ell_2), \quad \epsilon \in \{0, 1\}, \quad (\text{B.16})$$

so that the denominators appearing in (B.15) have the form

$$(2j \cdot \pi(\ell_2) + f^0(\ell_1, \ell_2))(2j \cdot \pi(\ell_2) + f^1(\ell_1, \ell_2));$$

arguing as to obtain estimates (B.10), we observe that $\forall \epsilon \in \{0, 1\}$, if $\langle b \rangle \geq |f^\epsilon(\ell_1, \ell_2)|$, one has

$$\left| \frac{2\langle b \rangle}{(\omega \cdot \ell_2 + 2b(j) \frac{|\pi(\ell_2)|}{|v(j)|} - 2\epsilon\pi(\ell_1) \cdot \pi(\ell_2) - |\pi(\ell_2)|^2)} \right| \leq 2, \quad (\text{B.17})$$

while if $\langle b \rangle < |f^\epsilon(\ell_1, \ell_2)|$, one has

$$\begin{aligned} & \left| \frac{2\langle b \rangle}{(\omega \cdot \ell_2 + 2b(j) \frac{|\pi(\ell_2)|}{|v(j)|} - 2\epsilon\pi(\ell_1) \cdot \pi(\ell_2) - |\pi(\ell_2)|^2)} \right| \leq \\ & \frac{2|f^\epsilon(\ell_1, \ell_2)|}{\left| (\omega \cdot \ell_2 + 2b(j) \frac{|\pi(\ell_2)|}{|v(j)|} - 2\epsilon\pi(\ell_1) \cdot \pi(\ell_2) - |\pi(\ell_2)|^2) \right|} \lesssim \frac{|\ell_2|^2 |\ell_1|}{\gamma |\ell_2|^{-\tau}} \lesssim \gamma^{-1} |\ell_2|^{\tau+2} |\ell_1|. \end{aligned} \quad (\text{B.18})$$

Let now $\theta := (\mathbf{a} - 2a')/2$. Combining (B.17) and (B.18) and recalling that, since $\pi(\ell_2) \parallel v$ and v is a generator, one has $c|v| \leq c|\pi(\ell_2)| \leq |\ell_2|$, one obtains

$$\begin{aligned} & [[S_0, M_V]^T]_{a', -2}^T \lesssim \sum_{\ell} e^{a'c|v| + a'|\ell|} \sum_{\substack{\ell_1 + \ell_2 = \ell \\ \pi(\ell_2) \parallel v}} |V(\ell_1)V(\ell_2)| |\ell_2|^{2(\tau+2)+1} |\ell_1|^3 \\ & \lesssim \sum_{\ell} e^{a'|\ell|} \sum_{\substack{\ell_1 + \ell_2 = \ell \\ \pi(\ell_2) \parallel v}} e^{a'|\ell_2|} |V(\ell_1)V(\ell_2)| |\ell_2|^{2(\tau+2)+1} |\ell_1|^3 \\ & \lesssim \sum_{\ell_1, \ell_2} e^{a'|\ell_1| + 2a'|\ell_2|} |\ell_2|^{2(\tau+2)+1} |\ell_1|^3 |V(\ell_1)| |V(\ell_2)| \lesssim_{c, \tau} (\mathbf{a} - 2a')^{-2(\tau+2)-4} \|V\|_{\mathbf{a}, \mathbf{p}}^2, \end{aligned}$$

provided $2a' \leq \mathbf{a}$.

We now turn to prove that R defined as in (B.15) belongs to $\mathcal{L}_{\frac{a'}{2}; -(2,0)}$. Letting $j_{\min} := 2^{\frac{1}{2\delta}}$, we decompose $R = R_1 + R_2 + R_3$ as follows:

$$(R_1)_j^h(\ell) = R_j^h(\ell) \quad \text{if } \max\{|\ell|, |\ell_1|, |\ell_2|\} > c\langle j \rangle^\delta, \quad (R_1)_j^h(\ell) = 0 \quad \text{otherwise};$$

$$(R_2)_j^h(\ell) = R_j^h(\ell) \quad \text{if } \langle j \rangle \leq j_{\min}, \quad (R_2)_j^h(\ell) = 0 \quad \text{otherwise};$$

$$(R_3)_j^h(\ell) = R_j^h(\ell) \quad \text{if } \max\{|\ell|, |\ell_1|, |\ell_2|\} \leq c\langle j \rangle^\delta \quad \text{and } \langle j \rangle \geq j_{\min}, \quad (R_3)_j^h(\ell) = 0 \quad \text{otherwise}.$$

Arguing as to prove estimate (B.12), one easily sees that $R_1, R_2 \in \mathcal{L}_{\frac{a'}{2}; -(N,0)}$ for any $N \in \mathbb{N}$. We

now prove that $R_3 \in \mathcal{L}_{\frac{a'}{2}; -(2,0)}$.

First of all, we claim the following:

$$(R_3)_j^h(\ell) \neq 0 \quad \implies \quad |j \cdot \pi(\ell_2)| > \langle j \rangle^\mu. \quad (\text{B.19})$$

Indeed, if $(R_3)_j^h(\ell) \neq 0$, then by definition of R_3 and R one has $\langle j \rangle \geq j_{\min}$, $|\ell_2| < c\langle j \rangle^\delta$ and $\pi(\ell_2) \not\parallel v(j)$. But then $|\ell_2| < c\langle j \rangle^\delta$ implies that $|\pi(\ell_2)| < \langle j \rangle^\delta$. Fixing $w \in \mathcal{V}$ to be the direction parallel to $\pi(\ell_2)$ one has $|w| < \langle j \rangle^\delta$, hence

$$|\pi(\ell_2) \cdot j| \geq |w \cdot j| \geq |b(j)| := \min_{v \in \mathcal{V}: |v| < \langle j \rangle^\delta} |j \cdot v|$$

Then, if $|b(j)| > \langle j \rangle^\mu$ our claim follows. Otherwise, we prove the claim by contradiction. Recall that $v(j)$ attains the minimum above and, since $\pi(\ell_2) \not\parallel v(j)$, $v(j) \not\parallel w$. If $|w \cdot j| \leq \langle j \rangle^\mu$ then, by Lemma 2.11

$$\langle j \rangle < \sqrt{2}\langle j \rangle^{1-\delta}$$

which contradicts $\langle j \rangle \geq 2^{\frac{1}{2\delta}}$. This proves that (B.19) holds.

As a consequence, taking $f^\epsilon(\ell_1, \ell_2)$ as in (B.16), one has that, if $(R_3)_j^j(\ell) \neq 0$, then $\forall \epsilon \in \{0, 1\}$

$$\begin{aligned} \frac{\langle j \rangle^\mu}{|\omega \cdot \ell_2 + 2j \cdot \pi(\ell_2) - \epsilon 2\pi(\ell_1) \cdot \pi(\ell_2) - |\pi(\ell_2)|^2|} &= \frac{\langle j \rangle^\mu}{|2j \cdot \pi(\ell_2) + f^\epsilon(\ell_1, \ell_2)|} \\ &\leq \frac{\langle j \rangle^\mu}{2\langle j \rangle^\mu - |\omega||\ell_2| - c^{-2}|\ell_2|^2 - 2c^{-2}|\ell_1||\ell_2|} \leq \frac{\langle j \rangle^\mu}{2\langle j \rangle^\mu - (|\omega| + 2c^{-2})c^2\langle j \rangle^{2\delta}} \leq 1, \end{aligned}$$

since $\langle j \rangle \geq j_{\min}$. This proves that

$$\begin{aligned} \left| ((R_3)_{\frac{a'}{2}; -(2,0)}^h)_j^h \right| &:= \sum_{\ell: j-h=\pi(\ell)} e^{\frac{a'}{2}|\ell|} |(R_3)_j^j(\ell)| \langle j \rangle^{2\mu} \\ &\lesssim_c \sum_{\ell: j-h=\pi(\ell)} e^{\frac{a'}{2}|\ell|} \sum_{\substack{\ell_1+\ell_2=\ell \\ \pi(\ell_2) \not\parallel v(j)}} |V(\ell_1)V(\ell_2)| |\pi(\ell_1)| |\pi(\ell_2)| \lesssim_c a^{-2} e^{-(a-\frac{a'}{2})|j-h|} \|V\|_{\mathbf{a}, \mathbf{p}}^2, \end{aligned}$$

and thus estimate (4.10) follows. \square

C Covariant properties

Assume that H is an Hamiltonian function which is invariant by translation and gauge:

$$H \circ \tau_\zeta = H, \quad H \circ e^{it} = H, \quad \forall \zeta \in \mathbb{R}, \quad \forall t \in \mathbb{R}. \quad (\text{C.1})$$

Its Hamiltonian vector field X_H thus fulfill

$$\tau_\zeta X_H(u) = X_H(\tau_\zeta u), \quad e^{it} X_H(u) = e^{it} X_H(u), \quad \forall \zeta \in \mathbb{R}, \quad \forall t \in \mathbb{R}. \quad (\text{C.2})$$

Consider now a quasi-periodic traveling wave $q(\varphi, x)$ fulfilling (1.8), (1.9). Denote by

$$A(\varphi)u := dX_H(q(\varphi, \cdot))u$$

and

$$\mathcal{N}(\varphi, u) := X_H(q(\varphi, \cdot) + u) - X_H(q(\varphi, \cdot)) - dX_H(q(\varphi, \cdot))u$$

Following [2, 3, Section 3.4], now we prove the following lemma.

Lemma C.1. *For any $(\varphi, \zeta, t) \in \mathbb{T}^d \times \mathbb{R}^2 \times \mathbb{R}$ one has*

$$A(\varphi + \mathbf{K}\zeta) \circ \tau_\zeta = \tau_\zeta \circ A(\varphi), \quad A(\varphi + t\vec{1}) \circ e^{it} = e^{it} \circ A(\varphi), \quad (\text{C.3})$$

$$\mathcal{N}(\varphi + \mathbf{K}\zeta, \tau_\zeta u) = \tau_\zeta \circ \mathcal{N}(\varphi, u), \quad \mathcal{N}(\varphi + t\vec{1}, e^{it}u) = e^{it} \circ \mathcal{N}(\varphi, u). \quad (\text{C.4})$$

Proof. We prove just the first identities in both line, the second one being similar. We start with the first identity of (C.3). Differentiating the first of (C.2) at $q(\varphi, x)$ in direction w we get

$$\tau_\zeta [dX_H(q(\varphi, \cdot))w] = dX_H(\tau_\zeta q(\varphi, \cdot))\tau_\zeta w . \quad (\text{C.5})$$

Then the covariance property (1.9) implies immediately

$$A(\varphi + \mathbf{K}\zeta) \circ \tau_\zeta = \tau_\zeta \circ A(\varphi) \quad (\text{C.6})$$

To prove (C.4) just use again that, by (C.2), (C.5) and the covariance properties (1.9) of the quasi-periodic traveling wave,

$$\begin{aligned} \tau_\zeta \circ \mathcal{N}(\varphi, u) &= X_H(\tau_\zeta q(\varphi, \cdot) + \tau_\zeta u) - X_H(\tau_\zeta q(\varphi, \cdot)) - dX_H(\tau_\zeta q(\varphi, \cdot))\tau_\zeta w \\ &= X_H(q(\varphi + \mathbf{K}\zeta, \cdot) + \tau_\zeta u) - X_H(q(\varphi + \mathbf{K}\zeta, \cdot)) - dX_H(q(\varphi + \mathbf{K}\zeta, \cdot))\tau_\zeta w \\ &= \mathcal{N}(\varphi + \mathbf{K}\zeta, \tau_\zeta u) . \end{aligned}$$

□

References

- [1] M. Berti and L. Biasco. Branching of Cantor manifolds of elliptic tori and applications to PDEs. *Comm. Math. Phys.*, 305(3):741–796, 2011.
- [2] M. Berti, L. Franzoi and A. Maspero. *Traveling quasi-periodic water waves with constant vorticity*, Archive for Rational Mechanics, 240: 99–202, 2021.
- [3] M. Berti, L. Franzoi and A. Maspero. *Pure gravity traveling quasi-periodic water waves with constant vorticity*, arXiv:2101.12006, 2021, to appear on Communications on Pure and Applied Mathematics.
- [4] J. Bernier, E. Faou, and B. Grébert. Rational normal forms and stability of small solutions to nonlinear Schrödinger equations. *Annals of PDE*, 6(14), 2020.
- [5] D. Bambusi and B. Grébert. Birkhoff normal form for partial differential equations with tame modulus. *Duke Math. J.*, 135(3):507–567, 2006.
- [6] J. Bernier and B. Grébert. Birkhoff normal forms for hamiltonian pdes in their energy space. *Journal de l’Ecole Polytechnique - Mathématiques*, 9:681–745, 2022.
- [7] D. Bambusi, B. Langella, and R. Montalto. On the spectrum of the Schrödinger operator on \mathbb{T}^d : a normal form approach. *Comm. Partial Differential Equations*, 45(4):303–320, 2020.
- [8] D. Bambusi, B. Langella, and R. Montalto. Spectral asymptotics of all the eigenvalues of Schrödinger operators on flat tori. *Nonlinear Anal.*, 216:Paper No. 112679, 37, 2022.
- [9] L. Biasco, J. E. Massetti, and M. Procesi. An Abstract Birkhoff Normal Form Theorem and Exponential Type Stability of the 1d NLS. *Comm. Math. Phys.*, 375(3):2089–2153, 2020.
- [10] L. Biasco, J.E. Massetti, and M. Procesi. Almost-periodic invariant tori for the NLS on the circle. *Ann. Inst. H. Poincaré Anal. Non Linéaire*, 38(3):711–758, 2021.
- [11] J. Bourgain. Fourier transform restriction phenomena for certain lattice subsets and application to nonlinear evolution equations. i. Schrödinger equations. *Geom. Funct. Anal.*, 3(2):107–156, 1993.

- [12] J. Bourgain. On the growth in time of higher Sobolev norms of smooth solutions of Hamiltonian PDE. *Internat. Math. Res. Notices*, 6:277–304, 1996.
- [13] J. Bourgain. On invariant tori of full dimension for 1D periodic NLS. *J. Funct. Anal.*, 229(1):62–94, 2005.
- [14] J. Colliander, M. Keel, G. Staffilani, H. Takaoka, and T. Tao. Transfer of energy to high frequencies in the cubic defocusing nonlinear Schrödinger equation. *Invent. Math.*, 181(1):39–113, 2010.
- [15] H. L. Eliasson and S. B. Kuksin. On reducibility of Schrödinger equations with quasiperiodic in time potentials. *Comm. Math. Phys.*, 286(1):125–135, 2009.
- [16] L. H. Eliasson and S. B. Kuksin. KAM for the nonlinear Schrödinger equation. *Ann. of Math. (2)*, 172(1):371–435, 2010.
- [17] E. Faou and B. Grébert. A Nekhoroshev-type theorem for the nonlinear Schrödinger equation on the torus. *Anal. PDE*, 6(6):1243–1262, 2013.
- [18] E. Faou, L. Gauckler, and C. Lubich. Sobolev stability of plane wave solutions to the cubic nonlinear Schrödinger equation on a torus. *Comm. Partial Differential Equations*, 38(7):1123–1140, 2013.
- [19] J. Feldman, H. Knörrer, and E. Trubowitz. The perturbatively stable spectrum of a periodic Schrödinger operator. *Invent. Math.*, 100(2):259–300, 1990.
- [20] J. Feldman, H. Knörrer, and E. Trubowitz. Perturbatively unstable eigenvalues of a periodic Schrödinger operator. *Comment. Math. Helv.*, 66(4):557–579, 1991.
- [21] M. Guardia, Z. Hani, E. Haus, A. Maspero, and M. Procesi. Strong nonlinear instability and growth of Sobolev norms near quasiperiodic finite gap tori for the 2d cubic NLS equation. *J. Eur. Math. Soc., published online first*, 2022.
- [22] M. Guardia, E. Haus, and M. Procesi. Growth of Sobolev norms for the analytic NLS on \mathbb{T}^2 . *Adv. Math.*, 301:615–692, 2016.
- [23] A. Giorgilli. Exponential stability of Hamiltonian systems. In *Dynamical systems. Part I*, Pubbl. Cent. Ric. Mat. Ennio Giorgi, pages 87–198. Scuola Norm. Sup., Pisa, 2003.
- [24] M. Guardia and V. Kaloshin. Growth of Sobolev norms in the cubic defocusing nonlinear Schrödinger equation. *J. Eur. Math. Soc. (JEMS)*, 17(1):71–149, 2015.
- [25] G. Gentile and M. Procesi. Periodic solutions for a class of nonlinear partial differential equations in higher dimension. *Comm. Math. Phys.*, 289(3):863–906, 2009.
- [26] J. Geng, X. Xu, and J. You. An infinite dimensional KAM theorem and its application to the two dimensional cubic Schrödinger equation. *Adv. Math.*, 226(6):5361–5402, 2011.
- [27] Z. Hani. Long-time instability and unbounded Sobolev orbits for some periodic nonlinear Schrödinger equations. *Arch. Ration. Mech. Anal.*, 211(3):929–964, 2014.
- [28] E. Haus and M. Procesi. KAM for beating solutions of the quintic NLS. *Comm. Math. Phys.*, 354(3):1101–1132, 2017.
- [29] T. Kappeler and R. Montalto. On the Stability of Periodic Multi-Solitons of the KdV Equation. *Comm. Math. Phys.*, 385: 1871–1956, 2021.

- [30] Yu. E. Karpeshina. Perturbation series for the Schrödinger operator with a periodic potential near planes of diffraction. *Comm. Anal. Geom.*, 4(3):339–413, 1996.
- [31] Yu. E. Karpeshina. *Perturbation theory for the Schrödinger operator with a periodic potential*, volume 1663 of *Lecture Notes in Mathematics*. Springer-Verlag, Berlin, 1997.
- [32] S.B. Kuksin. Growth and oscillations of solutions of nonlinear Schrödinger equation. *Comm. Math. Phys.*, 178(2):265–280, 1996.
- [33] S.B. Kuksin. On turbulence in nonlinear Schrödinger equations. *Geom. Funct. Anal.*, 7(4):783–822, 1997.
- [34] S.B. Kuksin. Oscillations in space-periodic nonlinear Schrödinger equations. *Geom. Funct. Anal.*, 7(2):338–363, 1997.
- [35] A. Maspero and M. Procesi. Long time stability of small finite gap solutions of the cubic nonlinear Schrödinger equation on \mathbb{T}^2 . *J. Differential Equations*, 265(7):3212–3309, 2018.
- [36] N. N. Nekhorošev. An exponential estimate of the time of stability of nearly integrable Hamiltonian systems. *Uspehi Mat. Nauk*, 32(6(198)):5–66, 287, 1977.
- [37] M. Procesi and C. Procesi. A normal form for the Schrödinger equation with analytic non-linearities. *Comm. Math. Phys.*, 312(2):501–557, 2012.
- [38] C. Procesi and M. Procesi. A KAM algorithm for the resonant non-linear Schrödinger equation. *Adv. Math.*, 272:399–470, 2015.
- [39] L. Parnowski and A. V. Sobolev. Bethe-Sommerfeld conjecture for periodic operators with strong perturbations. *Invent. Math.*, 181(3):467–540, 2010.
- [40] L. Parnowski and R. Shterenberg. Complete asymptotic expansion of the integrated density of states of multidimensional almost-periodic Schrödinger operators. *Ann. of Math. (2)*, 176(2):1039–1096, 2012.
- [41] M. Procesi and X. Xu. Quasi-Töplitz Functions in KAM Theorem. *SIAM J. of Math. Anal.*, 45(4):2148 – 2181, 2013.
- [42] O. Veliev. *Multidimensional periodic Schrödinger operator*, volume 263 of *Springer Tracts in Modern Physics*. Springer, Cham, 2015. Perturbation theory and applications.
- [43] W.-M. Wang. Energy supercritical nonlinear Schrödinger equations: Quasiperiodic solutions. *Duke Math. J.*, 165(6): 1129–1192, 2016.