

Almost-periodic Response Solutions for a forced quasi-linear Airy equation

Livia Corsi[†], Riccardo Montalto^{*}, Michela Procesi[†]

^{*} Università degli Studi di Milano; riccardo.montalto@unimi.it

[†] Università di Roma Tre; lcorsi@mat.uniroma3.it, procesi@mat.uniroma3.it

Abstract

We prove the existence of almost-periodic solutions for quasi-linear perturbations of the Airy equation. This is the first result about the existence of this type of solutions for a quasi-linear PDE. The solutions turn out to be analytic in time and space. To prove our result we use a Craig-Wayne approach combined with a KAM reducibility scheme and pseudo-differential calculus on \mathbb{T}^∞ .

Keywords: Almost-periodic solutions for PDEs; Nash-Moser-KAM theory; small divisor problems; KdV

MSC classification: 37K55; 58C15; 35Q53; 35B15

Contents

1	Introduction	2
2	Functional setting	5
3	The iterative scheme	7
3.1	The zero-th step	10
3.2	The $n + 1$ -th step	11
4	Proof of Proposition 3.6	17
4.1	Elimination of the x -dependence from the highest order term	18
4.2	Elimination of the φ -dependence from the highest order term	19
4.3	Time dependent traslation of the space variable	20
4.4	Conclusion of the proof	21
5	Proof of Proposition 3.8	21
5.1	Reduction of the first order term	22
5.2	Reducibility	24

5.3	Variations	29
5.4	Conclusion of the proof	31
A Technical Lemmata		32

1 Introduction

In this paper we study response solutions for almost-periodically forced quasilinear PDEs close to an elliptic fixed point.

The problem of response solutions for PDEs has been widely studied in many contexts, starting from the papers [24, 25], where the Author considers a periodically forced PDE with dissipation. In the presence of dissipation, of course there is no small divisors problem. However as soon as the dissipation is removed, small divisors appear even in the easiest possible case of a periodic forcing when the spacial variable is one dimensional.

The first results of this type in absence of dissipation were obtained by means of a KAM approach [16, 17, 18, 28, 22, 19]. However, a more functional approach, via a combination of a Ljapunov-Schmidt reduction and a Newton scheme, in the spirit of [24, 25], was proposed by Craig-Wayne [14], and then generalized in many ways by Bourgain; see for instance [5, 6, 7] to mention a few. All the results mentioned above concern semi-linear PDEs and the forcing is quasi-periodic.

In more recent times, the Craig-Wayne-Bourgain approach has been fruitfully used and generalized in order to cover quasi-linear and fully nonlinear PDEs, again in the quasi-periodic case; see for instance [1, 15, 12, 2].

Regarding the almost-periodic case, most of the classical results are obtained via a KAM-like approach; see for instance [10, 23, 9]. A notable exception is [8], where the Craig-Wayne-Bourgain method is used. More recently there have been results such as [26, 27, 20], which use a KAM approach. We mention also [29, 11, 3, 4] which however are tailored for an autonomous PDE.

All the aforementioned results, concern semi-linear PDEs, with no derivative in the nonlinearity. Moreover they require a very strong analyticity condition on the forcing term. Indeed the difficulty of proving the existence of almost-periodic response solution is strongly related to the regularity of the forcing, since one can see an almost periodic function as the limit of quasi-periodic ones with an increasing number of frequencies. If such limit is reached sufficiently fast, the most direct strategy would be to iteratively find approximate quasi-periodic response solutions and then take the limit. This is the overall strategy of [23] and [26, 27, 20]. However this procedure works if one considers a sufficiently regular forcing term and a bounded nonlinearity, but becomes very delicate in the case of unbounded nonlinearities.

In the present paper we study the existence of almost-periodic response solutions, for a quasi-linear PDE on \mathbb{T} . To the best of our knowledge this is the first result of this type.

Specifically we consider a quasi-linear Airy equation

$$\partial_t u + \partial_{xxx} u + Q(u, u_x, u_{xx}, u_{xxx}) + \mathfrak{f}(t, x) = 0, \quad x \in \mathbb{T} := (\mathbb{R}/(2\pi\mathbb{Z})) \quad (1.1)$$

where Q is a Hamiltonian, quadratic nonlinearity and \mathfrak{f} is an analytic forcing term with zero average w.r.t. x . We assume \mathfrak{f} to be “almost-periodic” with frequency $\omega \in \ell^\infty$, in the sense of Definition 1.1.

We mention that in the context of reducibility of linear PDEs a problem of this kind has been solved in [21]. Our aim is to provide a link between the linear techniques of [21] and the nonlinear Craig-Wayne-Bourgain method. Note that such a link is nontrivial, and requires a delicate handling; see below.

The overall setting we use is the one of [1]. However their strategy is tailored for Sobolev regularity; the quasi-periodic analytic case has been covered in [13]. Unfortunately the ideas of [13] cannot be directly applied in the almost-periodic case. Roughly, it is well known that the regularity and the small-divisor problem conflict. Thus, in the almost-periodic case one expect this issue to be even more dramatic. Specifically, we were not able to define a “Sobolev” norm for almost-periodic functions, satisfying the interpolation estimates needed in the Nash-Moser scheme; this is why we cannot use the theorem of [13].

Let us now present our main result in a more detailed way.

First of all we note that (1.1) is an Hamiltonian PDE whose Hamiltonian is given by

$$H(u) := \frac{1}{2} \int_{\mathbb{T}} u_x^2 dx - \frac{1}{6} \int_{\mathbb{T}} G(u, u_x) dx - \int_{\mathbb{T}} F(t, x) u dx, \quad \mathfrak{f}(t, x) = \partial_x F(t, x) \quad (1.2)$$

where $G(u, u_x)$ is a cubic Hamiltonian density of the form

$$G(u, u_x) := c_3 u_x^3 + c_2 u u_x^2 + c_1 u^2 u_x + c_0 u^3, \quad c_0, \dots, c_3 \in \mathbb{R} \quad (1.3)$$

and the symplectic structure is given by $J = \partial_x$. The Hamiltonian nonlinearity $Q(u, \dots, u_{xxx})$ is therefore given by

$$Q(u, u_x, u_{xx}, u_{xxx}) = \partial_{xx}(\partial_{u_x} G(u, u_x)) - \partial_x(\partial_u G(u, u_x)) \quad (1.4)$$

and the Hamilton equations are

$$\partial_t u = \partial_x \nabla_u H(u).$$

We look for an almost-periodic solution to (1.1) with frequency ω in the sense below.

For $\eta > 0$, define the set of infinite integer vectors with *finite support* as

$$\mathbb{Z}_*^\infty := \left\{ \ell \in \mathbb{Z}^{\mathbb{N}} : |\ell|_\eta := \sum_{i \in \mathbb{N}} i^\eta |\ell_i| < \infty \right\}. \quad (1.5)$$

Note that $\ell_i \neq 0$ only for finitely many indices $i \in \mathbb{N}$. In particular \mathbb{Z}_*^∞ does not depend on η .

Definition 1.1. Given $\omega \in [1, 2]^{\mathbb{N}}$ with rationally independent components¹ and a Banach space $(X, |\cdot|_X)$, we say that $F(t) : \mathbb{R} \rightarrow X$ is almost-periodic in time with frequency ω and analytic in the strip $\sigma > 0$ if we may write it in totally convergent Fourier series

$$F(t) = \sum_{\ell \in \mathbb{Z}_*^\infty} F(\ell) e^{i\ell \cdot \omega t} \quad \text{such that} \quad F(\ell) \in X, \forall \ell \in \mathbb{Z}_*^\infty$$

$$\text{and} \quad |F|_\sigma := \sum_{\ell \in \mathbb{Z}_*^\infty} |F(\ell)|_X e^{\sigma|\ell|_\eta} < \infty.$$

We shall be particularly interested in almost-periodic functions where $X = \mathcal{H}_0(\mathbb{T}_\sigma)$

$$\mathcal{H}_0(\mathbb{T}_\sigma) := \left\{ u = \sum_{j \in \mathbb{Z} \setminus \{0\}} u_j e^{ijx}, u_j = \bar{u}_{-j} \in \mathbb{C} : |u|_{\mathcal{H}(\mathbb{T}_\sigma)} := \sum_{j \in \mathbb{Z} \setminus \{0\}} |u_j| e^{\sigma|j|} < \infty \right\}$$

is the space of analytic, real on real functions $\mathbb{T}_s \rightarrow \mathbb{C}$ with zero-average, where $\mathbb{T}_s := \{\varphi \in \mathbb{C} : \text{Re}(\varphi) \in \mathbb{T}, |\text{Im}(\varphi)| \leq s\}$ is the thickened torus. We recall that a function $u : \mathbb{T}_s \rightarrow \mathbb{C}$ is real on real if for any $x \in \mathbb{T}$, $u(x) \in \mathbb{R}$.

Of course we need some kind of Diophantine condition on ω . We give the following, taken from [9, 21].

Definition 1.2. Given $\gamma \in (0, 1)$, we denote by \mathcal{D}_γ the set of Diophantine frequencies

$$\mathcal{D}_\gamma := \left\{ \omega \in [1, 2]^{\mathbb{N}} : |\omega \cdot \ell| > \gamma \prod_{i \in \mathbb{N}} \frac{1}{(1 + |\ell_i|^2 i^2)}, \quad \forall \ell \in \mathbb{Z}_*^\infty \setminus \{0\} \right\}. \quad (1.6)$$

We are now ready to state our main result.

Theorem 1.3 (Main Theorem). Fix $\bar{\gamma}$. Assume that \mathfrak{f} in (1.1) is almost-periodic in time and analytic in a strip S (both in time and space). Fix $\bar{s} < S$. If \mathfrak{f} has an appropriately small norm depending on $S - \bar{s}$, namely

$$|\mathfrak{f}|_S := \sum_{\ell \in \mathbb{Z}_*^\infty} |\mathfrak{f}(\ell)|_{\mathcal{H}_0(\mathbb{T}_S)} e^{S|\ell|_\eta} \leq \epsilon(S - \bar{s}) \ll 1, \quad \epsilon(0) = 0, \quad (1.7)$$

then there is a Cantor-like set $\mathcal{O}^{(\infty)} \subseteq \mathcal{D}_{\bar{\gamma}}$ with positive Lebesgue measure, and for all $\omega \in \mathcal{O}^{(\infty)}$ a solution to (1.1) which is almost-periodic in time with frequency ω and analytic in a strip \bar{s} (both in time and space).

Remark 1.4. Of course the same result holds verbatim if we replace the quadratic polynomial Q by a polynomial of arbitrary degree. We could also assume that the coefficients c_j appearing in (1.4) depend on x and ωt . In that case Theorem 1.3 holds provided we further require a condition of the type $\sup_j |\partial_x^2 c_j|_S \leq C$. Actually one could also take Q to be an analytic function with a zero of order two. However this leads to a number of long and non particularly enlightening calculations.

¹We say that ω has rationally independent components if for any $N > 0$ and any $k \in \mathbb{Z}^N$ one has $\sum_{i=1}^N \omega_i k_i \neq 0$.

To prove Theorem 1.3 we proceed as follows. First of all we regard (1.1) as a functional Implicit Function Problem on some appropriate space of functions defined on an infinite dimensional torus; see Definition 2.1 below. Then in Section 3 we prove an iterative ‘‘Nash-Moser-KAM’’ scheme to produce the solution of such Implicit Function Problem. It is well known that an iterative rapidly converging scheme heavily relies on a careful control on the invertibility of the linearized operator at any approximate solution. Of course, in the case of a quasi-linear PDE this amounts to study an unbounded non-constant coefficients operator. To deal with this problem, at each step we introduce a change of variables T_n which diagonalizes the highest order terms of the linearized operator. An interesting feature is that T_n preserves the PDE structure. As in [13] and differently from the classical papers, at each step we apply the change of variables T_n to the whole nonlinear operator. This is not a merely technical issue. Indeed, the norms we use are strongly coordinate-depending, and the change of variable T_n that we need to apply are not close-to-identity, in the sense that $T_n - \text{Id}$ is not a bounded operator small in size.

In Section 4 we show how to construct the change of variables T_n satisfying the properties above. Then in order to prove the invertibility of the linearized operator after the change of variables T_n is applied, one needs to perform a reducibility scheme: this is done in Section 5. For a more detailed description of the technical aspects see Remark 3.2.

ACKNOWLEDGEMENTS. Riccardo Montalto is supported by INDAM-GNFM.

2 Functional setting

As it is habitual in the theory of quasi-periodic functions we shall study almost periodic functions in the context of analytic functions on an infinite dimensional torus. To this purpose, for $\eta, s > 0$, we define the *thickened* infinite dimensional torus \mathbb{T}_s^∞ as

$$\varphi = (\varphi_i)_{i \in \mathbb{N}}, \quad \varphi_i \in \mathbb{C} : \text{Re}(\varphi_i) \in \mathbb{T}, \quad |\text{Im}(\varphi_i)| \leq s \langle i \rangle^\eta.$$

Given a Banach space $(X, |\cdot|_X)$ we consider the space \mathcal{F} of pointwise absolutely convergent formal Fourier series $\mathbb{T}_s^\infty \rightarrow X$

$$u(\varphi) = \sum_{\ell \in \mathbb{Z}_*^\infty} u(\ell) e^{i\ell \cdot \varphi}, \quad u(\ell) \in X \tag{2.1}$$

and define the analytic functions as follows.

Definition 2.1. *Given a Banach space $(X, |\cdot|_X)$ and $s > 0$, we define the space of analytic functions $\mathbb{T}_s^\infty \rightarrow X$ as the subspace*

$$\mathcal{H}(\mathbb{T}_s^\infty, X) := \left\{ u(\varphi) = \sum_{\ell \in \mathbb{Z}_*^\infty} u(\ell) e^{i\ell \cdot \varphi} \in \mathcal{F} : |u|_s := \sum_{\ell \in \mathbb{Z}_*^\infty} e^{s|\ell|^\eta} |u(\ell)|_X < \infty \right\}.$$

We denote by \mathcal{H}_s the subspace of $\mathcal{H}(\mathbb{T}_s^\infty, \mathcal{H}_0(\mathbb{T}_s))$ of the functions which are real on real. Moreover, we denote by $\mathcal{H}(\mathbb{T}_s^\infty \times \mathbb{T}_s)$, the space of analytic functions $\mathbb{T}_s^\infty \times \mathbb{T}_s \rightarrow \mathbb{C}$ which

are real on real. The space \mathcal{H}_s can be identified with the subspace of zero-average functions of $\mathcal{H}(\mathbb{T}_s^\infty \times \mathbb{T}_s)$. Indeed if $u \in \mathcal{H}_s$, then

$$u = \sum_{\ell \in \mathbb{Z}_*^\infty} u(\ell, x) e^{i\ell \cdot \varphi} = \sum_{(\ell, j) \in \mathbb{Z}_*^\infty \times \mathbb{Z} \setminus \{0\}} u_j(\ell) e^{i\ell \cdot \varphi + i j x},$$

with $u_j(\ell) = \overline{u_{-j}(-\ell)}$

For any $u \in \mathcal{H}(\mathbb{T}_s^\infty \times \mathbb{T}_s)$ let us denote

$$(\pi_0 u)(\varphi, x) := \langle u(\varphi, \cdot) \rangle_x := \frac{1}{2\pi} \int_{\mathbb{T}} u(\varphi, x) dx, \quad \pi_0^\perp := \mathbb{1} - \pi_0. \quad (2.2)$$

Throughout the algorithm we shall need to control the Lipschitz variation w.r.t. ω of functions in some $\mathcal{H}(\mathbb{T}_s^\infty, X)$, which are defined for ω in some Cantor set. Thus, for $\mathcal{O} \subset \mathcal{O}^{(0)}$ we introduce the following norm.

Parameter dependence. Let Y be a Banach space and $\gamma \in (0, 1)$. If $f : \Omega \rightarrow Y$, $\Omega \subseteq [1, 2]^{\mathbb{N}}$ is a Lipschitz function we define

$$|f|_Y^{\text{sup}} := \sup_{\omega \in \Omega} |f(\omega)|_Y, \quad |f|_Y^{\text{lip}} := \sup_{\substack{\omega_1, \omega_2 \in \Omega \\ \omega_1 \neq \omega_2}} \frac{|f(\omega_1) - f(\omega_2)|_Y}{|\omega_1 - \omega_2|_\infty}, \quad (2.3)$$

$$|f|_Y^\Omega := |f|_Y^{\text{sup}} + \gamma |f|_Y^{\text{lip}}.$$

If $Y = \mathcal{H}_s$ we simply write $|\cdot|_\sigma^{\text{sup}}$, $|\cdot|_\sigma^{\text{lip}}$, $|\cdot|_\sigma^\Omega$. If Y is a finite dimensional space, we write $|\cdot|_{\text{sup}}$, $|\cdot|_{\text{lip}}$, $|\cdot|^\Omega$.

Linear operators. For any $\sigma > 0$, $m \in \mathbb{R}$ we define the class of linear operators of order m (densely defined on $L^2(\mathbb{T})$) $\mathcal{B}^{\sigma, m}$ as

$$\mathcal{B}^{\sigma, m} := \left\{ \mathcal{R} : L^2(\mathbb{T}) \rightarrow L^2(\mathbb{T}) : \|\mathcal{R}\|_{\mathcal{B}^{\sigma, m}} < \infty \right\} \quad \text{where}$$

$$\|\mathcal{R}\|_{\mathcal{B}^{\sigma, m}} := \sup_{j' \in \mathbb{Z} \setminus \{0\}} \sum_{j \in \mathbb{Z} \setminus \{0\}} e^{\sigma|j-j'|} |R_j^{j'}| \langle j' \rangle^{-m}. \quad (2.4)$$

and for $\mathcal{T} \in \mathcal{H}(\mathbb{T}_\sigma^\infty, \mathcal{B}^{\sigma, m})$ we set

$$\|\mathcal{T}\|_{\sigma, m} := \sum_{\ell \in \mathbb{Z}_*^\infty} e^{\sigma|\ell|_\eta} \|\mathcal{T}(\ell)\|_{\mathcal{B}^{\sigma, m}}. \quad (2.5)$$

In particular we shall denote by $\|\cdot\|_{\sigma, m}^\Omega$ the corresponding Lipschitz norm. Moreover if $m = 0$ we shall drop it, and write simply $\|\cdot\|_\sigma$ or $\|\cdot\|_\sigma^\Omega$.

3 The iterative scheme

Let us rewrite (1.1) as

$$F_0(u) = 0 \tag{3.1}$$

where

$$F_0(u) := (\omega \cdot \partial_\varphi + \partial_{xxx})u + Q(u, u_x, u_{xx}, u_{xxx}) + f(\varphi, x) \tag{3.2}$$

where we $f(t, x) = f(\omega t, x)$ and, as customary the unknown u is a function of $(\varphi, x) \in \mathbb{T}^\infty \times \mathbb{T}$.

We introduce the (Taylor) notation

$$\begin{aligned} L_0 &:= (\omega \cdot \partial_\varphi + \partial_{xxx}) = F'_0(0), & f_0 &= F_0(0) = f(\varphi, x), \\ Q_0(u) &= Q(u, u_x, u_{xx}, u_{xxx}) \stackrel{(1.4)}{=} \partial_{xx} \left(3c_3 u_x^2 + 2c_2 u u_x + c_1 u^2 \right) \\ &\quad - \partial_x (c_2 u_x^2 + 2c_1 u u_x + 3c_0 u^2) \end{aligned} \tag{3.3}$$

so that (3.1) reads

$$f_0 + L_0 u + Q_0(u) = 0.$$

Note that Q_0 is of the form

$$Q_0(u) = \sum_{\substack{0 \leq i \leq 2, 0 \leq j \leq 3 \\ 0 \leq i+j \leq 4}} q_{i,j}^{(0)} (\partial_x^i u) (\partial_x^j u) \tag{3.4}$$

with the coefficients $q_{i,j}^{(0)}$ satisfying

$$\sum_{\substack{0 \leq i \leq 2, 0 \leq j \leq 3 \\ 0 \leq i+j \leq 4}} |q_{i,j}^{(0)}| \leq C, \tag{3.5}$$

where the constant C depends clearly on $|c_0|, \dots, |c_3|$. In particular, this implies that for all $u \in \mathcal{H}_s$ one has the following.

Q1. $|Q_0(u)|_{s-\sigma} \lesssim \sigma^{-4} |u|_s^2$

Q2. $|Q'_0(u)[h]|_{s-\sigma} \lesssim \sigma^{-4} |u|_s |h|_s$

We now fix the constants

$$\begin{aligned} \mu &> \max\left\{1, \frac{1}{\eta}\right\}, \\ \gamma_0 &< \frac{1}{2}\bar{\gamma}, \quad \gamma_n := (1 - 2^{-n})\gamma_{n-1}, \quad n \geq 1 \\ \sigma_{-1} &:= \frac{1}{8} \min\{(S - \bar{s}), 1\}, \quad \sigma_{n-1} = \frac{6\sigma_{-1}}{\pi^2 n^2}, \quad n \geq 1, \\ s_0 &= S - \sigma_{-1}, \quad s_n = s_{n-1} - 6\sigma_{n-1}, \quad n \geq 1, \\ \varepsilon_n &:= \varepsilon_0 e^{-\chi^n}, \quad \chi = \frac{3}{2}, \end{aligned} \tag{3.6}$$

where ε_0 is such that

$$e^{c_0\sigma_0^{-1}}|f|_S = e^{c_0\sigma_0^{-1}}|f_0|_S \ll \varepsilon_0. \quad (3.7)$$

Introduce

$$\mathbf{d}(\ell) := \prod_{i \in \mathbb{N}} (1 + |\ell_i|^5 \langle i \rangle^5), \quad \forall \ell \in \mathbb{Z}_*^\infty. \quad (3.8)$$

We also set $\mathcal{O}^{(-1)} := \mathcal{D}_{\bar{\gamma}}$ and

$$\mathcal{O}^{(0)} := \left\{ \omega \in \mathcal{D}_{\bar{\gamma}} : |\omega \cdot \ell + j^3| \geq \frac{\gamma_0}{\mathbf{d}(\ell)}, \quad \forall \ell \in \mathbb{Z}_*^\infty, \quad j \in \mathbb{N}, \quad (\ell, j) \neq (0, 0) \right\}. \quad (3.9)$$

Proposition 3.1. *There exists $\tau, \tau_1, \tau_2, \tau_3, \mathcal{C}, \varepsilon_0$ (pure numbers) such that for*

$$\varepsilon_0 \leq \sigma_0^\tau e^{-\mathcal{C}\sigma_0^{-\mu}} \varepsilon_0, \quad (3.10)$$

for all $n \geq 1$ the following hold.

1. *There exist a sequence of Cantor sets $\mathcal{O}^{(n)} \subseteq \mathcal{O}^{(n-1)}$, $n \geq 1$ such that*

$$\mathbb{P}(\mathcal{O}^{(n-1)} \setminus \mathcal{O}^{(n)}) \lesssim \frac{\gamma_0}{n^2}. \quad (3.11)$$

2. *For $n \geq 1$, there exists a sequence of linear, invertible, bounded and symplectic changes of variables defined for $\omega \in \mathcal{O}^{(n-1)}$, of the form*

$$T_n v(\varphi, x) = (1 + \xi_x^{(n)}) v(\varphi + \omega \beta^{(n)}(\varphi), x + \xi^{(n)}(\varphi, x) + p^{(n)}(\varphi)) \quad (3.12)$$

satisfying

$$|\xi^{(n)}|_{s_{n-1}-\sigma_{n-1}}^{\mathcal{O}^{(n-1)}}, |\beta^{(n)}|_{s_{n-1}-\sigma_{n-1}}^{\mathcal{O}^{(n-1)}}, |p^{(n)}|_{s_{n-1}-\sigma_{n-1}}^{\mathcal{O}^{(n-1)}} \lesssim \sigma_{n-1}^{-\tau_1} \varepsilon_{n-1} e^{C\sigma_{n-1}^{-\mu}}, \quad (3.13)$$

for some constant $C > 0$.

3. *For $n \geq 0$, there exists a sequence of functionals $F_n(u) \equiv F_n(\omega, u(\omega))$, defined for $\omega \in \mathcal{O}^{(n-1)}$, of the form*

$$F_n(u) = f_n + L_n u + Q_n(u), \quad (3.14)$$

such that

- (a) *L_n is invertible for $\omega \in \mathcal{O}^{(n)}$ and setting*

$$h_n := -L_n^{-1} f_n, \quad (3.15)$$

there exists $\mathbf{r}_n = \mathbf{r}_n(\varphi) \in \mathcal{H}(\mathbb{T}_{s_{n-1}-3\sigma_{n-1}}^\infty)$ such that

$$\begin{aligned} F_n(u) &= \mathbf{r}_n T_n^{-1} F_{n-1}(h_{n-1} + T_k u), \quad n \geq 1, \\ |\mathbf{r}_n - 1|_{s_{n-1}-3\sigma_{n-1}}^{\mathcal{O}^{(n-1)}} &\leq \sigma_{n-1}^{-\tau_2} e^{C\sigma_{n-1}^{-\mu}} \varepsilon_{n-1} \end{aligned} \quad (3.16)$$

(b) $f_n = f_n(\varphi, x)$ is a given function satisfying

$$|f_n|_{s_{n-1}-2\sigma_{n-1}}^{\mathcal{O}^{(n-1)}} \lesssim \sigma_{n-1}^{-4} \varepsilon_{n-1}^2, \quad n \geq 1 \quad (3.17)$$

(c) L_n is a linear operator of the form

$$L_n = \omega \cdot \partial_\varphi + (1 + A_n) \partial_{xxx} + B_n(\varphi, x) \partial_x + C_n(\varphi, x) \quad (3.18)$$

such that

$$\frac{1}{2\pi} \int_{\mathbb{T}} B_n(\varphi, x) dx = \bar{b}_n \quad (3.19)$$

and for $n \geq 1$

$$\begin{aligned} |A_n - A_{n-1}|^{\mathcal{O}^{(n-1)}} &\leq \sigma_{n-1}^{-\tau_2} e^{C\sigma_{n-1}^{-\mu}} \varepsilon_{n-1}, \\ |B_n - B_{n-1}|_{s_{n-1}-3\sigma_{n-1}}^{\mathcal{O}^{(n-1)}} &\lesssim \sigma_{n-1}^{-\tau_2} e^{C\sigma_{n-1}^{-\mu}} \varepsilon_{n-1} \\ |C_n - C_{n-1}|_{s_{n-1}-3\sigma_{n-1}}^{\mathcal{O}^{(n-1)}} &\lesssim \sigma_{n-1}^{-\tau_2} e^{C\sigma_{n-1}^{-\mu}} \varepsilon_{n-1}. \end{aligned} \quad (3.20)$$

(d) Q_n is of the form

$$Q_n(u) = \sum_{\substack{0 \leq i \leq 2, 0 \leq j \leq 3 \\ 0 \leq i+j \leq 4}} q_{i,j}^{(n)}(\varphi, x) (\partial_x^i u) (\partial_x^j u) \quad (3.21)$$

with the coefficients $q_{i,j}^{(n)}(\varphi, x)$ satisfying (3.5) for $n = 0$, while for $n \geq 1$

$$\begin{aligned} \sum_{\substack{0 \leq i \leq 2, 0 \leq j \leq 3 \\ 0 \leq i+j \leq 4}} |q_{i,j}^{(n)}|_{s_{n-1}-3\sigma_{n-1}}^{\mathcal{O}^{(n-1)}} &\leq C \sum_{l=1}^n 2^{-l}, \\ |q_{i,j}^{(n)} - q_{i,j}^{(n-1)}|_{s_{n-1}-3\sigma_{n-1}}^{\mathcal{O}^{(n-1)}} &\lesssim \sigma_{n-1}^{-\tau_3} e^{C\sigma_{n-1}^{-\mu}} \varepsilon_{n-1}. \end{aligned} \quad (3.22)$$

4. Finally one has

$$|h_n|_{s_n}^{\mathcal{O}^{(n)}} \leq \varepsilon_n \quad (3.23)$$

Moreover, setting

$$\mathcal{O}^{(\infty)} := \bigcap_{n \geq 0} \mathcal{O}^{(n)}, \quad (3.24)$$

and

$$u_n = h_0 + \sum_{j=1}^n T_1 \circ \dots \circ T_j h_j. \quad (3.25)$$

then

$$u_\infty := \lim_{n \rightarrow \infty} u_n$$

is well defined for $\omega \in \mathcal{O}^{(\infty)}$, belongs to $\mathcal{H}_{\bar{s}}$, and solves $F(u_\infty) = 0$. Finally the $\mathcal{O}^{(\infty)}$ has positive measure; precisely

$$\mathbb{P}(\mathcal{O}^{(\infty)}) = 1 - O(\gamma_0). \quad (3.26)$$

From Proposition 3.1 our main result Theorem 1.3 follows immediately by noting that (3.7) and (3.10) follow from (1.7) for an appropriate choice $\varepsilon(S - \bar{s})$.

Remark 3.2. *Let us spend few words on the strategy of the algorithm. At each step we apply an affine change of variables translating the approximate solution to zero; the translation is not particularly relevant and we perform it only to simplify the notation. On the other hand the linear change of variables is crucial.*

In (3.14) we denote by f_n the “constant term”, by L_n is the “linearized” term and by Q_n the “quadratic” part. In this way the approximate solution at the n -th step is $h_n = -L_n^{-1}f_n$.

In a classical KAM algorithm, in order to invert L_n one typically applies a linear change of variables that diagonalizes L_n ; this, together with the translation by h_n is the affine change of variables mentioned above, at least in the classical KAM scheme.

Unfortunately, in the case of unbounded nonlinearities this cannot be done. Indeed in order to diagonalize L_n in the unbounded case, one needs it to be a pseudo-differential operator. On the other hand, after the diagonalization is performed, one loses the pseudo-differential structure for the subsequent step. Thus we chose the operators T_n in (3.12) in such a way that we preserve the PDE structure and at the same time we diagonalize the highest order terms.

In the [1]-like algorithm the Authors do not apply any change of variables, but they use the reducibility of L_n only in order to deduce the estimates. However such a procedure works only in Sobolev class. Indeed in the analytic case, at each iterative step one needs to lose some analyticity, due to the small divisors. Since we are studying almost-periodic solutions, we need the analytic setting to deal with the small divisors. As usual, the problem is that the loss of the analyticity is related to the size of the perturbation; in the present case, at each step L_n is a diagonal term plus a perturbation $O(\varepsilon_0)$ with the same ε_0 for all n .

A more refined approach is to consider L_n as a small variation of L_{n-1} ; however the problem is that such small variation is unbounded. As a consequence, the operators T_n are not “close-to-identity”. However, since F_n is a differential operator, then the effect of applying T_n is simply a slight modification of the coefficients; see (3.20) and (3.22). Hence there is a strong motivation for applying the operators T_n . In principle we could have also diagonalized the terms up to order $-k$ for any $k \geq 0$; however the latter change of variables are close to the identity and they introduce pseudo-differential terms.

3.1 The zero-th step

Item 1., 2. are trivial for $n = 0$ while item 3.(b), (c), (d) amount to the definition of F_0 , see (3.2), (3.3), (3.4). Regarding item 3.(a) the invertibility of L_0 follows from the definition of $\mathcal{O}^{(0)}$. Indeed, consider the equation

$$L_0 h_0 = -f_0 \tag{3.27}$$

with

$$\langle f_0(\varphi, \cdot) \rangle_x = 0$$

we have the following result.

Lemma 3.3 (Homological equation). *Let $s > 0$, $0 < \sigma < 1$, $f_0 \in \mathcal{H}_{s+\sigma}$, $\omega \in \mathcal{O}^{(0)}$ (see (1.6)). Then there exists a unique solution $h_0 \in \mathcal{H}_s$ of (3.27). Moreover one has*

$$|h_0|_s^{\mathcal{O}^{(0)}} \lesssim \gamma^{-1} \exp\left(\frac{\tau}{\sigma^{\frac{1}{\eta}}} \ln\left(\frac{\tau}{\sigma}\right)\right) |f|_{s+\sigma}.$$

for some constant $\tau = \tau(\eta) > 0$.

Remark 3.4. *Note that from Lemma 3.3 above it follows that there is \mathfrak{C}_0 such that a solution h_0 of (3.27) actually satisfies*

$$|h_0|_s^{\mathcal{O}^{(0)}} \lesssim e^{\mathfrak{C}_0 \sigma^{-\mu}} |f|_{s+\sigma}. \quad (3.28)$$

where we recall that by (3.6), $\mu > \max\{1, \frac{1}{\eta}\}$. Of course the constant \mathfrak{C}_0 is correlated with the correction to the exponent $\frac{1}{\eta}$.

From Lemma 3.3 and (3.27) it follows that h_0 is analytic in a strip s_0 (where $S = s_0 + \sigma_{-1}$ is the analyticity of f , to be chosen). Moreover, by Lemma 3.3 the size of h_0 is

$$|h_0|_{s_0}^{\mathcal{O}^{(0)}} \sim e^{\mathfrak{C}_0 \sigma_{-1}^{-\mu}} |f_0|_S \quad (3.29)$$

proving item 4. for $|f_0|_S$ small enough, which is true by (3.7).

3.2 The $n + 1$ -th step

Assume now that we iterated the procedure above up to $n \geq 0$ times. This means that we arrived at a quadratic equation

$$F_n(u) = 0, \quad F_n(u) = f_n + L_n u + Q_n(u). \quad (3.30)$$

Defined on $\mathcal{O}^{(n-1)}$ (recall that $\mathcal{O}^{(-1)} = \mathcal{D}_\gamma$).

By the inductive hypothesis (3.22) we deduce that for all $0 < s - \sigma < s_{n-1} - 3\sigma_{n-1}$ one has

$$|Q_n(u)|_{s-\sigma}^{\mathcal{O}^{(n-1)}} \lesssim \sigma^{-4} (|u|_s^{\mathcal{O}^{(n-1)}})^2 \quad (3.31a)$$

$$|Q'_n(u)[h]|_{s-\sigma}^{\mathcal{O}^{(n-1)}} \lesssim \sigma^{-4} |u|_s^{\mathcal{O}^{(n-1)}} |h|_s^{\mathcal{O}^{(n-1)}} \quad (3.31b)$$

Moreover, again by the inductive hypothesis, we can invert L_n and define h_n by (3.15). Now we set

$$F_{n+1}(v) = \mathbf{r}_{n+1} T_{n+1}^{-1} F_n(h_n + T_{n+1} v) \quad (3.32)$$

where

$$T_{n+1} v(\varphi, x) = (1 + \xi_x^{(n+1)}) v(\varphi + \omega \beta^{(n+1)}(\varphi), x + \xi^{(n+1)}(\varphi, x) + p^{(n+1)}(\varphi)) \quad (3.33)$$

and r_{n+1} are to be chosen in order to ensure that $L_{n+1} := F'_{n+1}(0)$ has the form (3.18) with $n \rightsquigarrow n + 1$.

Of course by Taylor expansion we can identify

$$\begin{aligned}
f_{n+1} &= \mathbf{r}_{n+1} T_{n+1}^{-1} (f_n + L_n(h_n) + Q_n(h_n)) = \mathbf{r}_{n+1} T_{n+1}^{-1} Q_n(h_n), \\
L_{n+1} &= \mathbf{r}_{n+1} T_{n+1}^{-1} (L_n + Q'_n(h_n)) T_{n+1} \\
Q_{n+1}(v) &= \mathbf{r}_{n+1} (T_{n+1}^{-1} (Q_n(h_n + T_{n+1}v) - Q_n(h_n) - Q'_n(h_n) T_{n+1}v)) \\
&= \mathbf{r}_{n+1} T_{n+1}^{-1} Q_n(T_{n+1}v).
\end{aligned} \tag{3.34}$$

Remark 3.5. Note that the last equality in (3.34) follows from the fact that the nonlinearity Q in (1.1) is quadratic. In the general case, the last term is controlled by the second derivative, and thus one has to assume a bound of the type (3.31) for Q'' .

In section 4 we prove the following

Proposition 3.6. Assuming that

$$\varepsilon_n \leq \sigma_n^{\tau_1+1} e^{-C\sigma_n^{-\mu}} \tag{3.35}$$

for some $C > 0$, there exist $\xi^{(n+1)}$, $\beta^{(n+1)}$, $p^{(n+1)}$ and $\mathbf{r}_{n+1} \in \mathcal{H}(\mathbb{T}_{s_n-\sigma_n}^\infty \times \mathbb{T}_{s_n-\sigma_n})$, defined for all $\omega \in \mathcal{O}^{(n)}$ and satisfying

$$|\xi^{(n+1)}|_{s_n-\sigma_n}^{\mathcal{O}^{(n)}}, |\beta^{(n+1)}|_{s_n-\sigma_n}^{\mathcal{O}^{(n)}}, |p^{(n+1)}|_{s_n-\sigma_n}^{\mathcal{O}^{(n)}}, |\mathbf{r}_{n+1} - 1|_{s_n-\sigma_n}^{\mathcal{O}^{(n)}} \lesssim \sigma_n^{-\tau_1} \varepsilon_n e^{C\sigma_n^{-\mu}} \tag{3.36}$$

such that (3.33) is well defined and symplectic as well as its inverse, and moreover

$$\mathbf{r}_{n+1} T_{n+1}^{-1} (L_n + Q'_n(h_n)) T_{n+1} = \omega \cdot \partial_\varphi + (1 + A_{n+1}) \partial_{xxx} + B_{n+1}(\varphi, x) \partial_x + C_{n+1}(\varphi, x) \tag{3.37}$$

and (3.19) and (3.20) hold with $n \rightsquigarrow n+1$.

The assumption (3.35) follows from (3.10), provided that we choose the constants τ, \mathbf{C} and ε_0 appropriately.

We now prove (3.21) and (3.22) for $n \rightsquigarrow n+1$, namely the following result.

Lemma 3.7. One has

$$Q_{n+1}(v) = \mathbf{r}_{n+1} T_{n+1}^{-1} Q_n(T_{n+1}v) = \mathbf{r}_{n+1} \sum_{\substack{0 \leq i \leq 2, 0 \leq j \leq 3 \\ 0 \leq i+j \leq 4}} q_{i,j}^{(n+1)}(\varphi, x) (\partial_x^i v) (\partial_x^j v) \tag{3.38}$$

with the coefficients $q_{i,j}^{(n+1)}(\varphi, x)$ satisfying

$$\begin{aligned}
\sum_{\substack{0 \leq i \leq 2, 0 \leq j \leq 3 \\ 0 \leq i+j \leq 4}} |q_{i,j}^{(n+1)}|_{s_n-3\sigma_n}^{\mathcal{O}^{(n)}} &\leq C \sum_{l=1}^{n+1} 2^{-l}, \\
|q_{i,j}^{(n+1)} - q_{i,j}^{(n)}|_{s_n-3\sigma_n}^{\mathcal{O}^{(n)}} &\lesssim \sigma_n^{-\tau_3} e^{C\sigma_n^{-\mu}} \varepsilon_n.
\end{aligned} \tag{3.39}$$

Proof. By construction

$$Q_{n+1}(u) = \mathbf{r}_{n+1} \sum_{\substack{0 \leq i \leq 2, 0 \leq j \leq 3 \\ 0 \leq i+j \leq 4}} T_{n+1}^{-1}[q_{i,j}^{(n)}(\varphi, x)(\partial_x^i T_{n+1}v)(\partial_x^j T_{n+1}v)]. \quad (3.40)$$

Now we first note that

$$\partial_x(T_{n+1}v) = \xi_{xx}^{(n+1)}v(\theta, y) + (1 + \xi_x)^2v_y(\theta, y)$$

where

$$(\theta, y) = (\varphi + \omega\beta^{(n+1)}(\varphi), x + \xi^{(n+1)}(\varphi, x) + p^{(n+1)}(\varphi)).$$

Hence the terms $\partial_x^i T_{n+1}v$ are of the form

$$\partial_x^i T_{n+1}v = \partial_y^i v(\theta, y) + \sum_{l=0}^i g_{l,i}(\varphi, x) \partial_y^l v(\theta, y), \quad |g_{l,i}|_{s_n-2\sigma_n}^{\mathcal{O}^{(n)}} \lesssim \sigma_n^{-(i+2)} |\xi^{(n+1)}|_{s_n-\sigma_n}^{\mathcal{O}^{(n)}} \quad (3.41)$$

Inserting (3.41) into (3.40) we get

$$\begin{aligned} q_{l,m}^{(n+1)} = \mathbf{r}_{n+1} & \left(T_{n+1}^{-1} q_{l,m}^{(n)} + \sum_{j=0}^4 T_{n+1}^{-1} (q_{l,j}^{(n)} g_{m,j}) + \sum_{i=0}^4 T_{n+1}^{-1} (q_{i,m}^{(n)} g_{l,i}) \right. \\ & \left. + \sum_{\substack{0 \leq i \leq 2, 0 \leq j \leq 3 \\ 0 \leq i+j \leq 4}} T_{n+1}^{-1} (q_{i,j}^{(n)} g_{l,i} g_{m,j}) \right) \end{aligned} \quad (3.42)$$

so that

$$q_{i,j}^{(n+1)} = T_{n+1}^{-1} (q_{i,j}^{(n)} + O(\xi_{n+1})), \quad |T_{n+1}^{-1} O(\xi_{n+1})|_{s_n-3\sigma_n}^{\mathcal{O}^{(n)}} \lesssim \sigma_n^{-\tau_3} \varepsilon_n e^{C\sigma_n^\mu}. \quad (3.43)$$

In order to obtain the bound (3.43) we used the first line of (3.22) to control the sums appearing in (3.42).

Finally, since

$$T_{n+1}^{-1}(q) - q := (1 + \tilde{\xi}_x^{(n+1)})q(\varphi, x) - q(\theta, y)$$

the bound follows. ■

Now, by (3.31a) and (3.34) $f_{n+1} = f_{n+1}(\varphi, x)$ satisfies

$$|f_{n+1}|_{s_n-2\sigma_n}^{\mathcal{O}^{(n)}} \lesssim \sigma_n^{-4} \varepsilon_n^2. \quad (3.44)$$

In Section 5 we prove the existence of a Cantor set $\mathcal{O}^{(n+1)}$ where item 3.(a) of the iterative lemma holds with $n \rightsquigarrow n+1$.

Proposition 3.8. *Assume that*

$$2^n \sigma_n^{-\tau} e^{C\sigma_n^{-\mu}} \varepsilon_n \ll 1, \quad (3.45)$$

with $\tau \geq \tau_2$. Setting $\lambda_3^{(n+1)} := 1 + A_{n+1}$, there exist Lipschitz functions

$$\Omega^{(n+1)}(j) = \lambda_3^{(n+1)} j^3 + \lambda_1^{(n+1)} j + r_j^{(n+1)} \quad (3.46)$$

satisfying

$$|\lambda_1^{(n+1)} - \lambda_1^{(n)}|^{\mathcal{O}^{(n)}}, \sup_{j \in \mathbb{Z} \setminus \{0\}} |r_j^{(n+1)} - r_j^{(n)}|^{\mathcal{O}^{(n)}} \lesssim \sigma_n^{-\tau} \varepsilon_n e^{C\sigma_n^{-\mu}} \quad (3.47)$$

such that setting

$$\mathcal{E}^{(n+1)} := \left\{ \omega \in \mathcal{O}^{(n)} : |\omega \cdot \ell + \Omega^{(n+1)}(j) - \Omega^{(n+1)}(h)| \geq \frac{2\gamma_{n+1}|j^3 - h^3|}{\mathbf{d}(\ell)}, \forall (\ell, h, j) \neq (0, h, h) \right\} \quad (3.48)$$

for $\omega \in \mathcal{E}^{(n+1)}$ there exists an invertible and bounded linear operator $M^{(n+1)}$

$$\|M^{(n+1)} - \mathbf{Id}\|_{s_n - 5\sigma_n}^{\mathcal{E}^{(n+1)}} \leq \sigma_0^{-\tau} e^{C\sigma_0^{-\mu}} \varepsilon_0 \quad (3.49)$$

such that

$$(M^{(n+1)})^{-1} L_{n+1} M^{(n+1)} = D_{n+1} = \text{diag} \left(\omega \cdot \ell + \Omega^{(n+1)}(j) \right)_{(\ell, j) \in \mathbb{Z}_*^\infty \times \mathbb{Z} \setminus \{0\}} \quad (3.50)$$

The assumption (3.45) follows from (3.10), provided that we choose the constants τ, C and ε_0 appropriately.

Remark 3.9. *Note that in the context of [13] Proposition 3.8 is much simpler to prove, because in order to diagonalize the linearized operator one uses tame estimates coming from the Sobolev regularity on the boundary of the domain. Then the smallness conditions are much simpler to handle. Here we have to strongly rely on the fact that L_{n+1} is a “small” unbounded perturbation of L_n in order to show that the operators $M^{(n)}$ and $M^{(n+1)}$ are close to each other. This is a very delicate issue; see Lemma 5.2 and Section 5.3, which are probably the more technical parts of this paper.*

Lemma 3.10 (Homological equation). *Set*

$$\mathcal{U}^{(n+1)} := \left\{ \omega \in \mathcal{O}^{(n)} : |\omega \cdot \ell + \Omega^{(n+1)}(j)| \geq \gamma_{n+1} \frac{|j|^3}{\mathbf{d}(\ell)}, \forall (\ell, j) \neq (0, 0) \right\} \quad (3.51)$$

For $\omega \in \mathcal{O}^{(n+1)} := \mathcal{U}^{(n+1)} \cap \mathcal{E}^{(n+1)}$ one has

$$h_{n+1} := -L_{n+1}^{-1} f_{n+1} \in \mathcal{H}_{s_{n+1}} \quad (3.52)$$

and one has

$$|h_{n+1}|_{s_{n+1}}^{\mathcal{O}^{(n+1)}} \lesssim \exp \left(\tau \sigma_n^{-\frac{1}{\eta}} \ln \left(\frac{\tau}{\sigma_n} \right) \right) |f_{n+1}|_{s_{n+1} + \sigma_n}^{\mathcal{O}^{(n)}}.$$

Proof. The result follows simply by using the definition of $\mathcal{O}^{(n+1)}$ and applying Lemma A.7. ■

Of course from Lemma 3.10 it follows that,

$$|h_{n+1}|_{s_{n+1}}^{\mathcal{O}^{(n+1)}} \lesssim \sigma_n^{-4} e^{C\sigma_n^{-\mu}} \varepsilon_n^2 \quad (3.53)$$

Now we want to show inductively that

$$\sigma_n^{-4} e^{C\sigma_n^{-\mu}} \varepsilon_n^2 \leq \varepsilon_0 e^{-\chi^{n+1}}, \quad \chi = \frac{3}{2} \quad (3.54)$$

for ε_0 small enough.

By the definition of ε_n in (3.6), (3.54) is equivalent to

$$\varepsilon_0 \lesssim \sigma_0^4 n^{-8} e^{\chi^n(2-\chi)-C'n^\mu} \quad (3.55)$$

Since the r.h.s. of (3.55) admits a positive minimum, we can regard it as a smallness condition on ε_0 , which is precisely (3.10).

We now prove (3.11) with $n \rightsquigarrow n+1$. We only prove the bound for the set $\mathcal{E}^{(n)} \setminus \mathcal{E}^{(n+1)}$. The other one can be proved by similar arguments (it is actually even easier). Let us start by writing

$$\begin{aligned} \mathcal{E}^{(n)} \setminus \mathcal{E}^{(n+1)} &= \bigcup_{(\ell, j, j') \neq (0, j, j)} \mathcal{R}(\ell, j, j'), \\ \mathcal{R}(\ell, j, j') &:= \left\{ \omega \in \mathcal{E}^{(n)} : |\omega \cdot \ell + \Omega^{(n+1)}(j) - \Omega^{(n+1)}(j')| < \frac{2\gamma_{n+1}|j^3 - j'^3|}{\mathfrak{d}(\ell)} \right\}, \\ \forall (\ell, j, j') &\in \mathbb{Z}_*^\infty \times (\mathbb{Z} \setminus \{0\}) \times (\mathbb{Z} \setminus \{0\}), \quad (\ell, j, j') \neq (0, j, j). \end{aligned} \quad (3.56)$$

Lemma 3.11. *Denote $|\ell|_1$ as in (1.5) with $\eta \rightsquigarrow 1$. For any $(\ell, j, j') \neq (0, j, j)$ such that $|\ell|_1 \leq n^2$, one has that $\mathcal{R}(\ell, j, j') = \emptyset$.*

Proof. Let $(\ell, j, j') \in \mathbb{Z}_*^\infty \times (\mathbb{Z} \setminus \{0\}) \times (\mathbb{Z} \setminus \{0\})$, $(\ell, j, j') \neq (0, j, j)$, $|\ell|_1 \leq n^2$. If $j = j'$, clearly $\ell \neq 0$ and $\mathcal{R}(\ell, 0, 0) = \emptyset$ because $\omega \in \mathbb{D}_{\bar{\gamma}}$ with $\bar{\gamma} > 2\gamma_{n+1}$; recall (3.6). Hence we are left to analyze the case $j \neq j'$.

By (3.47), for any $j, j' \in \mathbb{Z} \setminus \{0\}$, $j \neq j'$

$$\left| \left(\Omega^{(n+1)}(j) - \Omega^{(n+1)}(j') \right) - \left(\Omega^{(n)}(j) - \Omega^{(n)}(j') \right) \right| \lesssim \sigma_n^{-\tau} \varepsilon_n e^{C\sigma_n^{-\mu}} |j^3 - j'^3|. \quad (3.57)$$

Therefore, for any $\omega \in \mathcal{E}^{(n)}$

$$\begin{aligned}
|\omega \cdot \ell + \Omega^{(n+1)}(j) - \Omega^{(n+1)}(j')| &\geq |\omega \cdot \ell + \Omega^{(n)}(j) - \Omega^{(n)}(j')| \\
&\quad - \left| \left(\Omega^{(n+1)}(j) - \Omega^{(n+1)}(j') \right) - \left(\Omega^{(n)}(j) - \Omega^{(n)}(j') \right) \right| \\
&\geq \frac{2\gamma_n |j^3 - j'^3|}{\mathbf{d}(\ell)} - C\sigma_n^{-\tau} \varepsilon_n e^{C\sigma_n^{-\mu}} |j^3 - j'^3| \\
&\geq \frac{2\gamma_{n+1} |j^3 - j'^3|}{\mathbf{d}(\ell)}
\end{aligned} \tag{3.58}$$

where in the last inequality we used (3.6) and the fact that, by (A.4) one has

$$\sigma_n^{-\tau} \varepsilon_n e^{C\sigma_n^{-\mu}} \mathbf{d}(\ell) \leq \sigma_n^{-\tau} \varepsilon_n e^{C\sigma_n^{-\mu}} (1+n^2)^{C(1)n} \leq \gamma_0 2^{-n}.$$

The estimate (3.58) clearly implies that $\mathcal{R}(\ell, j, j') = \emptyset$ for $|\ell|_1 \leq n^2$. \blacksquare

Lemma 3.12. *Let $\mathcal{R}(\ell, j, j') \neq \emptyset$. Then $\ell \neq 0$, $|j^3 - j'^3| \lesssim \|\ell\|_1$ and $\mathbb{P}(\mathcal{R}(\ell, j, j')) \lesssim \frac{\gamma_{n+1}}{\mathbf{d}(\ell)}$*

Proof. The proof is identical to the one for Lemma 6.2 in [21], simply replacing j^2 with j^3 . \blacksquare

By (3.56) and collecting Lemmata 3.11, 3.12, one obtains that

$$\mathbb{P}(\mathcal{E}^{(n)} \setminus \mathcal{E}^{(n+1)}) \lesssim \sum_{\substack{|\ell|_1 \geq n^2 \\ |j|, |j'| \leq C\|\ell\|_1}} \frac{\gamma_{n+1}}{\mathbf{d}(\ell)} \lesssim \gamma_{n+1} \sum_{|\ell|_1 \geq n^2} \frac{\|\ell\|_1^2}{\mathbf{d}(\ell)} \lesssim \gamma_{n+1} n^{-2} \sum_{\ell \in \mathbb{Z}_*^\infty} \frac{|\ell|_1^3}{\mathbf{d}(\ell)} \lesssim \gamma_{n+1} n^{-2}. \tag{3.59}$$

where in the last inequality we used Lemma A.8. Thus (3.11) follows.

We now the convergence of the scheme. Precisely we show that the series (3.25) converges totally in $\mathcal{H}_{\bar{s}}$. Note that

$$|T_i u|_{\bar{s}}^{\mathcal{O}(\infty)} \leq (1 + 2^{-i}) |u|_{\bar{s} + \sigma_i}^{\mathcal{O}(\infty)} \leq 2 |u|_{\bar{s} + \sigma_i}^{\mathcal{O}(\infty)}. \tag{3.60}$$

Thuse, using (3.60) into (3.25) we get

$$|u_n|_{\bar{s}}^{\mathcal{O}(\infty)} \leq |h_0|_{\bar{s}}^{\mathcal{O}(\infty)} + \sum_{j=1}^n 2^j |h_j|_{\bar{s} + (\sigma_1 + \dots + \sigma_j)}^{\mathcal{O}(\infty)} \tag{3.61}$$

Now since

$$\bar{s} + \sum_{n=1}^{\infty} \sigma_n = \bar{s} + \frac{6\sigma_{-1}}{\pi^2} \sum_{n \geq 1} \frac{1}{n^2} = s_\infty \leq s_j \tag{3.62}$$

we deduce that $u_\infty \in \mathcal{H}_{\bar{s}}$. Finally by continuity

$$F(u_\infty) = \lim_{n \rightarrow \infty} F(u_n) = \lim_{n \rightarrow \infty} T_1^{-1} T_2^{-1} \dots T_n^{-1} F_n(h_n) = 0.$$

so the assertion follows since (recall $\bar{s} := s_\infty - \sum_{n \geq 1} \sigma_n$ and (3.62))

$$|T_1^{-1} T_2^{-1} \dots T_n^{-1} F_n(h_n)|_{\bar{s}}^{\mathcal{O}^{(\infty)}} \leq 2^n \sigma_n^{-4} \varepsilon_n^2.$$

We finally conclude the proof of Proposition 3.1 by showing that (3.26) holds.

First of all, reasoning as in Lemma 3.12 and using Lemma A.8, we see that

$$\mathbb{P}(\mathcal{O}^{(0)}) = 1 - O(\gamma_0)$$

Then

$$\mathbb{P}(\mathcal{O}^{(\infty)}) = \mathbb{P}(\mathcal{O}^{(0)}) - \sum_{n \geq 0} \mathbb{P}(\mathcal{O}^{(n)} \setminus \mathcal{O}^{(n+1)})$$

so that (3.26) follows by (3.11). ■

4 Proof of Proposition 3.6

In order to prove Proposition 3.6, we start by dropping the index n , i.e. we set $\mathcal{L} \equiv L_n$ (see (3.18)) and $\mathcal{Q} \equiv Q'_n(h_n)$ (see (3.34)).

More generally, we consider a Hamiltonian operator of the form

$$\begin{aligned} \mathcal{L}^{(0)} &= \mathcal{L} + \mathcal{Q} \\ \mathcal{L} &:= \omega \cdot \partial_\varphi + \lambda_3 \partial_x^3 + a_1(\varphi, x) \partial_x + a_0(\varphi, x), \\ \mathcal{Q} &:= d_3(\varphi, x) \partial_x^3 + d_2(\varphi, x) \partial_x^2 + d_1(\varphi, x) \partial_x + d_0(\varphi, x) \end{aligned} \tag{4.1}$$

defined for all $\omega \in \Omega \subseteq \mathbb{D}_\gamma$ and $\lambda_3, a_0, a_1, d_0, \dots, d_3$ satisfy the following properties.

1. There is δ_0 small enough such that

$$|\lambda_3 - 1|^\Omega \leq \delta_0 \tag{4.2}$$

2. There is $\rho > 0$ such that $a_i \in \mathcal{H}(\mathbb{T}_\rho^\infty \times \mathbb{T}_\rho)$ and

$$|a_i|_\rho^\Omega \leq \delta_0, \quad i = 0, 1 \tag{4.3}$$

and moreover

$$\lambda_1 := \frac{1}{2\pi} \int_{\mathbb{T}} a_1(\varphi, x) dx \tag{4.4}$$

i.e. it does not depend on φ .

3. $d_0, \dots, d_3 \in \mathcal{H}(\mathbb{T}_\rho^\infty \times \mathbb{T}_\rho)$ (note that by the Hamiltonian structure $d_2 = \partial_x d_3$) and they satisfy the estimate

$$|d_i|_\rho^\Omega \lesssim \delta, \tag{4.5}$$

for some $\delta \ll \min\{\delta_0, \rho\}$.

Let us now choose ζ such that $0 < \zeta \ll \rho$ and

$$\zeta^{-\tau'} e^{2c_0\zeta^{-\mu}} \delta \ll 1. \quad (4.6)$$

for some $\tau' > 0$. We shall conjugate $\mathcal{L}^{(0)}$ to a new operator $\frac{1}{\zeta}\mathcal{L}_+$ with $\mathbf{r} = \mathbf{r}(\varphi)$ an explicit function with

$$\mathcal{L}_+ = \omega \cdot \partial_\varphi + \lambda_3^+ \partial_x^3 + a_1^+(\varphi, x) \partial_x + a_0^+(\varphi, x) \quad (4.7)$$

with the coefficients satisfying

$$|\lambda_3^+ - \lambda_3|^\Omega \lesssim \delta \quad (4.8)$$

and

$$|a_i^+ - a_i|_{\rho-2\zeta}^\Omega \leq \zeta^{-\tau'} e^{2c_0\zeta^{-\mu}} \delta, \quad \lambda_1 := \frac{1}{2\pi} \int_{\mathbb{T}} a_1(\varphi, x) dx. \quad (4.9)$$

This will allow us to conclude the proof of Proposition 3.6.

4.1 Elimination of the x -dependence from the highest order term

Consider an analytic function $\alpha(\varphi, x)$ (to be determined) and let

$$\mathcal{T}_1 u(\varphi, x) := (1 + \alpha_x(\varphi, x))(\mathcal{A}u)(\varphi, x), \quad \mathcal{A}u(\varphi, x) := u(\varphi, x + \alpha(\varphi, x)).$$

We choose $\alpha(\varphi, x)$ and $m_3(\varphi)$ in such a way that

$$(\lambda_3 + d_3(\varphi, x))(1 + \alpha_x(\varphi, x))^3 = m_3(\varphi), \quad (4.10)$$

which implies

$$\alpha(\varphi, x) := \partial_x^{-1} \left[\frac{m_3(\varphi)^{\frac{1}{3}}}{(\lambda_3 + d_3(\varphi, x))^{\frac{1}{3}}} - 1 \right], \quad m_3(\varphi) := \left(\frac{1}{2\pi} \int_{\mathbb{T}} \frac{dx}{(\lambda_3 + d_3(\varphi, x))^{\frac{1}{3}}} \right)^{-3}. \quad (4.11)$$

By (4.2), (4.5) and Lemma A.5 one has

$$|m_3 - \lambda_3|_\rho^\Omega, |\alpha|_\rho^\Omega \lesssim \delta \quad (4.12)$$

Note that for any $0 < \zeta \ll \rho$ such that $\delta\zeta^{-1} \ll 1$, by Lemma A.1, $x \mapsto x + \alpha(\varphi, x)$ is invertible and the inverse is given by $y \mapsto y + \tilde{\alpha}(\varphi, y)$ with

$$\tilde{\alpha} \in \mathcal{H}(\mathbb{T}_{\rho-\zeta}^\infty \times \mathbb{T}_{\rho-\zeta}), \quad |\tilde{\alpha}|_{\rho-\zeta}^\Omega, |\alpha|_\rho^\Omega \lesssim \delta. \quad (4.13)$$

A direct calculations shows that

$$\mathcal{A}^{-1}u(\varphi, y) = u(\varphi, y + \tilde{\alpha}(\varphi, y)), \quad \mathcal{T}_1^{-1} = (1 + \tilde{\alpha}_y)\mathcal{A}^{-1} \quad (4.14)$$

and the following conjugation rules hold:

$$\begin{aligned}
\mathcal{T}_1^{-1} a(\varphi, x) \mathcal{T}_1 &= \mathcal{A}^{-1} a(\varphi, x) \mathcal{A} = (\mathcal{A}^{-1} a)(\varphi, y), \\
\mathcal{T}_1^{-1} \partial_x \mathcal{T}_1 &= \left(1 + \mathcal{A}^{-1}(\alpha_x)\right) \partial_y + (1 + \tilde{\alpha}_y) \mathcal{A}^{-1}(\alpha_{xx}), \\
\mathcal{T}_1^{-1} \omega \cdot \partial_\varphi \mathcal{T}_1 &= \omega \cdot \partial_\varphi + \mathcal{A}^{-1}(\omega \cdot \partial_\varphi \alpha) \partial_y + (1 + \tilde{\alpha}_y) \mathcal{A}^{-1}(\omega \cdot \partial_\varphi \alpha_x).
\end{aligned} \tag{4.15}$$

Clearly one can get similar conjugation formulae for higher order derivatives, having expression similar to (3.41). In conclusion

$$\begin{aligned}
\mathcal{L}^{(1)} &:= \mathcal{T}_1^{-1}(\mathcal{L} + \mathcal{Q})\mathcal{T}_1 \\
&= \omega \cdot \partial_\varphi + \mathcal{A}^{-1} \left[(\lambda_3 + q_3)(1 + \alpha_x)^3 \right] \partial_y^3 + b_2(\varphi, y) \partial_y^2 + b_1(\varphi, y) \partial_y + b_0(\varphi, y) \\
&= \omega \cdot \partial_\varphi + m_3(\varphi) \partial_x^3 + b_1(\varphi, x) \partial_x + b_0(\varphi, x)
\end{aligned} \tag{4.16}$$

for some (explicitly computable) coefficients b_i , where in the last equality we used (4.10) and the fact that \mathcal{T}_1 is symplectic, so that $b_2(\varphi, x) = 2\partial_x m_3(\varphi) = 0$.

Furthermore, the estimates (4.2), (4.3), (4.12), (4.13), Corollary A.2 and Lemmata A.3, A.4 imply that for $0 < \zeta \ll \rho$

$$|b_i|_{\rho-2\zeta}^\Omega \lesssim \delta_0, \quad |b_i - a_i|_{\rho-2\zeta}^\Omega \lesssim \zeta^{-\tau} \delta, \quad \text{for some } \tau > 0. \tag{4.17}$$

4.2 Elimination of the φ -dependence from the highest order term

We now consider a quasi periodic reparametrization of time of the form

$$\mathcal{T}_2 u(\varphi, x) := u(\varphi + \omega \beta(\varphi), x) \tag{4.18}$$

where $\beta : \mathbb{T}_{\rho-\zeta}^\infty \rightarrow \mathbb{R}$ is an analytic function to be determined. Precisely we choose $\lambda_3^+ \in \mathbb{R}$ and $\beta(\varphi)$ in such a way that

$$\lambda_3^+ \left(1 + \omega \cdot \partial_\varphi \beta(\varphi)\right) = m_3(\varphi), \tag{4.19}$$

obtaining thus

$$\lambda_3^+ := \int_{\mathbb{T}^\infty} m_3(\varphi) d\varphi, \quad \beta(\varphi) := (\omega \cdot \partial_\varphi)^{-1} \left[\frac{m_3}{\lambda_3^+} - 1 \right] \tag{4.20}$$

where we recall the definition A.3. By the estimates (4.12) and by Lemma 3.3, one obtains that for $0 < \zeta \ll \rho$

$$|\lambda_3^+ - \lambda_3|^\Omega \lesssim \delta, \quad |\beta|_{\rho-\zeta}^\Omega \lesssim e^{c_0 \zeta^{-\mu}} \delta. \tag{4.21}$$

By Lemma A.1 and (4.6) we see that $\varphi \mapsto \varphi + \omega \beta(\varphi)$ is invertible and the inverse is given by $\vartheta \mapsto \vartheta + \omega \tilde{\beta}(\vartheta)$ with

$$\tilde{\beta} \in \mathcal{H}(\mathbb{T}_{\rho-2\zeta}^\infty), \quad |\tilde{\beta}|_{\rho-2\zeta}^\Omega \lesssim e^{c_0 \zeta^{-\mu}} \delta. \tag{4.22}$$

The inverse of the operator \mathcal{T}_2 is then given by

$$\mathcal{T}_2^{-1} u(\vartheta, x) = u(\vartheta + \omega \tilde{\beta}(\vartheta), x). \tag{4.23}$$

so that

$$\begin{aligned}\mathcal{T}_2^{-1}\mathcal{L}^{(1)}\mathcal{T}_2 &= \mathcal{T}_2^{-1}(1 + \omega \cdot \partial_\varphi\beta)\omega \cdot \partial_\vartheta + \mathcal{T}_2^{-1}(m_3)\partial_x^3 + \mathcal{T}_2^{-1}(b_1)\partial_x + \mathcal{T}_2^{-1}(b_0) \\ &=: \frac{1}{\mathbf{r}}\mathcal{L}^{(2)}\end{aligned}\tag{4.24}$$

where

$$\begin{aligned}\mathcal{L}^{(2)} &:= \omega \cdot \partial_\vartheta + \lambda_3^+\partial_x^3 + c_1(\vartheta, x)\partial_x + c_0(\vartheta, x), \\ \mathbf{r} &:= \frac{1}{\mathcal{T}_2^{-1}(1 + \omega \cdot \partial_\varphi\beta)} \stackrel{(4.19)}{=} \frac{\lambda_3^+}{\mathcal{T}_2^{-1}(m_3)}, \\ c_i &:= \mathbf{r}\mathcal{T}_2^{-1}(b_i), \quad i = 1, 0.\end{aligned}\tag{4.25}$$

Therefore by the estimates (4.12), (4.21), (4.22) and by applying Corollary A.2, Lemma A.5, and (4.6), one gets

$$\begin{aligned}|\mathbf{r} - 1|_{\rho-\zeta}^\Omega &\lesssim \delta \\ |c_i - a_i|_{\rho-\zeta}^\Omega &\lesssim \zeta^{-\tau} e^{c_0\zeta^{-\mu}} \delta, \quad i = 0, 1.\end{aligned}\tag{4.26}$$

4.3 Time dependent traslation of the space variable

Let $p : \mathbb{T}_{\rho-2\zeta}^\infty \rightarrow \mathbb{R}$ be an analytic function to be determined and let

$$\mathcal{T}_3 u(\varphi, x) := u(\varphi, x + p(\varphi)), \quad \text{with inverse} \quad \mathcal{T}_3^{-1}u(\varphi, y) = u(\varphi, y - p(\varphi)).\tag{4.27}$$

Computing explicitly

$$\begin{aligned}\mathcal{L}^{(3)} &:= \mathcal{T}_3^{-1}\mathcal{L}^{(2)}\mathcal{T}_3 = \omega \cdot \partial_\varphi + \lambda_3^+\partial_x^3 + a_1^+(\varphi, x)\partial_x + a_0^+(\varphi, x), \\ a_1^+ &:= \omega \cdot \partial_\varphi p + \mathcal{T}_3^{-1}(c_1), \quad a_0^+ := \mathcal{T}_3^{-1}(c_0),\end{aligned}\tag{4.28}$$

and by (4.4) one has

$$\begin{aligned}\frac{1}{2\pi} \int_{\mathbb{T}} \mathcal{T}_3^{-1}(c_1)(\varphi, y) dy &= \frac{1}{2\pi} \int_{\mathbb{T}} c_1(\varphi, x) dx \\ &= \frac{1}{2\pi} \int_{\mathbb{T}} a_1(\varphi, x) dx + \frac{1}{2\pi} \int_{\mathbb{T}} (c_1 - a_1)(\varphi, x) dx \\ &= \lambda_1 + \frac{1}{2\pi} \int_{\mathbb{T}} (c_1 - a_1)(\varphi, x) dx.\end{aligned}\tag{4.29}$$

We want to choose $p(\varphi)$ in such a way that the x -average of d_1 is constant. To this purpose we define

$$p(\varphi) := (\omega \cdot \partial_\varphi)^{-1} \left[\langle (c_1 - a_1) \rangle_{\varphi, x} - \frac{1}{2\pi} \int_{\mathbb{T}} (c_1 - a_1)(\varphi, x) dx \right]\tag{4.30}$$

where for any $a : \mathbb{T}_\sigma^\infty \times \mathbb{T}_\sigma \rightarrow \mathbb{C}$, $\langle a \rangle_{\varphi, x}$ is defined by

$$\langle a \rangle_{\varphi, x} := \frac{1}{(2\pi)} \int_{\mathbb{T}} \int_{\mathbb{T}^\infty} a(\varphi, x) d\varphi dx$$

(recall the definition A.3). By (4.26) and Lemma 3.3 one gets

$$|p|_{\rho-2\zeta}^{\Omega} \lesssim \zeta^{-\tau} e^{2c_0\zeta^{-\mu}} \delta \stackrel{(4.6)}{\ll} \zeta. \quad (4.31)$$

Moreover

$$\lambda_1^+ := \frac{1}{2\pi} \int_{\mathbb{T}} d_1(\varphi, x) dx = \lambda_1 + \langle (c_1 - a_1) \rangle_{\varphi, x}. \quad (4.32)$$

Finally using (4.26), (A.2) (with $\Phi_\alpha = \mathcal{T}_3^{-1}$), (4.31), one gets

$$|a_i^+ - a_i|_{\rho-2\zeta}^{\Omega} \lesssim \zeta^{-\tau'} e^{2c_0\zeta^{-\mu}} \delta, \quad (4.33)$$

for some $\tau' > 0$.

4.4 Conclusion of the proof

We start by noting that $\mathcal{T} := \mathcal{T}_3 \circ \mathcal{T}_2 \circ \mathcal{T}_1$ has the form (3.33) with $p^{(n+1)} = p$, $\beta^{(n+1)} = \beta$ and $\xi^{(n+1)}(\varphi, x) = \alpha(\varphi + \omega\beta(\varphi), x + p(\varphi))$. Hence, setting $\mathbf{r} := \mathbf{r}_{n+1}$, $\rho := s_n - \sigma_n$, $\delta := \sigma_n^{-4}\varepsilon_n$, $\delta_0 := 2\varepsilon_0$ and $\zeta := \sigma_n$ we denote

$$1 + A_{n+1} = \lambda_3^+, \quad B_{n+1}(\varphi, x) := a_1^+(\varphi, x), \quad C_{n+1} = a_0^+(\varphi, x),$$

and thus Proposition 3.6 follows. ■

5 Proof of Proposition 3.8

In order to prove Proposition 3.8, we start by considering a linear Hamiltonian operator defined for $\omega \in \mathcal{O} \subseteq \mathcal{D}_\gamma$ of the form

$$\mathcal{L} = \mathcal{L}(\lambda_3, a_1, a_0) := \omega \cdot \partial_\varphi + \lambda_3 \partial_x^3 + a_1(\varphi, x) \partial_x + a_0(\varphi, x). \quad (5.1)$$

We want to show that, for any choice of the coefficients λ_3, a_1, a_0 satisfying some hypotheses (see below), it is possible to reduce \mathcal{L} to constant coefficients. Moreover we want to show that such reduction is ‘‘Lipshitz’’ w.r.t. the parameters λ_3, a_1, a_0 , in a sense that will be clarified below.

Regarding the coefficients, we need to require that

$$\begin{aligned} a_i &:= \sum_{k=0}^m a_i^{(k)}, \quad |a_i^{(k)}|_{\rho_k}^{\mathcal{O}} \lesssim \delta_k, \quad \forall k = 0, \dots, m, \quad i = 0, 1, \\ |\lambda_3 - 1|^{\mathcal{O}} &\lesssim \delta_0, \\ \lambda_1 &\equiv \lambda_1(a_1) = \sum_{k=0}^m \lambda_1^{(k)}, \quad \lambda_1^{(k)} := \frac{1}{2\pi} \int_{\mathbb{T}} a_1^{(k)}(\varphi, x) dx = \text{const}. \end{aligned} \quad (5.2)$$

for some $0 < \dots < \rho_m < \dots < \rho_0$ and $0 < \dots \ll \delta_m \ll \dots \ll \delta_0 \ll 1$ so that there is a third sequence ζ_i such that $0 < \zeta_i < \rho_i$ and

$$\sum_{i \geq 0} \zeta_i^{-\tau} e^{C\zeta_i^{-\mu}} \delta_i \lesssim \delta_0, \quad (5.3)$$

for some $\tau, C > 0$.

5.1 Reduction of the first order term

We consider an operator \mathcal{L} of the form (5.1) satisfying the hypotheses above. We start by showing that it is possible to reduce it to constant coefficients up to a bounded reminder, and that such reduction is ‘‘Lipshitz’’ w.r.t. the parameters λ_3, a_1, a_0 .

Lemma 5.1. *There exists a symplectic invertible operator $\mathcal{M} = \exp(\mathcal{G})$, with $\mathcal{G} \equiv \mathcal{G}(\lambda_3, a_1)$ and an operator $\mathcal{R}_0 \equiv \mathcal{R}_0(\lambda_3, a_1, a_0)$ satisfying*

$$\begin{aligned} \mathcal{G} &= \sum_{i=0}^m \mathcal{G}^{(i)}, \quad \|\mathcal{G}^{(i)}\|_{\rho_i, -1}^{\mathcal{O}} \lesssim \delta_i, \\ \mathcal{R}_0 &= \sum_{i=0}^m \mathcal{R}_0^{(i)}, \quad \|\mathcal{R}_0^{(i)}\|_{\rho_i - \zeta_i}^{\mathcal{O}} \lesssim \zeta_i^{-\tau} e^{C\zeta_i^{-\mu}} \delta_i \end{aligned} \quad (5.4)$$

for some $C, \tau \gg 1$, such that

$$\mathcal{L}_0 := \mathcal{M}^{-1} \mathcal{L} \mathcal{M} = \omega \cdot \partial_\varphi + \lambda_3 \partial_x^3 + \lambda_1 \partial_x + \mathcal{R}_0. \quad (5.5)$$

Proof. We look for \mathcal{G} of the form

$$\mathcal{G} = \pi_0^\perp g(\varphi, x) \partial_x^{-1}$$

and we choose the function $g(\varphi, x)$ where $g = g(\lambda_3, a_1)$ in order to solve

$$3\lambda_3 \partial_x g(\varphi, x) + a_1(\varphi, x) = \lambda_1. \quad (5.6)$$

By (5.2), one obtains that

$$g := \frac{1}{3\lambda_3} \partial_x^{-1} [\lambda_1 - a_1] \quad (5.7)$$

and therefore

$$\begin{aligned} g &= \sum_{i=0}^m g_i, \quad g_i := \frac{1}{3\lambda_3} \partial_x^{-1} [\lambda_1^{(i)} - a_1^{(i)}], \\ |g_i|_{\rho_i}^{\mathcal{O}} &\lesssim \delta_i, \quad i = 0, \dots, m. \end{aligned} \quad (5.8)$$

Of course we can also write the operator $\mathcal{G} := \pi_0^\perp g(\varphi, x) \partial_x^{-1} = \sum_{i=0}^m \mathcal{G}_i$ where $\mathcal{G}_i := \pi_0^\perp g_i(\varphi, x) \partial_x^{-1}$ and one has

$$\|\mathcal{G}_i\|_{\rho_i, -1}^{\mathcal{O}} \lesssim \delta_i, \quad i = 0, \dots, m. \quad (5.9)$$

Again by (5.2), defining $\mathcal{P} := a_1 \partial_x + a_0$, one has that $\mathcal{P} = \sum_{i=0}^m \mathcal{P}_i$, where $\mathcal{P}_i := a_1^{(i)} \partial_x + a_0^{(i)}$ satisfies

$$\|\mathcal{P}_i\|_{\rho_i, 1}^{\mathcal{O}} \lesssim \delta_i. \quad (5.10)$$

Therefore

$$\begin{aligned} \mathcal{L}_0 &= \mathcal{M}^{-1} \mathcal{L} \mathcal{M} = e^{-\mathcal{G}} \omega \cdot \partial_\varphi e^{\mathcal{G}} + \lambda_3 e^{-\mathcal{G}} \partial_x^3 e^{\mathcal{G}} + e^{-\mathcal{G}} \mathcal{P} e^{\mathcal{G}} \\ &= \omega \cdot \partial_\varphi + \lambda_3 \partial_x^3 + \left(3\lambda_3 g_x + a_1\right) \partial_x + \mathcal{R}_0 \\ &\stackrel{(5.6)}{=} \omega \cdot \partial_\varphi + \lambda_3 \partial_x^3 + \lambda_1 \partial_x + \mathcal{R}_0 \end{aligned} \quad (5.11)$$

where

$$\mathcal{R}_0 := \left(e^{-\mathcal{G}} \omega \cdot \partial_\varphi e^{\mathcal{G}} - \omega \cdot \partial_\varphi\right) + \lambda_3 \left(e^{-\mathcal{G}} \partial_x^3 e^{\mathcal{G}} - \partial_x^3 - 3g_x \partial_x\right) + \left(e^{-\mathcal{G}} \mathcal{P} e^{\mathcal{G}} - \mathcal{P}\right) + a_0. \quad (5.12)$$

Then (5.3), (5.9), (5.10) guarantee that the hypotheses of Lemmata A.10-A.11 are verified. Hence, we apply Lemma A.10-(ii) to expand the operator $e^{-\mathcal{G}} \mathcal{P} e^{\mathcal{G}} - \mathcal{P}$, Lemma A.11-(ii) to expand $e^{-\mathcal{G}} \partial_x^3 e^{\mathcal{G}} - \partial_x^3 - 3g_x \partial_x$ and Lemma A.11-(iii) to expand $e^{-\mathcal{G}} \omega \cdot \partial_\varphi e^{\mathcal{G}} - \omega \cdot \partial_\varphi$. The expansion of the multiplication operator a_0 is already provided by (5.2). Hence, one obtains that there exist $C, \tau \gg 1$ such that (5.4) is satisfied. \blacksquare

We now consider a ‘‘small modification’’ of the operator \mathcal{L} in the following sense. We consider an operator

$$\mathcal{L}^+ = \mathcal{L}(\lambda_3^+, a_1^+, a_0^+) := \omega \cdot \partial_\varphi + \lambda_3^+ \partial_x^3 + a_1^+(\varphi, x) \partial_x + a_0^+(\varphi, x) \quad (5.13)$$

with

$$\frac{1}{2\pi} \int_{\mathbb{T}} a_1^+(\varphi, x) dx =: \lambda_1^+ = \text{const}, \quad |a_i^+ - a_i|_{\rho_{m+1}}, |\lambda_3^+ - \lambda_3| \lesssim \delta_{m+1}. \quad (5.14)$$

Of course we can apply Lemma 5.1 and conjugate \mathcal{L}^+ to

$$\mathcal{L}_0^+ := \omega \cdot \partial_\varphi + \lambda_3^+ \partial_x^3 + \lambda_1^+ \partial_x + \mathcal{R}_0^+ \quad (5.15)$$

with \mathcal{R}_0^+ a bounded operator. We want to show that \mathcal{L}_0^+ is ‘‘close’’ to \mathcal{L}_0 , namely the following result.

Lemma 5.2. *One has*

$$|\lambda_1^+ - \lambda_1| \lesssim \delta_{m+1}, \quad \|\mathcal{R}_0^+ - \mathcal{R}_0\|_{\rho_{m+1} - \zeta_{m+1}} \lesssim \zeta_{m+1}^{-\tau} e^{C\zeta_{m+1}^{-\mu}} \delta_{m+1}. \quad (5.16)$$

Proof. The first bound follows trivially from (5.14). Regarding the second bound one can reason as follows. As in Lemma 5.1, one can define $\mathcal{G}^+ := \pi_0^\perp g^+(\varphi, x) \partial_x^{-1}$ with

$$g^+ := \frac{1}{3\lambda_3^+} \partial_x^{-1} \left[\lambda_1^+ - a_1^+ \right] \quad (5.17)$$

so that

$$\|\mathcal{G}^+ - \mathcal{G}\|_{\rho_{m+1}, -1} \lesssim \delta_{m+1}. \quad (5.18)$$

Defining $\mathcal{P}^+ := a_1^+ \partial_x + a_0^+$ and recalling that $\mathcal{P} := a_1 \partial_x + a_0$, by (5.14), one gets

$$\|\mathcal{P}^+ - \mathcal{P}\|_{\rho_{m+1},1} \lesssim \delta_{m+1}. \quad (5.19)$$

The estimate on $\mathcal{R}_0^+ - \mathcal{R}_0$ follows by applying Lemmata A.13, A.14, and by the estimates (5.14), (5.19), (5.18). \blacksquare

5.2 Reducibility

We now consider an operator \mathcal{L}_0 of the form

$$\mathcal{L}_0 \equiv \mathcal{L}_0(\lambda_1, \lambda_3, \mathcal{P}_0) := \omega \cdot \partial_\varphi + \mathcal{D}_0 + \mathcal{P}_0 \quad (5.20)$$

with \mathcal{P}_0 a bounded operator and

$$\mathcal{D}_0 \equiv \mathcal{D}_0(\lambda_1, \lambda_3) := i \operatorname{diag}_{j \in \mathbb{Z} \setminus \{0\}} \Omega_0(j), \quad \Omega_0(j) := -\lambda_3 j^3 + \lambda_1 j, \quad j \in \mathbb{Z} \setminus \{0\}, \quad (5.21)$$

and we show that, under some smallness conditions specified below it is possible to reduce it to constant coefficients, and that the reduction is ‘‘Lipschitz’’ w.r.t. the parameters $\lambda_1, \lambda_3, \mathcal{P}_0$.

In order to do so, we introduce three sequences $0 < \dots < \rho_m < \dots < \rho_0$, $0 < \dots \ll \delta_m \ll \dots \ll \delta_0$ and $1 \ll N_0 \ll N_1 \ll \dots$ and we assume that setting $\Delta_i = \rho_i - \rho_{i+1}$ one has

$$\sum_{i \geq 0} \Delta_i^{-\tau} e^{C \Delta_i^{-\mu}} \delta_i \lesssim \delta_0, \quad (5.22)$$

$$e^{-N_k \Delta_k} \delta_k + e^{C \Delta_k^{-\mu}} \delta_k^2 \ll 2^{-k} \delta_{k+1}, \quad (5.23)$$

$$\delta_k \ll (1 + N_k)^{-C N_k^{\frac{1}{1+\eta}}} \quad (5.24)$$

and

$$\begin{aligned} |\lambda_3 - 1|^\mathcal{O}, |\lambda_1|^\mathcal{O} &\leq \delta_0, \\ \mathcal{P}_0 &:= \sum_{i=0}^m \mathcal{P}_0^{(i)}, \quad \|\mathcal{P}_0^{(i)}\|_{\rho_i}^\mathcal{O} \leq \delta_i, \quad i = 0, \dots, m, \end{aligned} \quad (5.25)$$

for some $\tau, C > 0$.

We have the following result.

Lemma 5.3. *Fix $\gamma \in [\gamma_0/2, 2\gamma_0]$. For $k = 0, \dots, m$ there is a sequence of sets $\mathcal{E}_k \subseteq \mathcal{E}_{k-1}$ and a sequence of symplectic maps Φ_k defined for $\omega \in \mathcal{E}_{k+1}$ such that setting \mathcal{L}_0 as in (5.20) and for $k \geq 1$,*

$$\mathcal{L}_k := \Phi_{k-1}^{-1} \mathcal{L}_{k-1} \Phi_{k-1}, \quad (5.26)$$

one has the following.

1. \mathcal{L}_k is of the form

$$\mathcal{L}_k := \omega \cdot \partial_\varphi + \mathcal{D}_k + \mathcal{P}_k \quad (5.27)$$

where

- The operator \mathcal{D}_k is of the form

$$\mathcal{D}_k := \text{diag}_{j \in \mathbb{Z} \setminus \{0\}} \Omega_k(j), \quad \Omega_k(j) = \Omega_0(j) + r_k(j) \quad (5.28)$$

with $r_0(j) = 0$ and for $k \geq 1$, $r_k(j)$ is defined for $\omega \in \mathcal{E}_0 = \mathcal{O}$ and satisfies

$$\sup_{j \in \mathbb{Z} \setminus \{0\}} |r_k(j) - r_{k-1}(j)|^{\mathcal{O}} \leq \delta_{k-1} \sum_{i=1}^{k-1} 2^{-i}. \quad (5.29)$$

- The operator \mathcal{P}_k is such that

$$\text{for } 0 \leq k \leq m, \quad \mathcal{P}_k = \sum_{i=k}^m \mathcal{P}_k^{(i)}, \quad \|\mathcal{P}_k^{(i)}\|_{\rho_i}^{\mathcal{E}_k} \leq \delta_i \sum_{j=1}^k 2^{-j}, \quad \forall i = k, \dots, m. \quad (5.30)$$

2. One has $\Phi_{k-1} := \exp(\Psi_{k-1})$, such that

$$\|\Psi_{k-1}\|_{\rho_k}^{\mathcal{E}_k} \lesssim e^{C\Delta_{k-1}^{-\mu}} \|\mathcal{P}_{k-1}^{(k-1)}\|_{\rho_{k-1}}^{\mathcal{E}_{k-1}} \lesssim e^{C\Delta_{k-1}^{-\mu}} \delta_{k-1} \quad (5.31)$$

3. The sets \mathcal{E}_k are defined as

$$\mathcal{E}_k := \left\{ \omega \in \mathcal{E}_{k-1} : |\omega \cdot \ell + \Omega_{k-1}(j) - \Omega_{k-1}(j')| \geq \frac{\gamma |j^3 - j'^3|}{\mathbf{d}(\ell)}, \right. \\ \left. \forall (\ell, j, j') \neq (0, j, j), \quad |\ell|_\eta \leq N_{k-1} \right\}. \quad (5.32)$$

Proof. The statement is trivial for $k = 0$ so we assume it to hold up to $k < m$ and let us prove it for $k + 1$. For any $\Phi_k := \exp(\Psi_k)$ one has

$$\mathcal{L}_{k+1} = \Phi_k^{-1} \mathcal{L}_k \Phi_k = \omega \cdot \partial_\varphi + \mathcal{D}_k + \omega \cdot \partial_\varphi \Psi_k + [\mathcal{D}_k, \Psi_k] + \Pi_{N_k} \mathcal{P}_k^{(k)} + \mathcal{P}_{k+1} \quad (5.33)$$

where the operator \mathcal{P}_{k+1} is defined by

$$\mathcal{P}_{k+1} := \Pi_{N_k}^\perp \mathcal{P}_k^{(k)} + \sum_{p \geq 2} \frac{\text{Ad}_{\Psi_k}^p(\omega \cdot \partial_\varphi + \mathcal{D}_k)}{p!} + \sum_{i=k+1}^m e^{-\Psi_k} \mathcal{P}_k^{(i)} e^{\Psi_k} + \sum_{p \geq 1} \frac{\text{Ad}_{\Psi_k}^p(\mathcal{P}_k^{(k)})}{p!}. \quad (5.34)$$

Then we choose Ψ_k in such a way that

$$\omega \cdot \partial_\varphi \Psi_k + [\mathcal{D}_k, \Psi_k] + \Pi_{N_k} \mathcal{P}_k^{(k)} = \mathcal{Z}_k, \quad (5.35) \\ \mathcal{Z}_k := \text{diag}_{j \in \mathbb{Z} \setminus \{0\}} (\mathcal{P}_k^{(k)})_j^j(0),$$

namely for $\omega \in \mathcal{E}_{k+1}$ we set

$$(\Psi_k)_j^{j'}(\ell) := \begin{cases} \frac{(\mathcal{P}_k^{(k)})_j^{j'}(\ell)}{i(\omega \cdot \ell + \Omega_k(j) - \Omega_k(j'))}, & \forall (\ell, j, j') \neq (0, j, j), \quad |\ell|_\eta \leq N_k, \\ 0 & \text{otherwise.} \end{cases} \quad (5.36)$$

Therefore,

$$|(\Psi_k)_j^{j'}(\ell)| \lesssim \mathbf{d}(\ell) |(\mathcal{P}_k^{(k)})_j^{j'}(\ell)|, \quad \forall \omega \in \mathcal{E}_{k+1}. \quad (5.37)$$

and by applying Lemma A.6, using the induction estimate (5.30), one obtains

$$\|\Psi_k\|_{\rho_k - \zeta}^{\mathcal{E}_{k+1}} \lesssim e^{C\zeta - \mu} \|\mathcal{P}_k^{(k)}\|_{\rho_k}^{\mathcal{E}_k} \stackrel{(5.30)}{\lesssim} e^{C\zeta - \mu} \delta_k, \quad (5.38)$$

for any $\zeta < \rho_k$.

We now define the diagonal part \mathcal{D}_{k+1} .

For any $j \in \mathbb{Z} \setminus \{0\}$ and any $\omega \in \mathcal{E}_k$ one has $|(\mathcal{P}_k^{(k)})_j^j(0)| \lesssim \|\mathcal{P}_k^{(k)}\|_{\rho_k}^{\mathcal{E}_k} \stackrel{(5.30)}{\leq} \delta_k$. The Hamiltonian structure guarantees that $\mathcal{P}_k^{(k)}(0)_j^j$ is purely imaginary and by the Kiszbraun Theorem there exists a Lipschitz extension $\omega \in \mathcal{O} \rightarrow iz_k(j)$ (with $z_k(j)$ real) of this function satisfying the bound $|z_k(j)|^\mathcal{O} \lesssim \delta_k$. Then, we define

$$\begin{aligned} \mathcal{D}_{k+1} &:= \text{diag}_{j \in \mathbb{Z} \setminus \{0\}} \Omega_{k+1}(j), \\ \Omega_{k+1}(j) &:= \Omega_k(j) + z_k(j) = \Omega_0(j) + r_{k+1}(j), \quad \forall j \in \mathbb{Z} \setminus \{0\}, \\ r_{k+1}(j) &:= r_k(j) + z_k(j) \end{aligned} \quad (5.39)$$

and one has

$$|r_{k+1}(j) - r_k(j)|^\mathcal{O} = |z_k(j)|^\mathcal{O} \leq \|\mathcal{P}_k^{(k)}\|_{\rho_k}^{\mathcal{E}_k} \stackrel{(5.30)}{\leq} \delta_k \sum_{j=1}^k 2^{-j} \quad (5.40)$$

which is the estimate (5.29) at the step $k+1$.

We now estimate the remainder \mathcal{P}_{k+1} in (5.34). Using (5.35) we see that

$$\mathcal{P}_{k+1} = \Pi_{N_k}^\perp \mathcal{P}_k^{(k)} + \sum_{p \geq 2} \frac{\text{Ad}_{\Psi_k}^{p-1}(\mathcal{Z}_k - \Pi_{N_k} \mathcal{P}_k^{(k)})}{p!} + \sum_{i=k+1}^m e^{-\Psi_k} \mathcal{P}_k^{(i)} e^{\Psi_k} + \sum_{p \geq 1} \frac{\text{Ad}_{\Psi_k}^p(\mathcal{P}_k^{(k)})}{p!}. \quad (5.41)$$

Denote

$$\begin{aligned} \mathcal{P}_{k+1} &= \sum_{i=k+1}^m \mathcal{P}_{k+1}^{(i)} \quad \text{where} \\ \mathcal{P}_{k+1}^{(k+1)} &:= \Pi_{N_k}^\perp \mathcal{P}_k^{(k)} + \sum_{p \geq 2} \frac{\text{Ad}_{\Psi_k}^{p-1}(\mathcal{Z}_k - \Pi_{N_k} \mathcal{P}_k^{(k)})}{p!} + e^{-\Psi_k} \mathcal{P}_k^{(k+1)} e^{\Psi_k} + \sum_{p \geq 1} \frac{\text{Ad}_{\Psi_k}^p(\mathcal{P}_k^{(k)})}{p!}, \\ \mathcal{P}_{k+1}^{(i)} &:= e^{-\Psi_k} \mathcal{P}_k^{(i)} e^{\Psi_k}, \quad i = k+2, \dots, m. \end{aligned} \quad (5.42)$$

ESTIMATE OF $\mathcal{P}_{k+1}^{(i)}$, $i = k + 2, \dots, m$. By the induction estimate, one has

$$\begin{aligned} \|e^{-\Psi_k} \mathcal{P}_k^{(i)} e^{\Psi_k}\|_{\rho_{k+1}}^{\mathcal{E}_{k+1}} &\leq \|\mathcal{P}_k^{(i)}\|_{\rho_i}^{\mathcal{E}_k} + \|\mathcal{P}_k^{(i)} - e^{-\Psi_k} \mathcal{P}_k^{(i)} e^{\Psi_k}\|_{\rho_i}^{\mathcal{E}_{k+1}} \\ &\lesssim \delta_i \sum_{j=1}^k 2^{-j} + \|\Psi_k\|_{\rho_i}^{\mathcal{E}_{k+1}} \|\mathcal{P}_k^{(i)}\|_{\rho_i}^{\mathcal{E}_{k+1}} \stackrel{(5.23)}{\lesssim} \delta_i \sum_{j=1}^{k+1} 2^{-j}. \end{aligned} \quad (5.43)$$

ESTIMATE OF $\mathcal{P}_{k+1}^{(k+1)}$. We estimate separately the four terms in the definition of $\mathcal{P}_{k+1}^{(k+1)}$ in (5.42). By Lemma A.9-(ii), one has

$$\|\Pi_{N_k}^\perp \mathcal{P}_k^{(k)}\|_{\rho_{k+1}}^{\mathcal{E}_k} \lesssim e^{-N_k \Delta_k} \|\mathcal{P}_k^{(k)}\|_{\rho_k}^{\mathcal{E}_k} \lesssim e^{-N_k \Delta_k} \delta_k. \quad (5.44)$$

By applying (A.7) and the estimate of Lemma A.9-(iii), one obtains

$$\begin{aligned} \left\| \sum_{p \geq 2} \frac{\text{Ad}_{\Psi_k}^{p-1}(\mathcal{Z}_k - \Pi_{N_k} \mathcal{P}_k^{(k)})}{p!} \right\|_{\rho_{k+1}}^{\mathcal{E}_{k+1}} &\leq \sum_{p \geq 2} \frac{C^{p-1}}{p!} (\|\Psi_k\|_{\rho_{k+1}}^{\mathcal{E}_{k+1}})^{p-1} \|\mathcal{P}_k^{(k)}\|_{\rho_k}^{\mathcal{E}_k} \\ &\lesssim \|\Psi_k\|_{\rho_{k+1}}^{\mathcal{E}_{k+1}} \|\mathcal{P}_k^{(k)}\|_{\rho_k}^{\mathcal{E}_k} \lesssim e^{C \Delta_k^{-\mu}} \delta_k^2 \end{aligned} \quad (5.45)$$

and similarly

$$\left\| \sum_{m \geq 1} \frac{\text{Ad}_{\Psi_k}^m(\mathcal{P}_k^{(k)})}{m!} \right\|_{\rho_{k+1}}^{\mathcal{E}_{k+1}} \lesssim e^{C \Delta_k^{-\mu}} \delta_k^2. \quad (5.46)$$

In conclusion we obtained

$$\|\mathcal{P}_{k+1}^{(k+1)}\|_{\rho_{k+1}}^{\mathcal{E}_{k+1}} \leq C' e^{-N_k \Delta_k} \delta_k + C' e^{C \Delta_k^{-\mu}} \delta_k^2 + \delta_{k+1} \sum_{j=1}^k 2^{-j} \quad (5.47)$$

where C' is an appropriate constant and the last summand is a bound for the term $e^{-\Psi_k} \mathcal{P}_k^{(k+1)} e^{\Psi_k}$, which can be obtained reasoning as in (5.43). Thus we obtain

$$\|\mathcal{P}_{k+1}^{(k+1)}\|_{\rho_{k+1}}^{\mathcal{E}_{k+1}} \leq \delta_{k+1} \sum_{j=1}^{k+1} 2^{-j} \quad (5.48)$$

provided

$$C' e^{-N_k \Delta_k} \delta_k + C' e^{C \Delta_k^{-\mu}} \delta_k^2 + \delta_{k+1} \sum_{j=1}^k 2^{-j} \leq \delta_{k+1} \sum_{j=1}^{k+1} 2^{-j},$$

which is of course follows from (5.23). ■

Now that we reduced \mathcal{L}_0 to the form $\mathcal{L}_m = \omega \cdot \partial_\varphi + \mathcal{D}_m + \mathcal{P}_m$ we can apply a ‘‘standard’’ KAM scheme to complete the diagonalization. This is a super-exponentially convergent iterative scheme based on iterating the following KAM step.

Lemma 5.4 (The $(m+1)$ -th step). *Following the notation of Lemma 5.3 we define*

$$\mathcal{E}_{m+1} := \left\{ \omega \in \mathcal{E}_m : |\omega \cdot \ell + \Omega_m(j) - \Omega_m(j')| \geq \frac{\gamma |j^3 - j'^3|}{\mathfrak{d}(\ell)}, \quad \forall (\ell, j, j') \neq (0, j, j), \quad |\ell|_\eta \leq N_m \right\}$$

and fix any ζ such that

$$e^{-N_m \zeta} \delta_m + e^{C\zeta^{-\mu}} \delta_m^2 \ll \delta_{m+1} \quad (5.49)$$

Then there exists a change of variables $\Phi_m := \exp(\Psi_m)$, such that

$$\|\Psi_m\|_{\rho_m - \zeta}^{\mathcal{E}_{m+1}} \lesssim e^{C\zeta^{-\mu}} \delta_m \quad (5.50)$$

which conjugates \mathcal{L}_m to the operator

$$\mathcal{L}_{m+1} = \omega \cdot \partial_\varphi + \mathcal{D}_{m+1} + \mathcal{P}_{m+1}.$$

The operator \mathcal{D}_{m+1} is of the form (5.28) and satisfies (5.29), with $k \rightsquigarrow m+1$, while the operator \mathcal{P}_{m+1} is such that

$$\|\mathcal{P}_{m+1}\|_{\rho_m - \zeta}^{\mathcal{E}_{m+1}} \leq \delta_{m+1}. \quad (5.51)$$

Proof. We reason similarly to Lemma 5.3 i.e. we fix Ψ_m in such a way that

$$\begin{aligned} \omega \cdot \partial_\varphi \Psi_m + [\mathcal{D}_m, \Psi_m] + \Pi_{N_m} \mathcal{P}_m &= \mathcal{Z}_m, \\ \mathcal{Z}_m &:= \text{diag}_{j \in \mathbb{Z} \setminus \{0\}} (\mathcal{P}_m)_j^j(0), \end{aligned} \quad (5.52)$$

so that we obtains

$$\|\Psi_m\|_{\rho_m - \zeta}^{\mathcal{E}_{m+1}} \lesssim e^{C\zeta^{-\mu}} \|\mathcal{P}_m\|_{\rho_m}^{\mathcal{E}_m} \lesssim e^{C\zeta^{-\mu}} \delta_m, \quad (5.53)$$

for any $\zeta < \rho_m$.

Now, for any $j \in \mathbb{Z} \setminus \{0\}$ and any $\omega \in \mathcal{E}_m$ one has $|(\mathcal{P}_m)_j^j(0)| \lesssim \|\mathcal{P}_m\|_{\rho_m}^{\mathcal{E}_m} \leq 2\delta_m$. The Hamiltonian structure guarantees that $\mathcal{P}_m(0)_j^j$ is purely imaginary and by the Kiszbraun Theorem there exists a Lipschitz extension $\omega \in \mathcal{O} \rightarrow iz_m(j)$ (with $z_m(j)$ real) of this function satisfying the bound $|z_m(j)|^{\mathcal{O}} \lesssim \delta_m$. Then, we define

$$\begin{aligned} \mathcal{D}_{m+1} &:= \text{diag}_{j \in \mathbb{Z} \setminus \{0\}} \Omega_{m+1}(j), \\ \Omega_{m+1}(j) &:= \Omega_m(j) + z_m(j) = \Omega_0(j) + r_{m+1}(j), \quad \forall j \in \mathbb{Z} \setminus \{0\}, \\ r_{m+1}(j) &:= r_m(j) + z_m(j) \end{aligned} \quad (5.54)$$

and (5.29), with $k \rightsquigarrow m+1$.

In order to obtain the bound 5.51 we start by recalling that

$$\mathcal{P}_{m+1} := \Pi_{N_m}^\perp \mathcal{P}_m + \sum_{p \geq 2} \frac{\text{Ad}_{\Psi_m}^{p-1} (\mathcal{Z}_m - \Pi_{N_m} \mathcal{P}_m)}{p!} + \sum_{p \geq 1} \frac{\text{Ad}_{\Psi_m}^p (\mathcal{P}_m)}{p!}, \quad (5.55)$$

so that reasoning as in (5.47) we obtain

$$\|\mathcal{P}_{m+1}\|_{\rho_m - \zeta}^{\mathcal{E}_{m+1}} \leq C' e^{-N_m \zeta} \delta_m + C' e^{C\zeta^{-\mu}} \delta_m^2 \quad (5.56)$$

and by (5.49) the assertion follows. \blacksquare

We now iterate the step of Lemma 5.4, using at each step a smaller loss of analyticity, namely at the p -th step we take ζ_p with

$$\sum_{p \geq m+1} \zeta_p = \zeta,$$

so that we obtain the following standard reducibility result; for a complete proof see [21].

Proposition 5.5. *For any $j \in \mathbb{Z} \setminus \{0\}$, the sequence $\Omega_k(j) = \Omega_0(j) + r_k(j)$, $k \geq 1$ provided in Lemmata 5.3, 5.4, and defined for any $\omega \in \mathcal{O}$ converges to $\Omega_\infty(j) = \Omega_0(j) + r_\infty(j)$ with $|r_\infty(j) - r_k(j)|^{\mathcal{O}} \lesssim \delta_k$. Defining the Cantor set*

$$\mathcal{E}_\infty := \left\{ \omega \in \mathcal{O} : |\omega \cdot \ell + \Omega_\infty(j) - \Omega_\infty(j')| \geq \frac{2\gamma|j^3 - j'^3|}{\mathfrak{d}(\ell)}, \quad \forall (\ell, j, j') \neq (0, j, j) \right\} \quad (5.57)$$

and

$$\mathcal{L}_\infty := \omega \cdot \partial_\varphi + \mathcal{D}_\infty, \quad \mathcal{D}_\infty := i \operatorname{diag}_{j \in \mathbb{Z} \setminus \{0\}} \Omega_\infty(j), \quad (5.58)$$

one has $\mathcal{E}_\infty \subseteq \bigcap_{k \geq 0} \mathcal{E}_k$.

Defining also

$$\tilde{\Phi}_k := \Phi_0 \circ \dots \circ \Phi_k \quad \text{with inverse} \quad \tilde{\Phi}_k^{-1} = \Phi_k^{-1} \circ \dots \circ \Phi_0^{-1}, \quad (5.59)$$

the sequence $\tilde{\Phi}_k$ converges for any $\omega \in \mathcal{E}_\infty$ to a symplectic, invertible map Φ_∞ w.r.t. the norm $\|\cdot\|_{\rho_m - 2\zeta}^{\mathcal{E}_\infty}$ and $\|\Phi_\infty^{\pm 1} - \operatorname{Id}\|_{\rho_m - 2\zeta}^{\mathcal{E}_\infty} \lesssim \delta_0$. Moreover for any $\omega \in \mathcal{E}_\infty$, one has that $\Phi_\infty^{-1} \mathcal{L}_0 \Phi_\infty = \mathcal{L}_\infty$.

5.3 Variations

We now consider an operator

$$\begin{aligned} \mathcal{L}_0^+ &\equiv \mathcal{L}_0(\lambda_1^+, \lambda_3^+, \mathcal{P}_0^+) = \omega \cdot \partial_\varphi + \mathcal{D}_0^+ + \mathcal{P}_0^+, \\ \mathcal{D}_0^+ &:= \lambda_3^+ \partial_x^3 + \lambda_1^+ \partial_x = i \operatorname{diag}_{j \in \mathbb{Z} \setminus \{0\}} \Omega_0^+(j), \\ \Omega_0^+(j) &:= -\lambda_3^+ j^3 + \lambda_1^+ j, \quad j \in \mathbb{Z} \setminus \{0\}. \end{aligned} \quad (5.60)$$

such that

$$|\lambda_1^+ - \lambda_1|^{\mathcal{O}^+}, |\lambda_3^+ - \lambda_3|^{\mathcal{O}^+}, \|\mathcal{P}_0^+ - \mathcal{P}_0\|_{\rho_{m+1}}^{\mathcal{O}^+} \leq \delta_{m+1} \quad (5.61)$$

where \mathcal{L} , λ_1 , λ_3 , \mathcal{P}_0 are given in (5.21) and $\mathcal{O}^+ \subseteq \mathcal{O}$. In other words, \mathcal{L}_0^+ is a small variation of \mathcal{L}_0 in (5.20) with also $m \rightsquigarrow m+1$.

Of course we can apply Proposition 5.5 to \mathcal{L}_0^+ ; our aim is to compare the ‘‘final frequencies’’ of \mathcal{L}_∞^+ with those of \mathcal{L}_∞ .

To this aim, we first apply Lemma 5.3 with $\mathcal{L}_0 \rightsquigarrow \mathcal{L}_0^+$ and $\gamma \rightsquigarrow \gamma_+ < \gamma$. In this way we obtain a sequence of sets $\mathcal{E}_k^+ \subseteq \mathcal{E}_{k-1}^+$ and a sequence of symplectic maps Φ_k^+ defined for $\omega \in \mathcal{E}_{k+1}^+$ such that setting \mathcal{L}_0^+ as in (5.60) and

$$\mathcal{L}_k := \Phi_{k-1}^{-1} \mathcal{L}_{k-1} \Phi_{k-1}, \quad (5.62)$$

one has

$$\mathcal{L}_k^+ := \omega \cdot \partial_\varphi + \mathcal{D}_k^+ + \mathcal{P}_k^+, \quad k \leq m+1, \quad (5.63)$$

where

$$\mathcal{D}_k^+ := \text{diag}_{j \in \mathbb{Z} \setminus \{0\}} \Omega_k^+(j), \quad \Omega_k^+(j) = \Omega_0^+(j) + r_k^+(j) \quad (5.64)$$

The sets \mathcal{E}_k^+ are defined as $\mathcal{E}_0^+ := \mathcal{O}^+$ and for $k \geq 1$

$$\mathcal{E}_k^+ := \left\{ \omega \in \mathcal{E}_{k-1}^+ : |\omega \cdot \ell + \Omega_{k-1}^+(j) - \Omega_{k-1}^+(j')| \geq \frac{\gamma_+ |j^3 - j'^3|}{\mathbf{d}(\ell)}, \right. \\ \left. \forall (\ell, j, j') \neq (0, j, j), \quad |\ell_\eta| \leq N_{k-1} \right\}. \quad (5.65)$$

Moreover one has $\Phi_{k-1}^+ := \exp(\Psi_{k-1}^+)$, with

$$\|\Psi_{k-1}^+\|_{\rho_k^+}^{\mathcal{E}_k^+} \lesssim e^{C\Delta_{k-1}^{-\mu}} \delta_{k-1}. \quad (5.66)$$

The following lemma holds.

Lemma 5.6. *For all $k = 1, \dots, m+1$ one has*

$$\|\mathcal{P}_k^+ - \mathcal{P}_k\|_{\rho_k}^{\mathcal{E}_k \cap \mathcal{E}_k^+} \leq \delta_{m+1}, \quad (5.67a)$$

$$|r_k^+(j) - r_k(j)|_{\mathcal{O} \cap \mathcal{O}^+} \leq \delta_{m+1} \quad (5.67b)$$

and

$$\|\Psi_{k-1}^+ - \Psi_{k-1}\|_{\rho_k}^{\mathcal{E}_k \cap \mathcal{E}_k^+} \lesssim \delta_{m+1}, \quad (5.68)$$

Proof. We procede differently for $k = 1, \dots, m$ and $k = m+1$.

For the first case we argue by induction. Assume the statement to hold up to some $k < m$. We want to prove

$$\|\Psi_k^+ - \Psi_k\|_{\rho_{k+1}}^{\mathcal{E}_{k+1} \cap \mathcal{E}_{k+1}^+} \leq \delta_{m+1}. \quad (5.69)$$

By Lemma 5.3, one has for $\omega \in \mathcal{E}_{k+1}^+$

$$(\Psi_k^+)_j^{j'}(\ell) := \begin{cases} \frac{((\mathcal{P}_k^+)^{(k)})_j^{j'}(\ell)}{\mathbf{i}(\omega \cdot \ell + \Omega_k^+(j) - \Omega_k^+(j'))}, & \forall (\ell, j, j') \neq (0, j, j), \quad |\ell|_\eta \leq N_k, \\ 0 & \text{otherwise,} \end{cases} \quad (5.70)$$

and direct calculation shows that for $\omega \in \mathcal{E}_{k+1} \cap \mathcal{E}_{k+1}^+$, one has

$$|(\Omega_k^+(j) - \Omega_k^+(j')) - (\Omega_k(j) - \Omega_k(j'))| \leq \delta_{m+1} |j^3 - j'^3| \quad (5.71)$$

and hence

$$|(\Psi_k^+)_j^{j'}(\ell) - (\Psi_k)_j^{j'}(\ell)|_{\mathcal{E}_{k+1} \cap \mathcal{E}_{k+1}^+} \lesssim \delta_{m+1} \mathbf{d}(\ell)^3 |(\mathcal{P}_k^+)_j^{j'}(\ell)|_{\mathcal{E}_{k+1} \cap \mathcal{E}_{k+1}^+} \\ + \mathbf{d}(\ell)^2 |(\mathcal{P}_k^+)_j^{j'}(\ell) - ((\mathcal{P}_k^+)^{(k)})_j^{j'}(\ell)|_{\mathcal{E}_{k+1} \cap \mathcal{E}_{k+1}^+}. \quad (5.72)$$

Therefore, reasoning as in (5.37)–(5.38), one uses Lemma A.6, the smallness condition (5.23) and the induction estimate (5.67a) so that (5.69) follows.

Now, from the definition of r_{k+1} in (5.39) it follows

$$|r_{k+1}^+(j) - r_{k+1}(j)|^{\mathcal{E}_{k+1} \cap \mathcal{E}_{k+1}^+} \leq \delta_{m+1}, \quad (5.73)$$

and by Kiszbraun Theorem applied to $r_{k+1}^+(j) - r_{k+1}(j)$, (5.67b) holds.

The estimate of $\mathcal{P}_{k+1}^+ - \mathcal{P}_{k+1}$ follows by explicit computation the difference by using the expressions provided in (5.41), using the induction estimates (5.30), (5.67a), the estimate (5.69) and by applying Lemma A.12.

For $k = m + 1$ the proof can be repeated word by word, the only difference being that Ψ_m is defined in (5.52) while Ψ_m^+ is defined in (5.36) with $k = m$. \blacksquare

5.4 Conclusion of the proof

To conclude the proof of Proposition 3.8 we start by noting that, setting \mathcal{O} appearing in (5.2) as $\mathcal{O}^{(n)}$ appearing in (3.11), the operator L_{n+1} appearing in (3.18) with of course $n \rightsquigarrow n + 1$ is of the form (5.1) with

$$\begin{aligned} \lambda_3 &= 1 + A_{n+1}, \\ a_1^{(k)}(\varphi, x) &= B_{k+1}(\varphi, x) - B_k(\varphi, x), \\ a_0^{(k)}(\varphi, x) &= C_{k+1}(\varphi, x) - C_k(\varphi, x). \end{aligned}$$

Moreover from (3.20) we have

$$\delta_k = \sigma_k^{-\tau_2} e^{\mathcal{C}\sigma_k^{-\mu}} \varepsilon_k, \quad \rho_k = s_k - 3\sigma_k$$

where s_k , σ_k and ε_k are defined in (3.6), so that L_{n+1} satisfies (5.2) with $m = n$. Thus, fixing

$$\zeta_k = \sigma_k, \quad 2\zeta = \sigma_k,$$

the smallness conditions (5.3) follows by definition. Hence we can apply Lemma 5.1 to L_{n+1} obtaining an operator of the form (5.5). In particular the conjugating operator \mathcal{M} satisfies

$$\|\mathcal{M} - \text{Id}\|_{s_n - 3\sigma_n}^{\mathcal{O}} \lesssim \sigma_0^{-\tau_2} e^{\mathcal{C}\sigma_0^{-\mu}} \varepsilon_0.$$

We are now in the setting of Section 5.2 with

$$\rho_k = s_k - 4\sigma_k, \quad \delta_k = \sigma_k^{-\tau_3} e^{2\mathcal{C}\sigma_k^{-\frac{1}{\eta}+}} \varepsilon_k$$

for some $\tau_3 > 0$. A direct calculation shows that the smallness conditions (5.22), (5.23), (5.24), (5.49) are satisfied provided we choose N_k appropriately, so that we can apply Proposition 5.5.

In conclusion we obtain an operator $M_{n+1} = \mathcal{M} \circ \Phi_\infty$ (recall that \mathcal{M} is constructed in Lemma 5.1) satisfying (3.49), (3.50), where $\Omega^{(n+1)}(j) := \Omega_\infty(j)$ and $\mathcal{E}^{(n+1)} = \mathcal{E}_\infty$. Note that in particular the functions $\Omega^{(n+1)}(j)$ turn out to be of the form (3.46).

Finally (3.47) follows from Lemmata 5.2 and 5.6 where \mathcal{L}_+ has the role of L_{n+1} while \mathcal{L} has the role of L_n . This means that here we are taking $m \rightsquigarrow n - 1$.

A Technical Lemmata

We start by recalling few results proved in [21]. Of course, as already noted in [21]-Remark 2.2, all the properties holding for $\mathcal{H}(\mathbb{T}_{\sigma+\rho}^\infty, \ell^\infty)$ hold verbatim for $\mathcal{H}(\mathbb{T}_{\sigma+\rho}^\infty \times \mathbb{T}_{\sigma+\rho}, \ell^\infty)$. In particular, all the estimates below hold also for the Lipschitz norms $|\cdot|_\sigma^\Omega$ and $\|\cdot\|_\sigma^\Omega$. Given two Banach spaces X, Y we denote by $\mathcal{B}(X, Y)$ the space of bounded linear operators from X to Y .

Proposition A.1 (Torus diffeomorphism). *Let $\alpha \in \mathcal{H}(\mathbb{T}_{\sigma+\rho}^\infty, \ell^\infty)$ be real on real. Then there exists a constant $\delta \in (0, 1)$ such that if $\rho^{-1}|\alpha|_{\sigma+\rho} \leq \delta$, then the map $\varphi \mapsto \varphi + \alpha(\varphi)$ is an invertible diffeomorphism of \mathbb{T}_σ^∞ (w.r.t. the ℓ^∞ -topology) and its inverse is of the form $\vartheta \mapsto \vartheta + \tilde{\alpha}(\vartheta)$, where $\tilde{\alpha} \in \mathcal{H}(\mathbb{T}_{\sigma+\frac{\rho}{2}}^\infty, \ell^\infty)$ is real on real and satisfies the estimate $|\tilde{\alpha}|_{\sigma+\frac{\rho}{2}} \lesssim |\alpha|_{\sigma+\rho}$.*

Corollary A.2. *Given $\alpha \in \mathcal{H}(\mathbb{T}_{\sigma+\rho}^\infty, \ell^\infty)$ as in Proposition A.1, the operators*

$$\begin{aligned} \Phi_\alpha &: \mathcal{H}(\mathbb{T}_{\sigma+\rho}^\infty, X) \rightarrow \mathcal{H}(\mathbb{T}_\sigma^\infty, X), & u(\varphi) &\mapsto u(\varphi + \alpha(\varphi)), \\ \Phi_{\tilde{\alpha}} &: \mathcal{H}(\mathbb{T}_{\sigma+\frac{\rho}{2}}^\infty, X) \rightarrow \mathcal{H}(\mathbb{T}_\sigma^\infty, X), & u(\vartheta) &\mapsto u(\vartheta + \tilde{\alpha}(\vartheta)) \end{aligned} \quad (\text{A.1})$$

are bounded, satisfy

$$\|\Phi_\alpha\|_{\mathcal{B}(\mathcal{H}(\mathbb{T}_{\sigma+\rho}^\infty, X), \mathcal{H}(\mathbb{T}_\sigma^\infty, X))}, \|\Phi_{\tilde{\alpha}}\|_{\mathcal{B}(\mathcal{H}(\mathbb{T}_{\sigma+\frac{\rho}{2}}^\infty, X), \mathcal{H}(\mathbb{T}_\sigma^\infty, X))} \leq 1,$$

and for any $\varphi \in \mathbb{T}_\sigma^\infty$, $u \in \mathcal{H}(\mathbb{T}_{\sigma+\rho}^\infty, X)$, $v \in \mathcal{H}(\mathbb{T}_{\sigma+\frac{\rho}{2}}^\infty, X)$ one has

$$\Phi_{\tilde{\alpha}} \circ \Phi_\alpha u(\varphi) = u(\varphi), \quad \Phi_\alpha \circ \Phi_{\tilde{\alpha}} v(\varphi) = v(\varphi).$$

Moreover Φ is close to the identity in the sense that

$$\|\Phi_\alpha(u) - u\|_\sigma \lesssim \rho^{-1}|\alpha|_\sigma |u|_{\sigma+\rho}. \quad (\text{A.2})$$

Given a function $u \in \mathcal{H}(\mathbb{T}_\sigma^\infty, X)$, we define its average on the infinite dimensional torus as

$$\int_{\mathbb{T}^\infty} u(\varphi) d\varphi := \lim_{N \rightarrow +\infty} \frac{1}{(2\pi)^N} \int_{\mathbb{T}^N} u(\varphi) d\varphi_1 \dots d\varphi_N. \quad (\text{A.3})$$

By Lemma 2.6 in [21], this definition is well posed and

$$\int_{\mathbb{T}^\infty} u(\varphi) d\varphi = u(0)$$

where $u(0)$ is the zero-th Fourier coefficient of u .

Lemma A.3 (Algebra). *One has $|uv|_\sigma \leq |u|_\sigma |v|_\sigma$ for $u, v \in \mathcal{H}(\mathbb{T}_\sigma^\infty \times \mathbb{T}_\sigma)$.*

Lemma A.4 (Cauchy estimates). *Let $u \in \mathcal{H}(\mathbb{T}_{\sigma+\rho}^\infty \times \mathbb{T}_{\sigma+\rho})$. Then $|\partial^k u|_\sigma \lesssim_k \rho^{-k} |u|_{\sigma+\rho}$.*

Lemma A.5 (Moser composition lemma). *Let $f : B_R(0) \rightarrow \mathbb{C}$ be an holomorphic function defined in a neighbourhood of the origin $B_R(0)$ of the complex plane \mathbb{C} . Then the composition operator $F(u) := f \circ u$ is a well defined non linear map $\mathcal{H}(\mathbb{T}_\sigma^\infty \times \mathbb{T}_\sigma) \rightarrow \mathcal{H}(\mathbb{T}_\sigma^\infty \times \mathbb{T}_\sigma)$ and if $|u|_\sigma \leq r < R$, one has the estimate $|F(u)|_\sigma \lesssim 1 + |u|_\sigma$. If f has a zero of order k at 0, then for any $|u|_\sigma \leq r < R$, one gets the estimate $|F(u)|_\sigma \lesssim |u|_\sigma^k$.*

For any function $u \in \mathcal{H}(\mathbb{T}_\sigma^\infty, X)$, given $N > 0$, we define the projector $\Pi_N u$ as

$$\Pi_N u(\varphi) := \sum_{|\ell|_\eta \leq N} u(\ell) e^{i\ell \cdot \varphi} \quad \text{and} \quad \Pi_N^\perp u := u - \Pi_N u.$$

Lemma A.6. (i) *Let $\rho > 0$. Then*

$$\sup_{\substack{\ell \in \mathbb{Z}_*^\infty \\ |\ell|_\eta < \infty}} \prod_i (1 + \langle i \rangle^5 |\ell_i|^5) e^{-\rho |\ell|_\eta} \leq e^{\tau \ln \left(\frac{\tau}{\rho} \right) \rho^{-\frac{1}{\eta}}}$$

for some constant $\tau = \tau(\eta) > 0$.

(ii) *Let $\rho > 0$. Then*

$$\sum_{\ell \in \mathbb{Z}_*^\infty} e^{-\rho |\ell|_\eta} \lesssim e^{\tau \ln \left(\frac{\tau}{\rho} \right) \rho^{-\frac{1}{\eta}}},$$

for some constant $\tau = \tau(\eta) > 0$.

(iii) *Let $\alpha > 0$. For $N \gg 1$ one has*

$$\sup_{\ell \in \mathbb{Z}_*^\infty: |\ell|_\alpha < N} \prod_i (1 + \langle i \rangle^5 |\ell_i|^5) \leq (1 + N)^{C(\alpha) N^{\frac{1}{1+\alpha}}} \quad (\text{A.4})$$

for some constant $C(\alpha) > 0$ such that $C(\alpha) \rightarrow \infty$ as $\alpha \rightarrow 0$.

Lemma A.7. *Given $u \in \mathcal{H}(\mathbb{T}_\sigma^\infty, X)$ for X some Banach space, let g be a pointwise absolutely convergent Formal Fourier series such that*

$$|g(\ell)|_X \leq \prod_i (1 + \langle i \rangle^5 |\ell_i|^5)^{\tau'} |u|_X,$$

for some $\tau' > 0$. Then for any $0 < \rho < \sigma$, then $g \in \mathcal{H}(\mathbb{T}_{\sigma-\rho}^\infty, X)$ and satisfies

$$|g|_{\sigma-\rho} \leq e^{\tau \ln \left(\frac{\tau}{\rho} \right) \rho^{-\frac{1}{\eta}}} |u|_\sigma$$

Proof. Follows directly from Lemma A.6 and Definition 2.1. ■

Lemma A.8. *Recalling (3.8) and the definition of $|\ell|_1$ in (1.5), one has*

$$\sum_{\ell \in \mathbb{Z}_*^\infty} \frac{|\ell|_1^3}{\mathbf{d}(\ell)} < \infty. \quad (\text{A.5})$$

Proof. First of all note that for all $\ell \in \mathbb{Z}_\infty^*$ one has

$$|\ell|_1^3 \leq \prod_i (1 + \langle i \rangle |\ell_i|)^3,$$

which implies

$$\frac{|\ell|_1^3}{\mathfrak{d}(\ell)} \lesssim \frac{1}{\prod_i (1 + \langle i \rangle^2 |\ell_i|^2)}. \quad (\text{A.6})$$

Then we recall that (see [9])

$$\sum_{\ell \in \mathbb{Z}_\infty^*} \frac{1}{\prod_i (1 + \langle i \rangle^2 |\ell_i|^2)} < \infty$$

which implies (A.5). ■

Lemma A.9. *Let $N, \sigma, \rho > 0$, $m, m' \in \mathbb{R}$, $\mathcal{R} \in \mathcal{H}(\mathbb{T}_\sigma^\infty, \mathcal{B}^{\sigma, m})$, $\mathcal{Q} \in \mathcal{H}(\mathbb{T}_{\sigma+\rho}^\infty, \mathcal{B}^{\sigma+\rho, m'})$.*

(i) The product operator $\mathcal{R}\mathcal{Q} \in \mathcal{H}(\mathbb{T}_\sigma^\infty, \mathcal{B}^{\sigma, m+m'})$ with $\|\mathcal{R}\mathcal{Q}\|_{\sigma, m+m'} \lesssim_m \rho^{-|m|} \|\mathcal{R}\|_{\sigma, m} \|\mathcal{Q}\|_{\sigma+\rho, m'}$. If $\mathcal{R}(\omega), \mathcal{Q}(\omega)$ depend on a parameter $\omega \in \Omega \subseteq \mathbb{D}_\gamma$, then $\|\mathcal{R}\mathcal{Q}\|_{\sigma, m+m'}^\Omega \lesssim_m \rho^{-(|m|+2)} \|\mathcal{R}\|_{\sigma, m}^\Omega \|\mathcal{Q}\|_{\sigma+\rho, m'}^\Omega$. If $m = m' = 0$, one has $\|\mathcal{R}\mathcal{Q}\|_\sigma^\Omega \lesssim \|\mathcal{R}\|_\sigma^\Omega \|\mathcal{Q}\|_\sigma^\Omega$.

(ii) The projected operator $\|\Pi_N^\perp \mathcal{R}\|_{\sigma, m} \leq e^{-\rho N} \|\mathcal{R}\|_{\sigma+\rho, m}$.

Given two linear operators \mathcal{A}, \mathcal{B} , we define for any $n \geq 0$, the operator $\text{Ad}_\mathcal{A}^n(\mathcal{B})$ as

$$\text{Ad}_\mathcal{A}^0(\mathcal{B}) := \mathcal{B}, \quad \text{Ad}_\mathcal{A}^{n+1}(\mathcal{B}) := [\text{Ad}_\mathcal{A}^n(\mathcal{B}), \mathcal{A}],$$

where

$$[\mathcal{B}, \mathcal{A}] := \mathcal{B}\mathcal{A} - \mathcal{A}\mathcal{B}.$$

By iterating the estimate (i) of Lemma A.9, one has that for any $n \geq 1$

$$\|\text{Ad}_\mathcal{A}^n(\mathcal{B})\|_\sigma \leq C^n \|\mathcal{A}\|_\sigma^n \|\mathcal{B}\|_\sigma \quad (\text{A.7})$$

for some constant $C > 0$.

Lemma A.10. *Let $0 < \dots < \rho_n < \dots < \rho_0$ and $0 < \dots \ll \delta_n \ll \dots \ll \delta_0$. Assume that $\sum_{i \geq 0} \delta_i < \infty$, choose any $n \geq 0$ and let \mathcal{A} and \mathcal{B} be linear operators such that*

$$\mathcal{A} = \sum_{i=0}^n \mathcal{A}_i \quad \mathcal{B} = \sum_{i=0}^n \mathcal{B}_i \quad \|\mathcal{A}_i\|_{\rho_i, -1}, \|\mathcal{B}_i\|_{\rho_i, 1} \leq \delta_i, \quad i = 0, \dots, n.$$

Then for any $0 < \zeta_i < \rho_i$ the following holds.

(i) For any $k \geq 1$, one has

$$\text{Ad}_\mathcal{A}^k(\mathcal{B}) = \sum_{i=0}^n \mathcal{R}_i^{(k)} \quad \text{with} \quad \|\mathcal{R}_i^{(k)}\|_{\rho_i - \zeta_i} \leq C_0^k \zeta_i^{-1} \delta_i \quad \forall i = 0, \dots, n$$

(ii) Let $\mathcal{R} := e^{-A}\mathcal{B}e^A - \mathcal{B}$. Then

$$\mathcal{R} = \sum_{i=0}^n \mathcal{R}_i \quad \text{with} \quad \|\mathcal{R}_i\|_{\rho_i - \zeta_i} \lesssim \zeta_i^{-1} \delta_i \quad \forall i = 0, \dots, n$$

Proof. PROOF OF ITEM (i). We prove the statement by induction on k . For $k = 1$, one has that

$$[\mathcal{B}, \mathcal{A}] = \sum_{i=0}^n \mathcal{R}_i^{(1)}, \quad \mathcal{R}_i^{(1)} := [\mathcal{B}_i, \mathcal{A}_i] + \sum_{j=0}^{i-1} ([\mathcal{B}_i, \mathcal{A}_j] - [\mathcal{A}_i, \mathcal{B}_j]).$$

Since for $j < i$ one has that $\rho_j > \rho_i$ and so all the terms in the above sum are analytic at least in the strip of width ρ_i . By applying Lemma A.9-(i) one has for any $0 < \zeta_i < \rho_i$

$$\|\mathcal{R}_i^{(1)}\|_{\rho_i - \zeta_i} \lesssim \zeta_i^{-1} \left(\delta_i^2 + \sum_{j=0}^i \delta_i \delta_j \right) \lesssim \zeta_i^{-1} \delta_i \sum_{j \geq 0} \delta_j \lesssim \zeta_i^{-1} \delta_i$$

for $i = 0, \dots, n$. Now we argue by induction. Assume that for some $k \geq 1$, $\mathcal{R}^{(k)} := \text{Ad}_{\mathcal{A}}^k(\mathcal{B}) = \sum_{i=0}^n \mathcal{R}_i^{(k)}$, with

$$\|\mathcal{R}_i^{(k)}\|_{\rho_i - \zeta_i} \leq C_0^k \zeta_i^{-1} \delta_i, \quad i = 0, \dots, n$$

for any $0 < \zeta_i < \rho_i$. Of course this implies that for all $j < i$ one has

$$\|\mathcal{R}_j^{(k)}\|_{\rho_i - \zeta_i} \leq C_0^k \zeta_i^{-1} \delta_j, \quad i = 0, \dots, n.$$

By definition

$$\begin{aligned} \text{Ad}_{\mathcal{A}}^{k+1}(\mathcal{B}) &= [\mathcal{R}^{(k)}, \mathcal{A}] = \sum_{i=0}^n \mathcal{R}_i^{(k+1)}, \\ \mathcal{R}_i^{(k+1)} &:= [\mathcal{R}_i^{(k)}, \mathcal{A}_i] + \sum_{j=0}^{i-1} ([\mathcal{R}_i^{(k)}, \mathcal{A}_j] - [\mathcal{A}_i, \mathcal{R}_j^{(k)}]), \quad \forall i = 0, \dots, n. \end{aligned}$$

Hence by applying Lemma A.9-(i) and using the induction hypothesis, one obtains

$$\begin{aligned} \|\mathcal{R}_i^{(k+1)}\|_{\rho_i - \zeta_i} &\leq C \left(\|\mathcal{R}_i^{(k)}\|_{\rho_i - \zeta_i} \|\mathcal{A}_i\|_{\rho_i - \zeta_i} + \sum_{j=0}^{i-1} \|\mathcal{R}_i^{(k)}\|_{\rho_i - \zeta_i} \|\mathcal{A}_j\|_{\rho_i - \zeta_i} + \|\mathcal{R}_j^{(k)}\|_{\rho_i - \zeta_i} \|\mathcal{A}_i\|_{\rho_i - \zeta_i} \right) \\ &\leq C \zeta_i^{-1} C_0^k \delta_i \sum_{j=0}^{i-1} \delta_j \leq C C_0^k \zeta_i^{-1} \delta_i \sum_{j \geq 0} \delta_j \leq C_0^{k+1} \zeta_i^{-1} \delta_i. \end{aligned}$$

PROOF OF (ii). One has

$$\mathcal{R} = e^{-A}\mathcal{B}e^A - \mathcal{B} = \sum_{k \geq 1} \frac{\text{Ad}_{\mathcal{A}}^k(\mathcal{B})}{k!} \stackrel{(i)}{=} \sum_{i=0}^n \mathcal{R}_i \quad \text{where} \quad \mathcal{R}_i = \sum_{k \geq 1} \frac{\mathcal{R}_i^{(k)}}{k!},$$

so that

$$\|\mathcal{R}_i\|_{\rho_i-\zeta_i} \leq \sum_{k \geq 1} \frac{\|\mathcal{R}_i^{(k)}\|_{\rho_i-\zeta_i}}{k!} \leq \sum_{k \geq 1} \frac{C_0^k}{k!} \zeta_i^{-1} \delta_i \lesssim \zeta_i^{-1} \delta_i.$$

Therefore the assertion follows. \blacksquare

Lemma A.11. *Let $\{\rho_n\}_{n \geq 0}$ and $\{\delta_n\}_{n \geq 0}$ as in Lemma A.10. Choose any $n > 0$ and consider*

$$g(\varphi, x) = \sum_{i=0}^n g_i(\varphi, x), \quad \text{with } g_i \in \mathcal{H}_{\rho_i}, \quad |g_i|_{\rho_i} \leq \delta_i, \quad i = 0, \dots, n.$$

Then the following holds.

(i) Consider the commutator $[\partial_x^3, \mathcal{G}]$ where $\mathcal{G} := \pi_0^\perp g(\varphi, x) \partial_x^{-1}$. Then, one has

$$[\partial_x^3, \mathcal{G}] = 3g_x \partial_x + \mathcal{R}, \quad \mathcal{R} := \sum_{k=0}^n \mathcal{R}_i, \quad \text{where } \|\mathcal{R}_i\|_{\rho_i-\zeta_i} \lesssim \zeta_i^{-3} \delta_i, \quad \text{for } 0 < \zeta_i < \rho_i.$$

(ii) Let $\zeta_0, \zeta_1, \dots, \zeta_n$ satisfying $0 < 2\zeta_i < \rho_i$, $0 < \rho_n - \zeta_n < \rho_{n-1} - \zeta_{n-1} < \dots < \rho_0 - \zeta_0$ and assume that $\sum_{i \geq 0} \zeta_i^{-3} \delta_i < \infty$. Then, one has

$$e^{-\mathcal{G}} \partial_x^3 e^{\mathcal{G}} = \partial_x^3 + 3g_x \partial_x + \mathcal{R}, \quad \mathcal{R} = \sum_{i=0}^n \mathcal{R}_i, \quad \|\mathcal{R}_i\|_{\rho_i-2\zeta_i} \lesssim \zeta_i^{-4} \delta_i, \quad i = 0, \dots, n.$$

(iii) Let $\zeta_0, \zeta_1, \dots, \zeta_n$ satisfying $0 < \zeta_i < \rho_i$, $0 < \rho_n - \zeta_n < \rho_{n-1} - \zeta_{n-1} < \dots < \rho_0 - \zeta_0$ and assume that $\sum_{i \geq 0} \zeta_i^{-1} \delta_i < \infty$. Then

$$e^{-\mathcal{G}} (\omega \cdot \partial_\varphi) e^{\mathcal{G}} = \omega \cdot \partial_\varphi + \mathcal{R}, \quad \mathcal{R} = \sum_{i=0}^n \mathcal{R}_i, \quad \|\mathcal{R}_i\|_{\rho_i-\zeta_i} \lesssim \zeta_i^{-1} \delta_i, \quad i = 0, \dots, n.$$

Proof. PROOF OF (i). One has

$$\begin{aligned} [\partial_x^3, \pi_0^\perp g \partial_x^{-1}] &= \pi_0^\perp (3g_x \partial_x + 3g_{xx} + g_{xxx} \partial_x^{-1}) = 3g_x \partial_x + \mathcal{R}, \\ \mathcal{R} &:= \sum_{i=0}^n \mathcal{R}_i, \quad \mathcal{R}_i := \pi_0^\perp (3(g_i)_{xx} + (g_i)_{xxx} \partial_x^{-1}) - 3\pi_0(g_i)_x \partial_x. \end{aligned}$$

Therefore

$$\|\mathcal{R}_i\|_{\rho_i-\zeta_i} \lesssim \zeta_i^{-3} \delta_i.$$

PROOF OF (ii). In view of the item (i), it is enough to estimate

$$\sum_{k \geq 2} \frac{\text{Ad}_{\mathcal{G}}^k(\partial_x^3)}{k!}.$$

Let

$$\mathcal{B} := [\partial_x^3, \mathcal{G}] = 3g_x \partial_x + \mathcal{R} = \sum_{i=0}^n \mathcal{B}_i, \quad \mathcal{B}_i := 3(g_i)_x \partial_x + \mathcal{R}_i, \quad i = 0, \dots, n, \quad (\text{A.8})$$

$$\mathcal{G} = \sum_{i=0}^n \mathcal{G}_i, \quad \mathcal{G}_i := \pi_0^\perp g_i(\varphi, x) \partial_x^{-1} \quad i = 0, \dots, n.$$

One has

$$\begin{aligned} \|\mathcal{B}_i\|_{\rho_i - \zeta_i, 1} &\lesssim \zeta_i^{-3} \delta_i, \quad i = 0, \dots, n, \\ \|\mathcal{G}_i\|_{\rho_i - \zeta_i, -1} &\leq \|\mathcal{G}_i\|_{\rho_i, -1} \lesssim |f_i|_{\rho_i} \lesssim \delta_i \leq \zeta_i^{-3} \delta_i, \quad i = 0, \dots, n \end{aligned} \quad (\text{A.9})$$

For any $k \geq 2$ one has

$$\text{Ad}_{\mathcal{G}}^k(\partial_x^3) = \text{Ad}_{\mathcal{G}}^{k-1}([\partial_x^3, \mathcal{G}]) = \text{Ad}_{\mathcal{G}}^{k-1}(\mathcal{B}),$$

hence, we can apply Lemma A.10 (replacing ρ_i with $\rho_i - \zeta_i$ and δ_i with $\zeta_i^{-3} \delta_i$) obtaining

$$\text{Ad}_{\mathcal{G}}^k(\partial_x^3) = \sum_{i=0}^n \mathcal{R}_i^{(k)}$$

where $\mathcal{R}_i^{(k)}$ satisfies

$$\|\mathcal{R}_i^{(k)}\|_{\rho_i - 2\zeta_i} \leq C_0^k \zeta_i^{-4} \delta_i, \quad i = 0, \dots, n \quad (\text{A.10})$$

and hence by setting

$$\mathcal{R} = \sum_{k \geq 2} \frac{\text{Ad}_{\mathcal{G}}^k(\partial_x^3)}{k!} = \sum_{i=0}^n \mathcal{R}_i$$

item (ii) follows.

PROOF OF ITEM (iii). The proof can be done arguing as in the item (ii), using that

$$e^{-\mathcal{G}}(\omega \cdot \partial_\varphi) e^{\mathcal{G}} = \omega \cdot \partial_\varphi + \sum_{k \geq 1} \frac{\text{Ad}_{\mathcal{G}}^{k-1}(\omega \cdot \partial_\varphi \mathcal{G})}{k!}, \quad \text{where } (\omega \cdot \partial_\varphi \mathcal{G}) := \pi_0^\perp \omega \cdot \partial_\varphi g(\varphi, x) \partial_x^{-1}.$$

■

Lemma A.12. *Let $\mathcal{A}, \mathcal{A}_+, \mathcal{B}, \mathcal{B}_+$ be bounded operators w.r.t. a norm $\|\cdot\|_\sigma$, and define*

$$M_{\mathcal{A}} := \max\{\|\mathcal{A}_+\|_\sigma, \|\mathcal{A}\|_\sigma\}, \quad M_{\mathcal{B}} := \max\{\|\mathcal{B}_+\|_\sigma, \|\mathcal{B}\|_\sigma\}. \quad (\text{A.11})$$

Then the following holds.

(i) *For any $k \geq 0$, one has*

$$\|\text{Ad}_{\mathcal{A}_+}^k(\mathcal{B}_+) - \text{Ad}_{\mathcal{A}}^k(\mathcal{B})\|_\sigma \leq C_*^k M_{\mathcal{A}}^k M_{\mathcal{B}} (\|\mathcal{A}_+ - \mathcal{A}\|_\sigma + \|\mathcal{B}_+ - \mathcal{B}\|_\sigma)$$

for some constant $C_ > 0$.*

(ii)

$$\|e^{-\mathcal{A}_+} \mathcal{B}_+ e^{\mathcal{A}_+} - e^{-\mathcal{A}} \mathcal{B} e^{\mathcal{A}}\|_\sigma \lesssim \|\mathcal{A}_+ - \mathcal{A}\|_\sigma + \|\mathcal{B}_+ - \mathcal{B}\|_\sigma.$$

Proof. PROOF OF (i). We argue by induction. Of course the result is trivial for $k = 0$. Assume that the estimate holds for some $k \geq 1$. Then

$$\begin{aligned} \text{Ad}_{\mathcal{A}_+}^{k+1}(\mathcal{B}_+) - \text{Ad}_{\mathcal{A}}^{k+1}(\mathcal{B}) &= \text{Ad}_{\mathcal{A}_+} \left(\text{Ad}_{\mathcal{A}_+}^k(\mathcal{B}_+) \right) - \text{Ad}_{\mathcal{A}} \left(\text{Ad}_{\mathcal{A}}^k(\mathcal{B}) \right) \\ &= \text{Ad}_{\mathcal{A}_+} \left(\text{Ad}_{\mathcal{A}_+}^k(\mathcal{B}_+) - \text{Ad}_{\mathcal{A}}^k(\mathcal{B}) \right) - \text{Ad}_{\mathcal{A}_+ - \mathcal{A}} \left(\text{Ad}_{\mathcal{A}}^k(\mathcal{B}) \right). \end{aligned}$$

Hence, by the induction hypothesis, using (A.11), (A.7) and Lemma A.9-(i), one obtains that

$$\begin{aligned} \|\text{Ad}_{\mathcal{A}_+}^{k+1}(\mathcal{B}_+) - \text{Ad}_{\mathcal{A}}^{k+1}(\mathcal{B})\|_{\sigma} &\lesssim \|\mathcal{A}_+\|_{\sigma} \|\text{Ad}_{\mathcal{A}_+}^k(\mathcal{B}_+) - \text{Ad}_{\mathcal{A}}^k(\mathcal{B})\|_{\sigma} + \|\mathcal{A}_+ - \mathcal{A}\|_{\sigma} C^k \|\mathcal{A}\|_{\sigma}^k \|\mathcal{B}\|_{\sigma} \\ &\lesssim C_*^k M_{\mathcal{A}}^{k+1} M_{\mathcal{B}} (\|\mathcal{A}_+ - \mathcal{A}\|_{\sigma} + \|\mathcal{B}_+ - \mathcal{B}\|_{\sigma}) + C^k M_{\mathcal{A}}^k M_{\mathcal{B}} \|\mathcal{A}_+ - \mathcal{A}\|_{\sigma} \\ &\leq C_*^{k+1} M_{\mathcal{A}}^{k+1} M_{\mathcal{B}} (\|\mathcal{A}_+ - \mathcal{A}\|_{\sigma} + \|\mathcal{B}_+ - \mathcal{B}\|_{\sigma}) \end{aligned}$$

for some $C_* > 0$ large enough.

PROOF OF (ii). It follows by item (i), using that

$$e^{-\mathcal{A}_+} \mathcal{B}_+ e^{\mathcal{A}_+} - e^{-\mathcal{A}} \mathcal{B} e^{\mathcal{A}} = \sum_{k \geq 0} \frac{\text{Ad}_{\mathcal{A}_+}^k(\mathcal{B}_+) - \text{Ad}_{\mathcal{A}}^k(\mathcal{B})}{k!}.$$

■

Lemma A.13. *Let $\mathcal{A}, \mathcal{A}^+, \mathcal{B}, \mathcal{B}^+$ be linear operators satisfying*

$$\|\mathcal{A}\|_{\rho, -1}, \|\mathcal{A}^+\|_{\rho, -1}, \|\mathcal{B}\|_{\rho, 1}, \|\mathcal{B}^+\|_{\rho, 1} < C_0.$$

Then the following holds.

(i) *For any $k \geq 1$,*

$$\|\text{Ad}_{\mathcal{A}_+}^k(\mathcal{B}_+) - \text{Ad}_{\mathcal{A}}^k(\mathcal{B})\|_{\rho - \zeta} \leq C^k \zeta^{-1} (\|\mathcal{A}_+ - \mathcal{A}\|_{\rho, -1} + \|\mathcal{B}_+ - \mathcal{B}\|_{\rho, 1})$$

for some constant $C > 0$ depending on C_0 .

(ii) *Setting $\mathcal{R} := e^{-\mathcal{A}} \mathcal{B} e^{\mathcal{A}} - \mathcal{B}$, and $\mathcal{R}_+ := e^{-\mathcal{A}_+} \mathcal{B}_+ e^{\mathcal{A}_+} - \mathcal{B}_+$, one has*

$$\|\mathcal{R} - \mathcal{R}_+\|_{\rho - \zeta} \lesssim \zeta^{-1} (\|\mathcal{A} - \mathcal{A}_+\|_{\rho, -1} + \|\mathcal{B} - \mathcal{B}_+\|_{\rho, 1}).$$

Proof. PROOF OF (i). We first estimate $\text{Ad}_{\mathcal{A}_+}(\mathcal{B}_+) - \text{Ad}_{\mathcal{A}}(\mathcal{B})$. One has

$$\text{Ad}_{\mathcal{A}_+}(\mathcal{B}_+) - \text{Ad}_{\mathcal{A}}(\mathcal{B}) = \text{Ad}_{\mathcal{A}_+}(\mathcal{B}_+ - \mathcal{B}) + \text{Ad}_{\mathcal{A}_+ - \mathcal{A}}(\mathcal{B}).$$

By Lemma A.9-(i), one has

$$\|\text{Ad}_{\mathcal{A}}(\mathcal{B})\|_{\rho - \zeta}, \|\text{Ad}_{\mathcal{A}_+}(\mathcal{B}_+)\|_{\rho - \zeta} \lesssim \zeta^{-1}, \tag{A.12}$$

and

$$\|\text{Ad}_{\mathcal{A}_+}(\mathcal{B}_+) - \text{Ad}_{\mathcal{A}}(\mathcal{B})\|_{\rho - \zeta} \lesssim \zeta^{-1} (\|\mathcal{A} - \mathcal{A}_+\|_{\rho, -1} + \|\mathcal{B} - \mathcal{B}_+\|_{\rho, 1}). \tag{A.13}$$

In order to estimate $\text{Ad}_{\mathcal{A}_+}^k(\mathcal{B}_+) - \text{Ad}_{\mathcal{A}}^k(\mathcal{B}) = \text{Ad}_{\mathcal{A}_+}^{k-1}\text{Ad}_{\mathcal{A}_+}(\mathcal{B}_+) - \text{Ad}_{\mathcal{A}}^{k-1}\text{Ad}_{\mathcal{A}}(\mathcal{B})$ for any $k \geq 2$, we apply Lemma A.12-(i) where we replace \mathcal{B}_+ with $\text{Ad}_{\mathcal{A}_+}(\mathcal{B}_+)$ and \mathcal{B} with $\text{Ad}_{\mathcal{A}}(\mathcal{B})$, together with the estimates (A.12), (A.13).

PROOF OF (ii). It follows by (i) using that $\mathcal{R}_+ - \mathcal{R} = \sum_{k \geq 1} \frac{\text{Ad}_{\mathcal{A}_+}^k(\mathcal{B}_+) - \text{Ad}_{\mathcal{A}}^k(\mathcal{B})}{k!}$. \blacksquare

Lemma A.14. *Let $g_+, g \in \mathcal{H}_\rho$, $\mathcal{G} := \pi_0^\perp g(\varphi, x)\partial_x^{-1}$, $\mathcal{G}_+ := \pi_0^\perp g_+(\varphi, x)\partial_x^{-1}$. Then the following holds.*

(i) *The operators $\mathcal{R} := e^{-\mathcal{G}}\partial_x^3 e^{\mathcal{G}} - \partial_x^3 - 3g_x\partial_x$, $\mathcal{R}_+ := e^{-\mathcal{G}_+}\partial_x^3 e^{\mathcal{G}_+} - \partial_x^3 - 3(g_+)_x\partial_x$ satisfy $\|\mathcal{R}_+ - \mathcal{R}\|_{\rho-\zeta} \lesssim \zeta^{-\tau}|g_+ - g|_\rho$ for some constant $\tau > 0$.*

(ii) *The operators $\mathcal{R} := e^{-\mathcal{G}}\omega \cdot \partial_\varphi e^{\mathcal{G}} - \omega \cdot \partial_\varphi$ and $\mathcal{R}_+ := e^{-\mathcal{G}_+}\omega \cdot \partial_\varphi e^{\mathcal{G}_+} - \omega \cdot \partial_\varphi$ satisfy the estimate $\|\mathcal{R}_+ - \mathcal{R}\|_{\rho-\zeta} \lesssim \zeta^{-\tau}|g_+ - g|_\rho$, for some constant $\tau > 0$.*

Proof. We only prove the item (i). The item (ii) can be proved by similar arguments. We compute

$$\begin{aligned} \mathcal{B} &:= [\partial_x^3, \pi_0^\perp g \partial_x^{-1}] = \pi_0^\perp (3g_x \partial_x + 3g_{xx} + g_{xxx} \partial_x^{-1}) = 3g_x \partial_x + \mathcal{R}_B, \\ \mathcal{R}_B &:= \pi_0^\perp (3g_{xx} + g_{xxx} \partial_x^{-1}) - \pi_0 (3g_x \partial_x), \\ \mathcal{B}_+ &:= [\partial_x^3, \pi_0^\perp g_+ \partial_x^{-1}] = \pi_0^\perp (3(g_+)_x \partial_x + 3(g_+)_{xx} + (g_+)_{xxx} \partial_x^{-1}) = 3(g_+)_x \partial_x + \mathcal{R}_{B_+}, \\ \mathcal{R}_{B_+} &:= \pi_0^\perp (3(g_+)_{xx} + (g_+)_{xxx} \partial_x^{-1}) - \pi_0 (3(g_+)_x \partial_x). \end{aligned} \tag{A.14}$$

Hence

$$\begin{aligned} \mathcal{R}_+ - \mathcal{R} &= \mathcal{R}_{B_+} - \mathcal{R}_B + \sum_{k \geq 2} \frac{\text{Ad}_{\mathcal{G}_+}^k(\partial_x^3) - \text{Ad}_{\mathcal{G}}^k(\partial_x^3)}{k!} \\ &\stackrel{(A.14)}{=} \mathcal{R}_{B_+} - \mathcal{R}_B + \sum_{k \geq 2} \frac{\text{Ad}_{\mathcal{G}_+}^{k-1}(\mathcal{B}_+) - \text{Ad}_{\mathcal{G}}^{k-1}(\mathcal{B})}{k!}. \end{aligned} \tag{A.15}$$

By a direct calculation one can show the estimates

$$\begin{aligned} \|\mathcal{B}\|_{\rho-\zeta,1} &\lesssim \zeta^{-3}|g|_\rho, \quad \|\mathcal{B}_+\|_{\rho-\zeta,1} \lesssim \zeta^{-3}|g_+|_\rho, \\ \|\mathcal{G}\|_{\rho,-1} &\lesssim |g|_\rho, \quad \|\mathcal{G}_+\|_{\rho,-1} \lesssim |g_+|_\rho, \\ \|\mathcal{R}_{B_+} - \mathcal{R}_B\|_{\rho-\zeta} &\lesssim \zeta^{-3}|g_+ - g|_\rho, \quad \|\mathcal{G}_+ - \mathcal{G}\|_{\rho,-1} \lesssim |g_+ - g|_\rho. \end{aligned} \tag{A.16}$$

The latter estimates, together with Lemma A.13-(i) allow to deduce

$$\|\text{Ad}_{\mathcal{G}_+}^{k-1}(\mathcal{B}_+) - \text{Ad}_{\mathcal{G}}^{k-1}(\mathcal{B})\|_{\rho-\zeta} \leq C^k \zeta^{-\tau}, \quad \forall k \geq 2, \tag{A.17}$$

for some constant $\tau > 0$. Thus (A.15)-(A.17) imply the desired bound. \blacksquare

References

- [1] P. Baldi, M. Berti, R. Montalto. *KAM for quasi-linear and fully nonlinear forced KdV*. *Math. Ann.*, 359:471–536, 2014.

- [2] P. Baldi, M. Berti, E. Haus, R. Montalto, *Time quasi-periodic gravity water waves in finite depth*, *Inventiones Math.* 214 (2), 739–911, 2018.
- [3] L. Biasco, J.E. Massetti, M. Procesi. *An abstract Birkhoff Normal Form Theorem and exponential type stability of the 1d NLS*.
- [4] L. Biasco, J.E. Massetti, M. Procesi. *Almost periodic solutions for the 1d NLS*, 2019. Preprint.
- [5] J. Bourgain, *Construction of quasi-periodic solutions for Hamiltonian perturbations of linear equations and applications to nonlinear PDE*, *Internat. Math. Res. Notices*, 1994
- [6] J. Bourgain, *Quasi-periodic solutions of Hamiltonian perturbations of 2D linear Schrödinger equations*, *Ann. of Math. (2)*, **148**, (1998)
- [7] J. Bourgain, *Green’s function estimates for lattice Schrödinger operators and applications*, Princeton University Press, 2005
- [8] J. Bourgain, *Construction of approximative and almost periodic solutions of perturbed linear Schrödinger and wave equations*, *Geom. Funct. Anal.* 6, no.2 (1996), 201–230
- [9] J. Bourgain. *On invariant tori of full dimension for 1D periodic NLS*. *J. Funct. Anal.*, 229(1):62–94, 2005.
- [10] L. Chierchia, P. Perfetti *Second order Hamiltonian equations on $f\mathbb{T}^\infty$ and almost-periodic solutions*, *J. Diff. Equations*, 116, 1995
- [11] H. Cong, J. Liu, Y. Shi, X. Yuan, *The stability of full dimensional KAM tori for nonlinear Schrödinger equation*, *J. Diff. Equations*, **264**, no.7, 2018
- [12] L. Corsi, R. Montalto. *Quasi-periodic solutions for the forced Kirchhoff equation on \mathbb{T}^d* . *Nonlinearity*, 31(11):5075–5109, 2018.
- [13] L. Corsi, R. Feola, M. Procesi. *Finite dimensional invariant KAM tori for tame vector fields*. *Transactions of the AMS*, 372 (2019), no.3 1913–1983
- [14] Craig W., Wayne C. E., *Newton’s method and periodic solutions of nonlinear wave equation*, *Comm. Pure Appl. Math.* 46, 1409–1498, 1993.
- [15] R. Feola and M. Procesi. *KAM for quasi-linear autonomous NLS*, preprint 2017.
- [16] S. Kuksin, *Hamiltonian perturbations of infinite-dimensional linear systems with imaginary spectrum*, *Funktsional Anal. i Prilozhen.*, 21, 22–37, 95, 1987.
- [17] S. Kuksin *Perturbations of quasiperiodic solutions of infinite-dimensional Hamiltonian systems* (*Math. USSR Izvestiya* 32 (1989), 39–62)
- [18] S. Kuksin *The perturbation theory for the quasiperiodic solutions of infinite-dimensional Hamiltonian systems and its applications to the Korteweg de Vries equation* (*Math. USSR Sbornik* 64 (1989), 397–413)
- [19] S. Kuksin, J. Pöschel. *Invariant Cantor manifolds of quasi-periodic oscillations for a nonlinear Schrödinger equation*. *Ann. of Math. (2)*, 143(1):149–179, 1996.

- [20] S. Liu *The existence of almost-periodic solutions for 1-dimensional nonlinear Schrödinger equation with quasi-periodic forcing* J. Math. Phys. **61**, 031502 (2020)
- [21] R. Montalto, M. Procesi *Linear Schrödinger equation with an almost periodic potential*, Preprint arXiv:1910.12300, 2019
- [22] J. Pöschel, *A KAM-Theorem for some nonlinear PDEs*, Ann. Sc. Norm. Pisa, 23, 119-148, 1996.
- [23] J. Pöschel. *On the construction of almost periodic solutions for a nonlinear Schrödinger equation.* Ergodic Theory Dynam. Systems, 22(5):1537–1549, 2002.
- [24] P. H. Rabinowitz. *Periodic solutions of nonlinear hyperbolic partial differential equations*, Comm. Pure Appl. Math., 20:145–205, 1967.
- [25] P. H. Rabinowitz. *Periodic solutions of nonlinear hyperbolic partial differential equations. II* Comm. Pure Appl. Math., 22:15–39, 1968.
- [26] J. Rui, B. Liu, J. Zhang, *Almost periodic solutions for a class of linear Schrödinger equations with almost periodic forcing*, J. Math. Phys. **57**, 092702 (2016). 18
- [27] J. Rui, B. Liu, *Almost-periodic solutions of an almost-periodically forced wave equation*, J. Math. Anal. Appl. **451**, no.2 (2017), 629–658
- [28] C.E. Wayne, *Periodic and quasi-periodic solutions of nonlinear wave equations via KAM theory*, Comm. Math. Phys. 127, 479-528, 1990.
- [29] X. Xu, J. Geng, *Almost periodic solutions of one dimensional Schrödinger equation with the external parameters*, J. Dyn. Differ. Equations 25, 435–450 (2013).