

# KAM for autonomous quasi-linear perturbations of KdV

Pietro Baldi, Massimiliano Berti, Riccardo Montalto

**Abstract.** We prove the existence and stability of Cantor families of quasi-periodic, small amplitude solutions of *quasi-linear* (i.e. *strongly nonlinear*) autonomous Hamiltonian perturbations of KdV.

*Keywords:* KdV, KAM for PDEs, quasi-linear PDEs, Nash-Moser theory, quasi-periodic solutions.

*MSC 2010:* 37K55, 35Q53.

## Contents

<b>1</b>	<b>Introduction and main results</b>	<b>1</b>
<b>2</b>	<b>Preliminaries</b>	<b>6</b>
2.1	Hamiltonian formalism of KdV . . . . .	6
2.2	Functional setting . . . . .	7
<b>3</b>	<b>Weak Birkhoff normal form</b>	<b>9</b>
<b>4</b>	<b>Action-angle variables</b>	<b>14</b>
<b>5</b>	<b>The nonlinear functional setting</b>	<b>15</b>
<b>6</b>	<b>Approximate inverse</b>	<b>17</b>
<b>7</b>	<b>The linearized operator in the normal directions</b>	<b>23</b>
7.1	Composition with the map $G_\delta$ . . . . .	24
7.2	The linearized operator in the normal directions . . . . .	25
<b>8</b>	<b>Reduction of the linearized operator in the normal directions</b>	<b>28</b>
8.1	Change of the space variable . . . . .	28
8.2	Reparametrization of time . . . . .	32
8.3	Translation of the space variable . . . . .	34
8.4	Linear Birkhoff normal form. Step 1 . . . . .	35
8.5	Linear Birkhoff normal form. Step 2 . . . . .	38
8.6	Descent method . . . . .	40
8.7	KAM reducibility and inversion of $\mathcal{L}_\omega$ . . . . .	41
<b>9</b>	<b>The Nash-Moser nonlinear iteration</b>	<b>42</b>

## 1 Introduction and main results

In this paper we prove the existence and stability of Cantor families of quasi-periodic solutions of Hamiltonian *quasi-linear* (also called “*strongly nonlinear*”, e.g. in [24]) perturbations of the KdV equation

$$u_t + u_{xxx} - 6uu_x + \mathcal{N}_4(x, u, u_x, u_{xx}, u_{xxx}) = 0, \quad (1.1)$$

under periodic boundary conditions  $x \in \mathbb{T} := \mathbb{R}/2\pi\mathbb{Z}$ , where

$$\mathcal{N}_4(x, u, u_x, u_{xx}, u_{xxx}) := -\partial_x[(\partial_u f)(x, u, u_x) - \partial_x((\partial_{u_x} f)(x, u, u_x))] \quad (1.2)$$

is the most general quasi-linear Hamiltonian (local) nonlinearity. Note that  $\mathcal{N}_4$  contains as many derivatives as the linear part  $\partial_{xxx}$ . The equation (1.1) is the Hamiltonian PDE  $u_t = \partial_x \nabla H(u)$  where  $\nabla H$  denotes the  $L^2(\mathbb{T}_x)$  gradient of the Hamiltonian

$$H(u) = \int_{\mathbb{T}} \frac{u_x^2}{2} + u^3 + f(x, u, u_x) dx \quad (1.3)$$

on the real phase space

$$H_0^1(\mathbb{T}_x) := \left\{ u(x) \in H^1(\mathbb{T}, \mathbb{R}) : \int_{\mathbb{T}} u(x) dx = 0 \right\}. \quad (1.4)$$

We assume that the ‘‘Hamiltonian density’’  $f \in C^q(\mathbb{T} \times \mathbb{R} \times \mathbb{R}; \mathbb{R})$  for some  $q$  large enough, and that

$$f = f_5(u, u_x) + f_{\geq 6}(x, u, u_x), \quad (1.5)$$

where  $f_5(u, u_x)$  denotes the homogeneous component of  $f$  of degree 5 and  $f_{\geq 6}$  collects all the higher order terms. By (1.5) the nonlinearity  $\mathcal{N}_4$  vanishes of order 4 at  $u = 0$  and (1.1) may be seen, close to the origin, as a ‘‘small’’ perturbation of the KdV equation

$$u_t + u_{xxx} - 6uu_x = 0, \quad (1.6)$$

which is completely integrable. Actually, the KdV equation (1.6) may be described by global analytic action-angle variables, see [20] and the references therein.

A natural question is to know whether the periodic, quasi-periodic or almost periodic solutions of (1.6) persist under small perturbations. This is the content of KAM theory.

The first KAM results for PDEs have been obtained for 1-d semilinear Schrödinger and wave equations by Kuksin [22], Wayne [31], Craig-Wayne [11], Pöschel [25], see [10], [24] and references therein. For PDEs in higher space dimension the theory has been more recently extended by Bourgain [9], Eliasson-Kuksin [12], and Berti-Bolle [5], Geng-Xu-You [13], Procesi-Procesi [28]-[27], Wang [30].

For *unbounded* perturbations the first KAM results have been proved by Kuksin [23] and Kappeler-Pöschel [20] for KdV (see also Bourgain [8]), and more recently by Liu-Yuan [19], Zhang-Gao-Yuan [32] for derivative NLS, and by Berti-Biasco-Procesi [3]-[4] for derivative NLW. For a recent survey of known results for KdV, we refer to [14].

The KAM theorems in [23], [20] prove the persistence of the finite-gap solutions of the integrable KdV (1.6) under semilinear Hamiltonian perturbations  $\varepsilon \partial_x(\partial_u f)(x, u)$ , namely when the density  $f$  is independent of  $u_x$ , so that (1.2) is a differential operator of order 1 (note that in [24] such nonlinearities are called ‘‘quasi-linear’’ and (1.2) ‘‘strongly nonlinear’’). The key point is that the frequencies of KdV grow as  $\sim j^3$  and the difference  $|j^3 - i^3| \geq (j^2 + i^2)/2$ ,  $i \neq j$ , so that KdV gains (outside the diagonal) two derivatives. This approach also works for Hamiltonian pseudo-differential perturbations of order 2 (in space), using the improved Kuksin’s lemma in [19]. However it does *not* work for a general quasi-linear perturbation as in (1.2), which is a nonlinear differential operator of the *same* order (i.e. 3) as the constant coefficient linear operator  $\partial_{xxx}$ . Such a strongly nonlinear perturbation term makes the KAM question quite delicate because of the possible phenomenon of formation of singularities in finite time, see Lax [18], Klainerman-Majda [21] for quasi-linear wave equations, see also section 1.4 of [24]. For example, Kappeler-Pöschel [20] (Remark 3, page 19) wrote: ‘‘*It would be interesting to obtain perturbation results which also include terms of higher order, at least in the region where the KdV approximation is valid. However, results of this type are still out of reach, if true at all.*’’

In this paper we give the first positive answer to this problem, proving the existence of small amplitude, linearly stable, quasi-periodic solutions of (1.1), see Theorem 1.1. Note that (1.1) does not depend on external parameters. Moreover the KdV equation (1.1) is a *completely resonant* PDE, namely the

linearized equation at the origin is the linear Airy equation  $u_t + u_{xxx} = 0$ , which possesses only the  $2\pi$ -periodic in time solutions

$$u(t, x) = \sum_{j \in \mathbb{Z} \setminus \{0\}} u_j e^{ij^3 t} e^{ijx}. \quad (1.7)$$

Thus the existence of quasi-periodic solutions of (1.1) is a purely nonlinear phenomenon (the diophantine frequencies in (1.9) are  $O(|\xi|)$ -close to integers with  $\xi \rightarrow 0$ ) and a perturbation theory is more difficult.

The solutions that we find are localized in Fourier space close to finitely many “tangential sites”

$$S^+ := \{\bar{j}_1, \dots, \bar{j}_\nu\}, \quad S := S^+ \cup (-S^+) = \{\pm j : j \in S^+\}, \quad \bar{j}_i \in \mathbb{N} \setminus \{0\}, \quad \forall i = 1, \dots, \nu. \quad (1.8)$$

The set  $S$  is required to be even because the solutions  $u$  of (1.1) have to be real valued. Moreover, we also assume the following explicit hypotheses on  $S$ :

- (S1)  $j_1 + j_2 + j_3 \neq 0$  for all  $j_1, j_2, j_3 \in S$ .
- (S2)  $\nexists j_1, \dots, j_4 \in S$  such that  $j_1 + j_2 + j_3 + j_4 \neq 0$ ,  $j_1^3 + j_2^3 + j_3^3 + j_4^3 - (j_1 + j_2 + j_3 + j_4)^3 = 0$ .

**Theorem 1.1.** *Given  $\nu \in \mathbb{N}$ , let  $f \in C^q$  (with  $q := q(\nu)$  large enough) satisfy (1.5). Then, for all the tangential sites  $S$  as in (1.8) satisfying (S1)-(S2), the KdV equation (1.1) possesses small amplitude quasi-periodic solutions with diophantine frequency vector  $\omega := \omega(\xi) = (\omega_j)_{j \in S^+} \in \mathbb{R}^\nu$ , of the form*

$$u(t, x) = \sum_{j \in S^+} 2\sqrt{\xi_j} \cos(\omega_j t + jx) + o(\sqrt{|\xi|}), \quad \omega_j := j^3 - 6\xi_j j^{-1}, \quad (1.9)$$

for a “Cantor-like” set of small amplitudes  $\xi \in \mathbb{R}_+^\nu$  with density 1 at  $\xi = 0$ . The term  $o(\sqrt{|\xi|})$  is small in some  $H^s$ -Sobolev norm,  $s < q$ . These quasi-periodic solutions are linearly stable.

This result is deduced from Theorem 5.1. Let us make some comments.

1. The set of tangential sites  $S$  satisfying (S1)-(S2) can be iteratively constructed in an explicit way, see the end of section 9. After fixing  $\{\bar{j}_1, \dots, \bar{j}_n\}$ , in the choice of  $\bar{j}_{n+1}$  there are only finitely many forbidden values, while all the other infinitely many values are good choices for  $\bar{j}_{n+1}$ . In this precise sense the set  $S$  is “generic”.
2. The linear stability of the quasi-periodic solutions is discussed after (9.41). In a suitable set of symplectic coordinates  $(\psi, \eta, w)$ ,  $\psi \in \mathbb{T}^\nu$ , near the invariant torus, the linearized equations at the quasi-periodic solutions assume the form (9.41), (9.42). Actually there is a complete KAM normal form near the invariant torus (remark 6.5), see also [6].
3. A similar result holds for perturbed (focusing/defocusing) mKdV equations

$$u_t + u_{xxx} \pm \partial_x u^3 + \mathcal{N}_4(x, u, u_x, u_{xx}, u_{xxx}) = 0 \quad (1.10)$$

for tangential sites  $S$  which satisfy  $\frac{2}{2\nu-1} \sum_{i=1}^\nu \bar{j}_i^2 \notin \mathbb{Z}$ . If the density  $f(u, u_x)$  is independent on  $x$ , the result holds for *all* the choices of the tangential sites. The KdV equation (1.1) is more difficult than (1.10) because the nonlinearity is quadratic and not cubic.

An important point is that the fourth order Birkhoff normal form of KdV and mKdV is completely integrable. The present strategy of proof — that we describe in detail below — is a rather general approach for constructing small amplitude quasi-periodic solutions of quasi-linear perturbed KdV equations. For example it could be applied to generalized KdV equations with leading nonlinearity  $u^p$ ,  $p \geq 4$ , by using the normal form techniques of Procesi-Procesi [27]-[28]. A further interesting open question concerns perturbations of the finite gap solutions of KdV.

Let us describe the strategy of proof of Theorem 1.1, which involves many different arguments.

*Weak Birkhoff normal form.* Once the finite set of tangential sites  $S$  has been fixed, the first step is to perform a “weak” Birkhoff normal form (weak BNF), whose goal is to find an invariant manifold of solutions of the third order approximate KdV equation (1.1), on which the dynamics is completely

integrable, see section 3. Since the KdV nonlinearity is quadratic, two steps of weak BNF are required. The present Birkhoff map is close to the identity up to *finite dimensional* operators, see Proposition 3.1. The key advantage is that it modifies  $\mathcal{N}_4$  very mildly, only up to finite dimensional operators (see for example Lemma 7.1), and thus the spectral analysis of the linearized equations (that we shall perform in section 8) is essentially the same as if we were in the original coordinates.

The weak normal form (3.5) does not remove (or normalize) the monomials  $O(z^2)$ . This could be done. However, we do not perform such stronger normal form (called “partial BNF” in Pöschel [26]) because the corresponding Birkhoff map is close to the identity only up to an operator of order  $O(\partial_x^{-1})$ , and so it would produce, in the transformed vector field  $\mathcal{N}_4$ , terms of order  $\partial_{xx}$  and  $\partial_x$ . A fortiori, we cannot either use the full Birkhoff normal form computed in [20] for KdV, which completely diagonalizes the fourth order terms, because such Birkhoff map is only close to the identity up to a bounded operator. For the same reason, we do not use the global nonlinear Fourier transform in [20] (Birkhoff coordinates), which is close to the Fourier transform up to smoothing operators of order  $O(\partial_x^{-1})$ .

The weak BNF procedure of section 3 is sufficient to find the first nonlinear (integrable) approximation of the solutions and to extract the “frequency-to-amplitude” modulation (4.10).

In Proposition 3.1 we also remove the terms  $O(v^5)$ ,  $O(v^4z)$  in order to have sufficiently good approximate solutions so that the Nash-Moser iteration of section 9 will converge. This is necessary for KdV whose nonlinearity is quadratic at the origin. These further steps of Birkhoff normal form are not required if the nonlinearity is yet cubic as for mKdV, see Remark 3.5. To this aim, we choose the tangential sites  $S$  such that (S2) holds. We also note that we assume (1.5) because we use the conservation of momentum up to the homogeneity order 5, see (2.7).

*Action-angle and rescaling.* At this point we introduce action-angle variables on the tangential sites (section 4) and, after the rescaling (4.5), we look for quasi-periodic solutions of the Hamiltonian (4.9). Note that the coefficients of the normal form  $\mathcal{N}$  in (4.11) depend on the angles  $\theta$ , unlike the usual KAM theorems [26], [22], where the whole normal form is reduced to constant coefficients. This is because the weak BNF of section 3 did *not* normalize the quadratic terms  $O(z^2)$ . These terms are dealt with the “linear Birkhoff normal form” (linear BNF) in sections 8.4, 8.5. In some sense here the “partial” Birkhoff normal form of [26] is split into the weak BNF of section 3 and the linear BNF of sections 8.4, 8.5.

The action-angle variables are convenient for proving the stability of the solutions.

*The nonlinear functional setting.* We look for a zero of the nonlinear operator (5.6), whose unknown is the embedded torus and the frequency  $\omega$  is seen as an “external” parameter. The solution is obtained by a Nash-Moser iterative scheme in Sobolev scales. The key step is to construct (for  $\omega$  restricted to a suitable Cantor-like set) an approximate inverse (*à la* Zehnder [33]) of the linearized operator at any approximate solution. Roughly, this means to find a linear operator which is an inverse at an exact solution. A major difficulty is that the tangential and the normal dynamics near an invariant torus are strongly coupled.

This difficulty is overcome by implementing the abstract procedure in Berti-Bolle [6]-[7] developed in order to prove existence of quasi-periodic solutions for autonomous NLW (and NLS) with a multiplicative potential. This approach reduces the search of an approximate inverse for (5.6) to the invertibility of a quasi-periodically forced PDE restricted on the normal directions. This method approximately decouples the “tangential” and the “normal” dynamics around an approximate invariant torus, introducing a suitable set of symplectic variables  $(\psi, \eta, w)$  near the torus, see (6.18). Note that, in the first line of (6.18),  $\psi$  is the “natural” angle variable which coordinates the torus, and, in the third line, the normal variable  $z$  is only translated by the component  $z_0(\psi)$  of the torus. The second line completes this transformation to a symplectic one. The canonicity of this map is proved in [6] using the isotropy of the approximate invariant torus  $i_\delta$ , see Lemma 6.3. The change of variable (6.18) brings the torus  $i_\delta$  “at the origin”. The advantage is that the second equation in (6.28) (which corresponds to the action variables of the torus) can be immediately solved, see (6.30). Then it remains to solve the third equation (6.31), i.e. to invert the linear operator  $\mathcal{L}_\omega$ . This is, up to finite dimensional remainders, a quasi-periodic Hamiltonian linear Airy equation perturbed by a variable coefficients differential operator of order  $O(\partial_{xxx})$ . The exact form of  $\mathcal{L}_\omega$  is obtained in Proposition 7.6.

*Reduction of the linearized operator in the normal directions.* In section 8 we conjugate the variable coef-

ficients operator  $\mathcal{L}_\omega$  in (7.34) to a diagonal operator with constant coefficients which describes infinitely many harmonic oscillators

$$\dot{v}_j + \mu_j^\infty v_j = 0, \quad \mu_j^\infty := i(-m_3 j^3 + m_1 j) + r_j^\infty \in i\mathbb{R}, \quad j \notin S, \quad (1.11)$$

where the constants  $m_3 - 1, m_1 \in \mathbb{R}$  and  $\sup_j |r_j^\infty|$  are small, see Theorem 8.25. The main perturbative effect to the spectrum (and the eigenfunctions) of  $\mathcal{L}_\omega$  is clearly due to the term  $a_1(\omega t, x)\partial_{xxx}$  (see (7.34)), and it is too strong for the usual reducibility KAM techniques to work directly. The conjugacy of  $\mathcal{L}_\omega$  with (1.11) is obtained in several steps. The first task (obtained in sections 8.1-8.6) is to conjugate  $\mathcal{L}_\omega$  to another Hamiltonian operator of  $H_S^\perp$  with constant coefficients

$$\mathcal{L}_6 := \omega \cdot \partial_\varphi + m_3 \partial_{xxx} + m_1 \partial_x + R_6, \quad m_1, m_3 \in \mathbb{R}, \quad (1.12)$$

up to a small bounded remainder  $R_6 = O(\partial_x^0)$ , see (8.113). This expansion of  $\mathcal{L}_\omega$  in “decreasing symbols” with constant coefficients is similar to [2], and it is somehow in the spirit of the works of Iooss, Plotnikov and Toland [17]-[16] in water waves theory, and Baldi [1] for Benjamin-Ono. It is obtained by transformations which are very different from the usual KAM changes of variables. There are several differences with respect to [2]:

1. The first step is to eliminate the  $x$ -dependence from the coefficient  $a_1(\omega t, x)\partial_{xxx}$  of the Hamiltonian operator  $\mathcal{L}_\omega$ . We cannot use the symplectic transformation  $\mathcal{A}$  defined in (8.1), used in [2], because  $\mathcal{L}_\omega$  acts on the normal subspace  $H_S^\perp$  only, and not on the whole Sobolev space as in [2]. We can not use the restricted map  $\mathcal{A}_\perp := \Pi_S^\perp \mathcal{A} \Pi_S^\perp$  which is *not* symplectic. In order to find a symplectic diffeomorphism of  $H_S^\perp$  near  $\mathcal{A}_\perp$ , the first observation is to realize  $\mathcal{A}$  as the flow map of the time dependent Hamiltonian transport linear PDE (8.3). Thus we conjugate  $\mathcal{L}_\omega$  with the flow map of the projected Hamiltonian equation (8.5). In Lemma 8.2 we prove that it differs from  $\mathcal{A}_\perp$  up to finite dimensional operators. A technical, but important, fact is that the remainders produced after this conjugation of  $\mathcal{L}_\omega$  remain of the finite dimensional form (7.7), see Lemma 8.3.

This step may be seen as a quantitative application of the Egorov theorem, see [29], which describes how the principal symbol of a pseudo-differential operator (here  $a_1(\omega t, x)\partial_{xxx}$ ) transforms under the flow of a linear hyperbolic PDE (here (8.5)).

2. Since the weak BNF procedure of section 3 did not touch the quadratic terms  $O(z^2)$ , the operator  $\mathcal{L}_\omega$  has variable coefficients also at the orders  $O(\varepsilon)$  and  $O(\varepsilon^2)$ , see (7.34)-(7.35). These terms cannot be reduced to constants by the perturbative scheme in [2], which applies to terms  $R$  such that  $R\gamma^{-1} \ll 1$  where  $\gamma$  is the diophantine constant of the frequency vector  $\omega$ . Here, since KdV is completely resonant, such  $\gamma = o(\varepsilon^2)$ , see (5.4). These terms are reduced to constant coefficients in sections 8.4-8.5 by means of purely algebraic arguments (linear BNF), which, ultimately, stem from the complete integrability of the fourth order BNF of the KdV equation (1.6), see [20].

The order of the transformations of sections 8.1-8.7 used to reduce  $\mathcal{L}_\omega$  is not accidental. The first two steps in sections 8.1, 8.2 reduce to constant coefficients the quasi-linear term  $O(\partial_{xxx})$  and eliminate the term  $O(\partial_{xx})$ , see (8.45) (the second transformation is a time quasi-periodic reparametrization of time). Then, in section 8.3, we apply the transformation  $\mathcal{T}$  (8.64) in such a way that the space average of the coefficient  $d_1(\varphi, \cdot)$  in (8.65) is constant. This is done in view of the applicability of the descent method in section 8.6. All these transformations are composition operators induced by diffeomorphisms of the torus. Therefore they are well-defined operators of a Sobolev space into itself, but their decay norm is infinite! We perform the transformation  $\mathcal{T}$  *before* the linear Birkhoff normal form steps of sections 8.4-8.5, because  $\mathcal{T}$  is a change of variable that preserves the form (7.7) of the remainders (it is not evident after the Birkhoff normal form). The Birkhoff transformations are symplectic maps of the form  $I + \varepsilon O(\partial_x^{-1})$ . Thanks to this property the coefficient  $d_1(\varphi, x)$  obtained in step 8.3 is *not* changed by these Birkhoff maps. The transformation in section 8.6 is one step of “descent method” which transforms  $d_1(\varphi, x)\partial_x$  into a constant  $m_1\partial_x$ . It is at this point of the regularization procedure that the assumption (S1) on the tangential sites is used, so that the space average of the function  $q_{>2}$  is zero, see Lemma 7.5. Actually we

only need that the average of the function in (7.33) is zero. If  $f_5 = 0$  (see (1.5)) then (S1) is not required. This completes the task of conjugating  $\mathcal{L}_\omega$  to  $\mathcal{L}_6$  in (1.12).

Finally, in section 8.7 we apply the abstract reducibility Theorem 4.2 in [2], based on a quadratic KAM scheme, which completely diagonalizes the linearized operator, obtaining (1.11). The required smallness condition (8.115) for  $R_6$  holds. Indeed the biggest term in  $R_6$  comes from the conjugation of  $\varepsilon \partial_x v_\varepsilon(\theta_0(\varphi), y_\delta(\varphi))$  in (7.35). The linear BNF procedure of section 8.4 had eliminated its main contribution  $\varepsilon \partial_x v_\varepsilon(\varphi, 0)$ . It remains  $\varepsilon \partial_x (v_\varepsilon(\theta_0(\varphi), y_\delta(\varphi)) - v_\varepsilon(\varphi, 0))$  which has size  $O(\varepsilon^{7-2b} \gamma^{-1})$  due to the estimate (6.4) of the approximate solution. This term enters in the variable coefficients of  $d_1(\varphi, x) \partial_x$  and  $d_0(\varphi, x) \partial_x^0$ . The first one had been reduced to the constant operator  $m_1 \partial_x$  by the descent method of section 8.6. The latter term is an operator of order  $O(\partial_x^0)$  which satisfies (8.115). Thus  $\mathcal{L}_6$  may be diagonalized by the iterative scheme of Theorem 4.2 in [2] which requires the smallness condition  $O(\varepsilon^{7-2b} \gamma^{-2}) \ll 1$ . This is the content of section 8.7.

*The Nash-Moser iteration.* In section 9 we perform the nonlinear Nash-Moser iteration which finally proves Theorem 5.1 and, therefore, Theorem 1.1. The optimal smallness condition required for the convergence of the scheme is  $\varepsilon \|\mathcal{F}(\varphi, 0, 0)\|_{s_0+\mu} \gamma^{-2} \ll 1$ , see (9.5). It is verified because  $\|X_P(\varphi, 0, 0)\|_s \leq_s \varepsilon^{6-2b}$  (see (5.15)), which, in turn, is a consequence of having eliminated the terms  $O(v^5), O(v^4 z)$  from the original Hamiltonian (3.1), see (3.5). This requires the condition (S2).

*Acknowledgements.* We thank M. Procesi, P. Bolle and T. Kappeler for many useful discussions. This research was supported by the European Research Council under FP7, and partially by the grants STAR 2013 and PRIN 2012 ‘‘Variational and perturbative aspects of nonlinear differential problems’’.

## 2 Preliminaries

### 2.1 Hamiltonian formalism of KdV

The Hamiltonian vector field  $X_H$  generated by a Hamiltonian  $H : H_0^1(\mathbb{T}_x) \rightarrow \mathbb{R}$  is  $X_H(u) := \partial_x \nabla H(u)$ , because

$$dH(u)[h] = (\nabla H(u), h)_{L^2(\mathbb{T}_x)} = \Omega(X_H(u), h), \quad \forall u, h \in H_0^1(\mathbb{T}_x),$$

where  $\Omega$  is the non-degenerate symplectic form

$$\Omega(u, v) := \int_{\mathbb{T}} (\partial_x^{-1} u) v \, dx, \quad \forall u, v \in H_0^1(\mathbb{T}_x), \quad (2.1)$$

and  $\partial_x^{-1} u$  is the periodic primitive of  $u$  with zero average. Note that

$$\partial_x \partial_x^{-1} = \partial_x^{-1} \partial_x = \pi_0, \quad \pi_0(u) := u - \frac{1}{2\pi} \int_{\mathbb{T}} u(x) \, dx. \quad (2.2)$$

A map is symplectic if it preserves the 2-form  $\Omega$ .

We also remind that the Poisson bracket between two functions  $F, G : H_0^1(\mathbb{T}_x) \rightarrow \mathbb{R}$  is

$$\{F(u), G(u)\} := \Omega(X_F, X_G) = \int_{\mathbb{T}} \nabla F(u) \partial_x \nabla G(u) \, dx. \quad (2.3)$$

The linearized KdV equation at  $u$  is

$$h_t = \partial_x (\partial_u \nabla H)(u)[h] = X_K(h),$$

where  $X_K$  is the KdV Hamiltonian vector field with quadratic Hamiltonian  $K = \frac{1}{2} ((\partial_u \nabla H)(u)[h], h)_{L^2(\mathbb{T}_x)} = \frac{1}{2} (\partial_{uu} H)(u)[h, h]$ . By the Schwartz theorem, the Hessian operator  $A := (\partial_u \nabla H)(u)$  is symmetric, namely  $A^T = A$ , with respect to the  $L^2$ -scalar product.

**Dynamical systems formulation.** It is convenient to regard the KdV equation also in the Fourier representation

$$u(x) = \sum_{j \in \mathbb{Z} \setminus \{0\}} u_j e^{ijx}, \quad u(x) \longleftrightarrow u := (u_j)_{j \in \mathbb{Z} \setminus \{0\}}, \quad u_{-j} = \overline{u_j}, \quad (2.4)$$



where the Fourier indices  $j \in \mathbb{Z} \setminus \{0\}$  by the definition (1.4) of the phase space and  $u_{-j} = \bar{u}_j$  because  $u(x)$  is real-valued. The symplectic structure writes

$$\Omega = \frac{1}{2} \sum_{j \neq 0} \frac{1}{ij} du_j \wedge du_{-j} = \sum_{j \geq 1} \frac{1}{ij} du_j \wedge du_{-j}, \quad \Omega(u, v) = \sum_{j \neq 0} \frac{1}{ij} u_j v_{-j} = \sum_{j \neq 0} \frac{1}{ij} u_j \bar{v}_j, \quad (2.5)$$

the Hamiltonian vector field  $X_H$  and the Poisson bracket  $\{F, G\}$  are

$$[X_H(u)]_j = ij(\partial_{u_{-j}} H)(u), \quad \forall j \neq 0, \quad \{F(u), G(u)\} = - \sum_{j \neq 0} ij(\partial_{u_{-j}} F)(u)(\partial_{u_j} G)(u). \quad (2.6)$$

**Conservation of momentum.** A Hamiltonian

$$H(u) = \sum_{j_1, \dots, j_n \in \mathbb{Z} \setminus \{0\}} H_{j_1, \dots, j_n} u_{j_1} \dots u_{j_n}, \quad u(x) = \sum_{j \in \mathbb{Z} \setminus \{0\}} u_j e^{ijx}, \quad (2.7)$$

homogeneous of degree  $n$ , *preserves the momentum* if the coefficients  $H_{j_1, \dots, j_n}$  are zero for  $j_1 + \dots + j_n \neq 0$ , so that the sum in (2.7) is restricted to integers such that  $j_1 + \dots + j_n = 0$ . Equivalently,  $H$  preserves the momentum if  $\{H, M\} = 0$ , where  $M$  is the momentum  $M(u) := \int_{\mathbb{T}} u^2 dx = \sum_{j \in \mathbb{Z} \setminus \{0\}} u_j u_{-j}$ . The homogeneous components of degree  $\leq 5$  of the KdV Hamiltonian  $H$  in (1.3) preserve the momentum because, by (1.5), the homogeneous component  $f_5$  of degree 5 does not depend on the space variable  $x$ .

**Tangential and normal variables.** Let  $\bar{j}_1, \dots, \bar{j}_\nu \geq 1$  be  $\nu$  distinct integers, and  $S^+ := \{\bar{j}_1, \dots, \bar{j}_\nu\}$ . Let  $S$  be the symmetric set in (1.8), and  $S^c := \{j \in \mathbb{Z} \setminus \{0\} : j \notin S\}$  its complementary set in  $\mathbb{Z} \setminus \{0\}$ . We decompose the phase space as

$$H_0^1(\mathbb{T}_x) := H_S \oplus H_{S^c}^\perp, \quad H_S := \text{span}\{e^{ijx} : j \in S\}, \quad H_{S^c}^\perp := \left\{ u = \sum_{j \in S^c} u_j e^{ijx} \in H_0^1(\mathbb{T}_x) \right\}, \quad (2.8)$$

and we denote by  $\Pi_S, \Pi_{S^c}^\perp$  the corresponding orthogonal projectors. Accordingly we decompose

$$u = v + z, \quad v = \Pi_S u := \sum_{j \in S} u_j e^{ijx}, \quad z = \Pi_{S^c}^\perp u := \sum_{j \in S^c} u_j e^{ijx}, \quad (2.9)$$

where  $v$  is called the *tangential* variable and  $z$  the *normal* one. We shall sometimes identify  $v \equiv (v_j)_{j \in S}$  and  $z \equiv (z_j)_{j \in S^c}$ . The subspaces  $H_S$  and  $H_{S^c}^\perp$  are *symplectic*. The dynamics of these two components is quite different. On  $H_S$  we shall introduce the action-angle variables, see (4.1). The linear frequencies of oscillations on the tangential sites are

$$\bar{\omega} := (\bar{j}_1^3, \dots, \bar{j}_\nu^3) \in \mathbb{N}^\nu. \quad (2.10)$$

## 2.2 Functional setting

**Norms.** Along the paper we shall use the notation

$$\|u\|_s := \|u\|_{H^s(\mathbb{T}^{\nu+1})} := \|u\|_{H_{\varphi, x}^s} \quad (2.11)$$

to denote the Sobolev norm of functions  $u = u(\varphi, x)$  in the Sobolev space  $H^s(\mathbb{T}^{\nu+1})$ . We shall denote by  $\|\cdot\|_{H_x^s}$  the Sobolev norm in the phase space of functions  $u := u(x) \in H^s(\mathbb{T})$ . Moreover  $\|\cdot\|_{H_\varphi^s}$  will denote the Sobolev norm of scalar functions, like the Fourier components  $u_j(\varphi)$ .

We fix  $s_0 := (\nu + 2)/2$  so that  $H^{s_0}(\mathbb{T}^{\nu+1}) \hookrightarrow L^\infty(\mathbb{T}^{\nu+1})$  and the spaces  $H^s(\mathbb{T}^{\nu+1})$ ,  $s > s_0$ , are an algebra. At the end of this section we report interpolation properties of the Sobolev norm that will be currently used along the paper. We shall also denote

$$H_{S^\perp}^s(\mathbb{T}^{\nu+1}) := \left\{ u \in H^s(\mathbb{T}^{\nu+1}) : u(\varphi, \cdot) \in H_S^\perp \quad \forall \varphi \in \mathbb{T}^\nu \right\}, \quad (2.12)$$

$$H_S^s(\mathbb{T}^{\nu+1}) := \left\{ u \in H^s(\mathbb{T}^{\nu+1}) : u(\varphi, \cdot) \in H_S \quad \forall \varphi \in \mathbb{T}^\nu \right\}. \quad (2.13)$$

For a function  $u : \Omega_o \rightarrow E$ ,  $\omega \mapsto u(\omega)$ , where  $(E, \|\cdot\|_E)$  is a Banach space and  $\Omega_o$  is a subset of  $\mathbb{R}^\nu$ , we define the sup-norm and the Lipschitz semi-norm

$$\|u\|_E^{\text{sup}} := \|u\|_{E, \Omega_o}^{\text{sup}} := \sup_{\omega \in \Omega_o} \|u(\omega)\|_E, \quad \|u\|_E^{\text{lip}} := \|u\|_{E, \Omega_o}^{\text{lip}} := \sup_{\omega_1 \neq \omega_2} \frac{\|u(\omega_1) - u(\omega_2)\|_E}{|\omega_1 - \omega_2|}, \quad (2.14)$$

and, for  $\gamma > 0$ , the Lipschitz norm

$$\|u\|_E^{\text{Lip}(\gamma)} := \|u\|_{E, \Omega_o}^{\text{Lip}(\gamma)} := \|u\|_E^{\text{sup}} + \gamma \|u\|_E^{\text{lip}}. \quad (2.15)$$

If  $E = H^s$  we simply denote  $\|u\|_{H^s}^{\text{Lip}(\gamma)} := \|u\|_s^{\text{Lip}(\gamma)}$ . We shall use the notation

$$a \leq_s b \iff a \leq C(s)b \quad \text{for some constant } C(s) > 0.$$

**Matrices with off-diagonal decay.** A linear operator can be identified, as usual, with its matrix representation. We recall the definition of the  $s$ -decay norm (introduced in [5]) of an infinite dimensional matrix. This norm is used in [2] for the KAM reducibility scheme of the linearized operators.

**Definition 2.1.** *The  $s$ -decay norm of an infinite dimensional matrix  $A := (A_{i_1}^{i_2})_{i_1, i_2 \in \mathbb{Z}^b}$ ,  $b \geq 1$ , is*

$$|A|_s^2 := \sum_{i \in \mathbb{Z}^b} \langle i \rangle^{2s} \left( \sup_{i_1 - i_2 = i} |A_{i_1}^{i_2}| \right)^2. \quad (2.16)$$

For parameter dependent matrices  $A := A(\omega)$ ,  $\omega \in \Omega_o \subseteq \mathbb{R}^\nu$ , the definitions (2.14) and (2.15) become

$$|A|_s^{\text{sup}} := \sup_{\omega \in \Omega_o} |A(\omega)|_s, \quad |A|_s^{\text{lip}} := \sup_{\omega_1 \neq \omega_2} \frac{|A(\omega_1) - A(\omega_2)|_s}{|\omega_1 - \omega_2|}, \quad |A|_s^{\text{Lip}(\gamma)} := |A|_s^{\text{sup}} + \gamma |A|_s^{\text{lip}}. \quad (2.17)$$

Such a norm is modeled on the behavior of matrices representing the multiplication operator by a function. Actually, given a function  $p \in H^s(\mathbb{T}^b)$ , the multiplication operator  $h \mapsto ph$  is represented by the Töplitz matrix  $T_i^{i'} = p_{i-i'}$  and  $|T|_s = \|p\|_s$ . If  $p = p(\omega)$  is a Lipschitz family of functions, then

$$|T|_s^{\text{Lip}(\gamma)} = \|p\|_s^{\text{Lip}(\gamma)}. \quad (2.18)$$

The  $s$ -norm satisfies classical algebra and interpolation inequalities, see [2].

**Lemma 2.1.** *Let  $A = A(\omega)$  and  $B = B(\omega)$  be matrices depending in a Lipschitz way on the parameter  $\omega \in \Omega_o \subset \mathbb{R}^\nu$ . Then for all  $s \geq s_0 > b/2$  there are  $C(s) \geq C(s_0) \geq 1$  such that*

$$|AB|_s^{\text{Lip}(\gamma)} \leq C(s) |A|_s^{\text{Lip}(\gamma)} |B|_s^{\text{Lip}(\gamma)}, \quad (2.19)$$

$$|AB|_s^{\text{Lip}(\gamma)} \leq C(s) |A|_s^{\text{Lip}(\gamma)} |B|_{s_0}^{\text{Lip}(\gamma)} + C(s_0) |A|_{s_0}^{\text{Lip}(\gamma)} |B|_s^{\text{Lip}(\gamma)}. \quad (2.20)$$

The  $s$ -decay norm controls the Sobolev norm, namely

$$\|Ah\|_s^{\text{Lip}(\gamma)} \leq C(s) (|A|_{s_0}^{\text{Lip}(\gamma)} \|h\|_s^{\text{Lip}(\gamma)} + |A|_s^{\text{Lip}(\gamma)} \|h\|_{s_0}^{\text{Lip}(\gamma)}). \quad (2.21)$$

Let now  $b := \nu + 1$ . An important sub-algebra is formed by the *Töplitz in time matrices* defined by

$$A_{(l_1, j_1)}^{(l_2, j_2)} := A_{j_1}^{j_2}(l_1 - l_2), \quad (2.22)$$

whose decay norm (2.16) is

$$|A|_s^2 = \sum_{j \in \mathbb{Z}, l \in \mathbb{Z}^\nu} \left( \sup_{j_1 - j_2 = j} |A_{j_1}^{j_2}(l)| \right)^2 \langle l, j \rangle^{2s}. \quad (2.23)$$

These matrices are identified with the  $\varphi$ -dependent family of operators

$$A(\varphi) := (A_{j_1}^{j_2}(\varphi))_{j_1, j_2 \in \mathbb{Z}}, \quad A_{j_1}^{j_2}(\varphi) := \sum_{l \in \mathbb{Z}^\nu} A_{j_1}^{j_2}(l) e^{il \cdot \varphi} \quad (2.24)$$



which act on functions of the  $x$ -variable as

$$A(\varphi) : h(x) = \sum_{j \in \mathbb{Z}} h_j e^{ijx} \mapsto A(\varphi)h(x) = \sum_{j_1, j_2 \in \mathbb{Z}} A_{j_1}^{j_2}(\varphi) h_{j_2} e^{ij_1 x}. \quad (2.25)$$

We still denote by  $|A(\varphi)|_s$  the  $s$ -decay norm of the matrix in (2.24). As in [2], all the transformations that we shall construct in this paper are of this type (with  $j, j_1, j_2 \neq 0$  because they act on the phase space  $H_0^1(\mathbb{T}_x)$ ). This observation allows to interpret the conjugacy procedure from a dynamical point of view, see [2]-section 2.2. Let us fix some terminology.

**Definition 2.2.** *We say that:*

*the operator  $(Ah)(\varphi, x) := A(\varphi)h(\varphi, x)$  is SYMPLECTIC if each  $A(\varphi)$ ,  $\varphi \in \mathbb{T}^\nu$ , is a symplectic map of the phase space (or of a symplectic subspace like  $H_S^\perp$ );*

*the operator  $\omega \cdot \partial_\varphi - \partial_x G(\varphi)$  is HAMILTONIAN if each  $G(\varphi)$ ,  $\varphi \in \mathbb{T}^\nu$ , is symmetric;*

*an operator is REAL if it maps real-valued functions into real-valued functions.*

As well known, a Hamiltonian operator  $\omega \cdot \partial_\varphi - \partial_x G(\varphi)$  is transformed, under a symplectic map  $\mathcal{A}$ , into another Hamiltonian operator  $\omega \cdot \partial_\varphi - \partial_x E(\varphi)$ , see e.g. [2]-section 2.3.

We conclude this preliminary section recalling the following well known lemmata, see Appendix of [2].

**Lemma 2.2. (Composition)** *Assume  $f \in C^s(\mathbb{T}^d \times B_1)$ ,  $B_1 := \{y \in \mathbb{R}^m : |y| \leq 1\}$ . Then  $\forall u \in H^s(\mathbb{T}^d, \mathbb{R}^m)$  such that  $\|u\|_{L^\infty} < 1$ , the composition operator  $\tilde{f}(u)(x) := f(x, u(x))$  satisfies  $\|\tilde{f}(u)\|_s \leq C\|f\|_{C^s}(\|u\|_s + 1)$  where the constant  $C$  depends on  $s, d$ . If  $f \in C^{s+2}$  and  $\|u + h\|_{L^\infty} < 1$ , then*

$$\|\tilde{f}(u + h) - \sum_{i=0}^k \frac{\tilde{f}^{(i)}(u)}{i!} [h^i]\|_s \leq C\|f\|_{C^{s+2}} \|h\|_{L^\infty}^k (\|h\|_s + \|h\|_{L^\infty} \|u\|_s), \quad k = 0, 1.$$

*The previous statement also holds replacing  $\|\cdot\|_s$  with the norms  $\|\cdot\|_{s, \infty}$ .*

**Lemma 2.3. (Tame product).** *For  $s \geq s_0 > d/2$ ,*

$$\|uv\|_s \leq C(s_0)\|u\|_s \|v\|_{s_0} + C(s)\|u\|_{s_0} \|v\|_s, \quad \forall u, v \in H^s(\mathbb{T}^d).$$

*For  $s \geq 0$ ,  $s \in \mathbb{N}$ ,*

$$\|uv\|_s \leq \frac{3}{2}\|u\|_{L^\infty} \|v\|_s + C(s)\|u\|_{W^{s, \infty}} \|v\|_0, \quad \forall u \in W^{s, \infty}(\mathbb{T}^d), v \in H^s(\mathbb{T}^d).$$

*The above inequalities also hold for the norms  $\|\cdot\|_s^{\text{Lip}(\gamma)}$ .*

**Lemma 2.4. (Change of variable)** *Let  $p \in W^{s, \infty}(\mathbb{T}^d, \mathbb{R}^d)$ ,  $s \geq 1$ , with  $\|p\|_{W^{1, \infty}} \leq 1/2$ . Then the function  $f(x) = x + p(x)$  is invertible, with inverse  $f^{-1}(y) = y + q(y)$  where  $q \in W^{s, \infty}(\mathbb{T}^d, \mathbb{R}^d)$ , and  $\|q\|_{W^{s, \infty}} \leq C\|p\|_{W^{s, \infty}}$ . If, moreover,  $p = p_\omega$  depends in a Lipschitz way on a parameter  $\omega \in \Omega \subset \mathbb{R}^\nu$ , and  $\|D_x p_\omega\|_{L^\infty} \leq 1/2$ ,  $\forall \omega$ , then  $\|q\|_{W^{s, \infty}}^{\text{Lip}(\gamma)} \leq C\|p\|_{W^{s+1, \infty}}^{\text{Lip}(\gamma)}$ . The constant  $C := C(d, s)$  is independent of  $\gamma$ .*

*If  $u \in H^s(\mathbb{T}^d, \mathbb{C})$ , then  $(u \circ f)(x) := u(x + p(x))$  satisfies*

$$\begin{aligned} \|u \circ f\|_s &\leq C(\|u\|_s + \|p\|_{W^{s, \infty}} \|u\|_1), & \|u \circ f - u\|_s &\leq C(\|p\|_{L^\infty} \|u\|_{s+1} + \|p\|_{W^{s, \infty}} \|u\|_2), \\ \|u \circ f\|_s^{\text{Lip}(\gamma)} &\leq C(\|u\|_{s+1}^{\text{Lip}(\gamma)} + \|p\|_{W^{s, \infty}}^{\text{Lip}(\gamma)} \|u\|_2^{\text{Lip}(\gamma)}). \end{aligned}$$

*The function  $u \circ f^{-1}$  satisfies the same bounds.*

### 3 Weak Birkhoff normal form

The Hamiltonian of the perturbed KdV equation (1.1) is  $H = H_2 + H_3 + H_{\geq 5}$  (see (1.3)) where

$$H_2(u) := \frac{1}{2} \int_{\mathbb{T}} u_x^2 dx, \quad H_3(u) := \int_{\mathbb{T}} u^3 dx, \quad H_{\geq 5}(u) := \int_{\mathbb{T}} f(x, u, u_x) dx, \quad (3.1)$$

and  $f$  satisfies (1.5). According to the splitting (2.9)  $u = v + z$ ,  $v \in H_S$ ,  $z \in H_S^\perp$ , we have

$$H_2(u) = \int_{\mathbb{T}} \frac{v_x^2}{2} dx + \int_{\mathbb{T}} \frac{z_x^2}{2} dx, \quad H_3(u) = \int_{\mathbb{T}} v^3 dx + 3 \int_{\mathbb{T}} v^2 z dx + 3 \int_{\mathbb{T}} v z^2 dx + \int_{\mathbb{T}} z^3 dx. \quad (3.2)$$

For a finite-dimensional space

$$E := E_C := \text{span}\{e^{ijx} : 0 < |j| \leq C\}, \quad C > 0, \quad (3.3)$$

let  $\Pi_E$  denote the corresponding  $L^2$ -projector on  $E$ .

The notation  $R(v^{k-q}z^q)$  indicates a homogeneous polynomial of degree  $k$  in  $(v, z)$  of the form

$$R(v^{k-q}z^q) = M[\underbrace{v, \dots, v}_{(k-q) \text{ times}}, \underbrace{z, \dots, z}_{q \text{ times}}], \quad M = k\text{-linear}.$$

**Proposition 3.1. (Weak Birkhoff normal form)** *Assume Hypothesis (S2). Then there exists an analytic invertible symplectic transformation of the phase space  $\Phi_B : H_0^1(\mathbb{T}_x) \rightarrow H_0^1(\mathbb{T}_x)$  of the form*

$$\Phi_B(u) = u + \Psi(u), \quad \Psi(u) = \Pi_E \Psi(\Pi_E u), \quad (3.4)$$

where  $E$  is a finite-dimensional space as in (3.3), such that the transformed Hamiltonian is

$$\mathcal{H} := H \circ \Phi_B = H_2 + \mathcal{H}_3 + \mathcal{H}_4 + \mathcal{H}_5 + \mathcal{H}_{\geq 6}, \quad (3.5)$$

where  $H_2$  is defined in (3.1),

$$\mathcal{H}_3 := \int_{\mathbb{T}} z^3 dx + 3 \int_{\mathbb{T}} v z^2 dx, \quad \mathcal{H}_4 := -\frac{3}{2} \sum_{j \in S} \frac{|u_j|^4}{j^2} + \mathcal{H}_{4,2} + \mathcal{H}_{4,3}, \quad \mathcal{H}_5 := \sum_{q=2}^5 R(v^{5-q}z^q), \quad (3.6)$$

$$\mathcal{H}_{4,2} := 6 \int_{\mathbb{T}} v z \Pi_S((\partial_x^{-1} v)(\partial_x^{-1} z)) dx + 3 \int_{\mathbb{T}} z^2 \pi_0 (\partial_x^{-1} v)^2 dx, \quad \mathcal{H}_{4,3} := R(v z^3), \quad (3.7)$$

and  $\mathcal{H}_{\geq 6}$  collects all the terms of order at least six in  $(v, z)$ .

The rest of this section is devoted to the proof of Proposition 3.1.

First, we remove the cubic terms  $\int_{\mathbb{T}} v^3 + 3 \int_{\mathbb{T}} v^2 z$  from the Hamiltonian  $H_3$  defined in (3.2). In the Fourier coordinates (2.4), we have

$$H_2 = \frac{1}{2} \sum_{j \neq 0} j^2 |u_j|^2, \quad H_3 = \sum_{j_1 + j_2 + j_3 = 0} u_{j_1} u_{j_2} u_{j_3}. \quad (3.8)$$

We look for a symplectic transformation  $\Phi^{(3)}$  of the phase space which eliminates the monomials  $u_{j_1} u_{j_2} u_{j_3}$  of  $H_3$  with at most one index outside  $S$ . Note that, by the relation  $j_1 + j_2 + j_3 = 0$ , they are *finitely* many. We look for  $\Phi^{(3)} := (\Phi_{F^{(3)}}^t)_{|t=1}$  as the time-1 flow map generated by the Hamiltonian vector field  $X_{F^{(3)}}$ , with an auxiliary Hamiltonian of the form

$$F^{(3)}(u) := \sum_{j_1 + j_2 + j_3 = 0} F_{j_1 j_2 j_3}^{(3)} u_{j_1} u_{j_2} u_{j_3}.$$

The transformed Hamiltonian is

$$\begin{aligned} H^{(3)} &:= H \circ \Phi^{(3)} = H_2 + H_3^{(3)} + H_4^{(3)} + H_{\geq 5}^{(3)}, \\ H_3^{(3)} &= H_3 + \{H_2, F^{(3)}\}, \quad H_4^{(3)} = \frac{1}{2} \{ \{H_2, F^{(3)}\}, F^{(3)} \} + \{H_3, F^{(3)}\}, \end{aligned} \quad (3.9)$$

where  $H_{\geq 5}^{(3)}$  collects all the terms of order at least five in  $(u, u_x)$ . By (3.8) and (2.6) we calculate

$$H_3^{(3)} = \sum_{j_1 + j_2 + j_3 = 0} \{1 - i(j_1^3 + j_2^3 + j_3^3) F_{j_1 j_2 j_3}^{(3)}\} u_{j_1} u_{j_2} u_{j_3}.$$

Hence, in order to eliminate the monomials with at most one index outside  $S$ , we choose

$$F_{j_1 j_2 j_3}^{(3)} := \begin{cases} \frac{1}{i(j_1^3 + j_2^3 + j_3^3)} & \text{if } (j_1, j_2, j_3) \in \mathcal{A}, \\ 0 & \text{otherwise,} \end{cases} \quad (3.10)$$

where  $\mathcal{A} := \{(j_1, j_2, j_3) \in (\mathbb{Z} \setminus \{0\})^3 : j_1 + j_2 + j_3 = 0, j_1^3 + j_2^3 + j_3^3 \neq 0, \text{ and at least 2 among } j_1, j_2, j_3 \text{ belong to } S\}$ . Note that

$$\mathcal{A} = \{(j_1, j_2, j_3) \in (\mathbb{Z} \setminus \{0\})^3 : j_1 + j_2 + j_3 = 0, \text{ and at least 2 among } j_1, j_2, j_3 \text{ belong to } S\} \quad (3.11)$$

because of the elementary relation

$$j_1 + j_2 + j_3 = 0 \quad \Rightarrow \quad j_1^3 + j_2^3 + j_3^3 = 3j_1 j_2 j_3 \neq 0 \quad (3.12)$$

being  $j_1, j_2, j_3 \in \mathbb{Z} \setminus \{0\}$ . Also note that  $\mathcal{A}$  is a finite set, actually  $\mathcal{A} \subseteq [-2C_S, 2C_S]^3$  where the tangential sites  $S \subseteq [-C_S, C_S]$ . As a consequence, the Hamiltonian vector field  $X_{F^{(3)}}$  has finite rank and vanishes outside the finite dimensional subspace  $E := E_{2C_S}$  (see (3.3)), namely

$$X_{F^{(3)}}(u) = \Pi_E X_{F^{(3)}}(\Pi_E u).$$

Hence its flow  $\Phi^{(3)} : H_0^1(\mathbb{T}_x) \rightarrow H_0^1(\mathbb{T}_x)$  has the form (3.4) and it is analytic.

By construction, all the monomials of  $H_3$  with at least two indices outside  $S$  are not modified by the transformation  $\Phi^{(3)}$ . Hence (see (3.2)) we have

$$H_3^{(3)} = \int_{\mathbb{T}} z^3 dx + 3 \int_{\mathbb{T}} v z^2 dx. \quad (3.13)$$

We now compute the fourth order term  $H_4^{(3)} = \sum_{i=0}^4 H_{4,i}^{(3)}$  in (3.9), where  $H_{4,i}^{(3)}$  is of type  $R(v^{4-i} z^i)$ .

**Lemma 3.2.** *One has (recall the definition (2.2) of  $\pi_0$ )*

$$H_{4,0}^{(3)} := \frac{3}{2} \int_{\mathbb{T}} v^2 \pi_0 [(\partial_x^{-1} v)^2] dx, \quad H_{4,2}^{(3)} := 6 \int_{\mathbb{T}} v z \Pi_S [(\partial_x^{-1} v)(\partial_x^{-1} z)] dx + 3 \int_{\mathbb{T}} z^2 \pi_0 [(\partial_x^{-1} v)^2] dx. \quad (3.14)$$

*Proof.* We write  $H_3 = H_{3, \leq 1} + H_3^{(3)}$  where  $H_{3, \leq 1}(u) := \int_{\mathbb{T}} v^3 dx + 3 \int_{\mathbb{T}} v^2 z dx$ . Then, by (3.9), we get

$$H_4^{(3)} = \frac{1}{2} \{H_{3, \leq 1}, F^{(3)}\} + \{H_3^{(3)}, F^{(3)}\}. \quad (3.15)$$

By (3.10), (3.12), the auxiliary Hamiltonian may be written as

$$F^{(3)}(u) = -\frac{1}{3} \sum_{(j_1, j_2, j_3) \in \mathcal{A}} \frac{u_{j_1} u_{j_2} u_{j_3}}{(ij_1)(ij_2)(ij_3)} = -\frac{1}{3} \int_{\mathbb{T}} (\partial_x^{-1} v)^3 dx - \int_{\mathbb{T}} (\partial_x^{-1} v)^2 (\partial_x^{-1} z) dx.$$

Hence, using that the projectors  $\Pi_S, \Pi_S^\perp$  are self-adjoint and  $\partial_x^{-1}$  is skew-selfadjoint,

$$\nabla F^{(3)}(u) = \partial_x^{-1} \{(\partial_x^{-1} v)^2 + 2\Pi_S [(\partial_x^{-1} v)(\partial_x^{-1} z)]\} \quad (3.16)$$

(we have used that  $\partial_x^{-1} \pi_0 = \partial_x^{-1}$  be the definition of  $\partial_x^{-1}$ ). Recalling the Poisson bracket definition (2.3), using that  $\nabla H_{3, \leq 1}(u) = 3v^2 + 6\Pi_S(vz)$  and (3.16), we get

$$\begin{aligned} \{H_{3, \leq 1}, F^{(3)}\} &= \int_{\mathbb{T}} \{3v^2 + 6\Pi_S(vz)\} \pi_0 \{(\partial_x^{-1} v)^2 + 2\Pi_S [(\partial_x^{-1} v)(\partial_x^{-1} z)]\} dx \\ &= 3 \int_{\mathbb{T}} v^2 \pi_0 (\partial_x^{-1} v)^2 dx + 12 \int_{\mathbb{T}} \Pi_S(vz) \Pi_S [(\partial_x^{-1} v)(\partial_x^{-1} z)] dx + R(v^3 z). \end{aligned} \quad (3.17)$$

Similarly, since  $\nabla H_3^{(3)}(u) = 3z^2 + 6\Pi_S^\perp(vz)$ ,

$$\{H_3^{(3)}, F^{(3)}\} = 3 \int_{\mathbb{T}} z^2 \pi_0 (\partial_x^{-1} v)^2 dx + R(v^3 z) + R(vz^3). \quad (3.18)$$

The lemma follows by (3.15), (3.17), (3.18).  $\square$

We now construct a symplectic map  $\Phi^{(4)}$  such that the Hamiltonian system obtained transforming  $H_2 + H_3^{(3)} + H_4^{(3)}$  possesses the invariant subspace  $H_S$  (see (2.8)) and its dynamics on  $H_S$  is integrable and non-isochronous. Hence we have to eliminate the term  $H_{4,1}^{(3)}$  (which is linear in  $z$ ), and to normalize  $H_{4,0}^{(3)}$  (which is independent of  $z$ ). We need the following elementary lemma (Lemma 13.4 in [20]).

**Lemma 3.3.** *Let  $j_1, j_2, j_3, j_4 \in \mathbb{Z}$  such that  $j_1 + j_2 + j_3 + j_4 = 0$ . Then*

$$j_1^3 + j_2^3 + j_3^3 + j_4^3 = -3(j_1 + j_2)(j_1 + j_3)(j_2 + j_3).$$

**Lemma 3.4.** *There exists a symplectic transformation  $\Phi^{(4)}$  of the form (3.4) such that*

$$H^{(4)} := H^{(3)} \circ \Phi^{(4)} = H_2 + H_3^{(3)} + H_4^{(4)} + H_{\geq 5}^{(4)}, \quad H_4^{(4)} := -\frac{3}{2} \sum_{j \in S} \frac{|u_j|^4}{j^2} + H_{4,2}^{(3)} + H_{4,3}^{(3)}, \quad (3.19)$$

where  $H_3^{(3)}$  is defined in (3.13),  $H_{4,2}^{(3)}$  in (3.14),  $H_{4,3}^{(3)} = R(vz^3)$  and  $H_{\geq 5}^{(4)}$  collects all the terms of degree at least five in  $(u, u_x)$ .

*Proof.* We look for a map  $\Phi^{(4)} := (\Phi_{F^{(4)}}^t)_{|t=1}$  which is the time 1-flow map of an auxiliary Hamiltonian

$$F^{(4)}(u) := \sum_{\substack{j_1+j_2+j_3+j_4=0 \\ \text{at least 3 indices are in } S}} F_{j_1 j_2 j_3 j_4}^{(4)} u_{j_1} u_{j_2} u_{j_3} u_{j_4}$$

with the same form of the Hamiltonian  $H_{4,0}^{(3)} + H_{4,1}^{(3)}$ . The transformed Hamiltonian is

$$H^{(4)} := H^{(3)} \circ \Phi^{(4)} = H_2 + H_3^{(3)} + H_4^{(4)} + H_{\geq 5}^{(4)}, \quad H_4^{(4)} = \{H_2, F^{(4)}\} + H_4^{(3)}, \quad (3.20)$$

where  $H_{\geq 5}^{(4)}$  collects all the terms of order at least five. We write  $H_4^{(4)} = \sum_{i=0}^4 H_{4,i}^{(4)}$  where each  $H_{4,i}^{(4)}$  is of type  $R(v^{4-i}z^i)$ . We choose the coefficients

$$F_{j_1 j_2 j_3 j_4}^{(4)} := \begin{cases} \frac{H_{j_1 j_2 j_3 j_4}^{(3)}}{i(j_1^3 + j_2^3 + j_3^3 + j_4^3)} & \text{if } (j_1, j_2, j_3, j_4) \in \mathcal{A}_4, \\ 0 & \text{otherwise,} \end{cases} \quad (3.21)$$

where

$$\mathcal{A}_4 := \{(j_1, j_2, j_3, j_4) \in (\mathbb{Z} \setminus \{0\})^4 : j_1 + j_2 + j_3 + j_4 = 0, j_1^3 + j_2^3 + j_3^3 + j_4^3 \neq 0, \text{ and at most one among } j_1, j_2, j_3, j_4 \text{ outside } S\}.$$

By this definition  $H_{4,1}^{(4)} = 0$  because there exist no integers  $j_1, j_2, j_3 \in S, j_4 \in S^c$  satisfying  $j_1 + j_2 + j_3 + j_4 = 0, j_1^3 + j_2^3 + j_3^3 + j_4^3 = 0$ , by Lemma 3.3 and the fact that  $S$  is symmetric. By construction, the terms  $H_{4,i}^{(4)} = H_{4,i}^{(3)}, i = 2, 3, 4$ , are not changed by  $\Phi^{(4)}$ . Finally, by (3.14)

$$H_{4,0}^{(4)} = \frac{3}{2} \sum_{\substack{j_1, j_2, j_3, j_4 \in S \\ j_1 + j_2 + j_3 + j_4 = 0 \\ j_1^3 + j_2^3 + j_3^3 + j_4^3 = 0 \\ j_1 + j_2, j_3 + j_4 \neq 0}} \frac{1}{(ij_3)(ij_4)} u_{j_1} u_{j_2} u_{j_3} u_{j_4}. \quad (3.22)$$

If  $j_1 + j_2 + j_3 + j_4 = 0$  and  $j_1^3 + j_2^3 + j_3^3 + j_4^3 = 0$ , then  $(j_1 + j_2)(j_1 + j_3)(j_2 + j_3) = 0$  by Lemma 3.3. We develop the sum in (3.22) with respect to the first index  $j_1$ . Since  $j_1 + j_2 \neq 0$  the possible cases are:

$$(i) \{j_2 \neq -j_1, j_3 = -j_1, j_4 = -j_2\} \quad \text{or} \quad (ii) \{j_2 \neq -j_1, j_3 \neq -j_1, j_3 = -j_2, j_4 = -j_1\}.$$

Hence, using  $u_{-j} = \bar{u}_j$  (recall (2.4)), and since  $S$  is symmetric, we have

$$\sum_{(i)} \frac{1}{j_3 j_4} u_{j_1} u_{j_2} u_{j_3} u_{j_4} = \sum_{j_1, j_2 \in S, j_2 \neq -j_1} \frac{|u_{j_1}|^2 |u_{j_2}|^2}{j_1 j_2} = \sum_{j, j' \in S} \frac{|u_j|^2 |u_{j'}|^2}{j j'} + \sum_{j \in S} \frac{|u_j|^4}{j^2} = \sum_{j \in S} \frac{|u_j|^4}{j^2}, \quad (3.23)$$

and in the second case (ii)

$$\sum_{(ii)} \frac{1}{j_3 j_4} u_{j_1} u_{j_2} u_{j_3} u_{j_4} = \sum_{j_1, j_2, j_2 \neq \pm j_1} \frac{1}{j_1 j_2} u_{j_1} u_{j_2} u_{-j_2} u_{-j_1} = \sum_{j \in S} \frac{1}{j} |u_j|^2 \left( \sum_{j_2 \neq \pm j} \frac{1}{j_2} |u_{j_2}|^2 \right) = 0. \quad (3.24)$$

Then (3.19) follows by (3.22), (3.23), (3.24).  $\square$

Note that the Hamiltonian  $H_2 + H_3^{(3)} + H_4^{(4)}$  (see (3.19)) possesses the invariant subspace  $\{z = 0\}$  and the system restricted to  $\{z = 0\}$  is completely integrable and non-isochronous (actually it is formed by  $\nu$  decoupled rotators). We shall construct quasi-periodic solutions which bifurcate from this invariant manifold.

In order to enter in a perturbative regime, we have to eliminate further monomials of  $H^{(4)}$  in (3.19). The minimal requirement for the convergence of the nonlinear Nash-Moser iteration is to eliminate the monomials  $R(v^5)$  and  $R(v^4 z)$ . Here we need the choice of the sites of Hypothesis (S2).

**Remark 3.5.** In the KAM theorems [24], [26] (and [28], [30]), as well as for the perturbed mKdV equations (1.10), these further steps of Birkhoff normal form are not required because the nonlinearity of the original PDE is yet cubic. A difficulty of KdV is that the nonlinearity is quadratic.  $\square$

We spell out Hypothesis (S2) as follows:

- (S2<sub>0</sub>). There is no choice of 5 integers  $j_1, \dots, j_5 \in S$  such that

$$j_1 + \dots + j_5 = 0, \quad j_1^3 + \dots + j_5^3 = 0. \quad (3.25)$$

- (S2<sub>1</sub>). There is no choice of 4 integers  $j_1, \dots, j_4$  in  $S$  and an integer in the complementary set  $j_5 \in S^c := (\mathbb{Z} \setminus \{0\}) \setminus S$  such that (3.25) holds.

The homogeneous component of degree 5 of  $H^{(4)}$  is

$$H_5^{(4)}(u) = \sum_{j_1 + \dots + j_5 = 0} H_{j_1, \dots, j_5}^{(4)} u_{j_1} \dots u_{j_5}.$$

We want to remove from  $H_5^{(4)}$  the terms with at most one index among  $j_1, \dots, j_5$  outside  $S$ . We consider the auxiliary Hamiltonian

$$F^{(5)} = \sum_{\substack{j_1 + \dots + j_5 = 0 \\ \text{at most one index outside } S}} F_{j_1, \dots, j_5}^{(5)} u_{j_1} \dots u_{j_5}, \quad F_{j_1, \dots, j_5}^{(5)} := \frac{H_{j_1, \dots, j_5}^{(5)}}{i(j_1^3 + \dots + j_5^3)}. \quad (3.26)$$

By Hypotheses (S2<sub>0</sub>), (S2<sub>1</sub>), if  $j_1 + \dots + j_5 = 0$  with at most one index outside  $S$  then  $j_1^3 + \dots + j_5^3 \neq 0$  and  $F^{(5)}$  is well defined. Let  $\Phi^{(5)}$  be the time 1-flow generated by  $X_{F^{(5)}}$ . The new Hamiltonian is

$$H^{(5)} := H^{(4)} \circ \Phi^{(5)} = H_2 + H_3^{(3)} + H_4^{(4)} + \{H_2, F^{(5)}\} + H_5^{(4)} + H_{\geq 6}^{(5)} \quad (3.27)$$

where, by (3.26),

$$H_5^{(5)} := \{H_2, F^{(5)}\} + H_5^{(4)} = \sum_{q=2}^5 R(v^{5-q} z^q).$$

Renaming  $\mathcal{H} := H^{(5)}$ , namely  $\mathcal{H}_n := H_n^{(n)}$ ,  $n = 3, 4, 5$ , and setting  $\Phi_B := \Phi^{(3)} \circ \Phi^{(4)} \circ \Phi^{(5)}$ , formula (3.5) follows.

The homogeneous component  $H_5^{(4)}$  preserves the momentum, see section 2.1. Hence  $F^{(5)}$  also preserves the momentum. As a consequence, also  $H_k^{(5)}$ ,  $k \leq 5$ , preserve the momentum.

Finally, since  $F^{(5)}$  is Fourier-supported on a finite set, the transformation  $\Phi^{(5)}$  is of type (3.4) (and analytic), and therefore also the composition  $\Phi_B$  is of type (3.4) (and analytic).

## 4 Action-angle variables

We now introduce action-angle variables on the tangential directions by the change of coordinates

$$\begin{cases} u_j := \sqrt{\xi_j + |j|y_j} e^{i\theta_j}, & \text{if } j \in S, \\ u_j := z_j, & \text{if } j \in S^c, \end{cases} \quad (4.1)$$

where (recall  $u_{-j} = \bar{u}_j$ )

$$\xi_{-j} = \xi_j, \quad \xi_j > 0, \quad y_{-j} = y_j, \quad \theta_{-j} = -\theta_j, \quad \theta_j, y_j \in \mathbb{R}, \quad \forall j \in S. \quad (4.2)$$

For the tangential sites  $S^+ := \{\bar{j}_1, \dots, \bar{j}_\nu\}$  we shall also denote  $\theta_{\bar{j}_i} := \theta_i$ ,  $y_{\bar{j}_i} := y_i$ ,  $\xi_{\bar{j}_i} := \xi_i$ ,  $i = 1, \dots, \nu$ . The symplectic 2-form  $\Omega$  in (2.5) (i.e. (2.1)) becomes

$$\mathcal{W} := \sum_{i=1}^{\nu} d\theta_i \wedge dy_i + \frac{1}{2} \sum_{j \in S^c \setminus \{0\}} \frac{1}{ij} dz_j \wedge dz_{-j} = \left( \sum_{i=1}^{\nu} d\theta_i \wedge dy_i \right) \oplus \Omega_{S^\perp} = d\Lambda \quad (4.3)$$

where  $\Omega_{S^\perp}$  denotes the restriction of  $\Omega$  to  $H_S^\perp$  (see (2.8)) and  $\Lambda$  is the contact 1-form on  $\mathbb{T}^\nu \times \mathbb{R}^\nu \times H_S^\perp$  defined by  $\Lambda_{(\theta, y, z)} : \mathbb{R}^\nu \times \mathbb{R}^\nu \times H_S^\perp \rightarrow \mathbb{R}$ ,

$$\Lambda_{(\theta, y, z)}[\widehat{\theta}, \widehat{y}, \widehat{z}] := -y \cdot \widehat{\theta} + \frac{1}{2} (\partial_x^{-1} z, \widehat{z})_{L^2(\mathbb{T})}. \quad (4.4)$$

Instead of working in a shrinking neighborhood of the origin, it is a convenient device to rescale the ‘‘unperturbed actions’’  $\xi$  and the action-angle variables as

$$\xi \mapsto \varepsilon^2 \xi, \quad y \mapsto \varepsilon^{2b} y, \quad z \mapsto \varepsilon^b z. \quad (4.5)$$

Then the symplectic 2-form in (4.3) transforms into  $\varepsilon^{2b} \mathcal{W}$ . Hence the Hamiltonian system generated by  $\mathcal{H}$  in (3.5) transforms into the new Hamiltonian system

$$\dot{\theta} = \partial_y H_\varepsilon(\theta, y, z), \quad \dot{y} = -\partial_\theta H_\varepsilon(\theta, y, z), \quad \dot{z} = \partial_x \nabla_z H_\varepsilon(\theta, y, z), \quad H_\varepsilon := \varepsilon^{-2b} \mathcal{H} \circ A_\varepsilon \quad (4.6)$$

where

$$A_\varepsilon(\theta, y, z) := \varepsilon v_\varepsilon(\theta, y) + \varepsilon^b z := \varepsilon \sum_{j \in S} \sqrt{\xi_j + \varepsilon^{2(b-1)} |j| y_j} e^{i\theta_j} e^{ijx} + \varepsilon^b z. \quad (4.7)$$

We shall still denote by  $X_{H_\varepsilon} = (\partial_y H_\varepsilon, -\partial_\theta H_\varepsilon, \partial_x \nabla_z H_\varepsilon)$  the Hamiltonian vector field in the variables  $(\theta, y, z) \in \mathbb{T}^\nu \times \mathbb{R}^\nu \times H_S^\perp$ .

We now write explicitly the Hamiltonian  $H_\varepsilon(\theta, y, z)$  in (4.6). The quadratic Hamiltonian  $H_2$  in (3.1) transforms into

$$\varepsilon^{-2b} H_2 \circ A_\varepsilon = \text{const} + \sum_{j \in S^+} j^3 y_j + \frac{1}{2} \int_{\mathbb{T}} z_x^2 dx, \quad (4.8)$$

and, recalling (3.6), (3.7), the Hamiltonian  $\mathcal{H}$  in (3.5) transforms into (shortly writing  $v_\varepsilon := v_\varepsilon(\theta, y)$ )

$$\begin{aligned} H_\varepsilon(\theta, y, z) &= e(\xi) + \alpha(\xi) \cdot y + \frac{1}{2} \int_{\mathbb{T}} z_x^2 dx + \varepsilon^b \int_{\mathbb{T}} z^3 dx + 3\varepsilon \int_{\mathbb{T}} v_\varepsilon z^2 dx \\ &+ \varepsilon^2 \left\{ 6 \int_{\mathbb{T}} v_\varepsilon z \Pi_S((\partial_x^{-1} v_\varepsilon)(\partial_x^{-1} z)) dx + 3 \int_{\mathbb{T}} z^2 \pi_0(\partial_x^{-1} v_\varepsilon)^2 dx \right\} - \frac{3}{2} \varepsilon^{2b} \sum_{j \in S^+} y_j^2 \\ &+ \varepsilon^{b+1} R(v_\varepsilon z^3) + \varepsilon^3 R(v_\varepsilon^3 z^2) + \varepsilon^{2+b} \sum_{q=3}^5 \varepsilon^{(q-3)(b-1)} R(v_\varepsilon^{5-q} z^q) + \varepsilon^{-2b} \mathcal{H}_{\geq 6}(\varepsilon v_\varepsilon + \varepsilon^b z) \end{aligned} \quad (4.9)$$

where  $e(\xi)$  is a constant, and the frequency-amplitude map is

$$\alpha(\xi) := \bar{\omega} + \varepsilon^2 \mathbb{A} \xi, \quad \mathbb{A} := -6 \text{diag}\{1/j\}_{j \in S^+}. \quad (4.10)$$



We write the Hamiltonian in (4.9) as

$$H_\varepsilon = \mathcal{N} + P, \quad \mathcal{N}(\theta, y, z) = \alpha(\xi) \cdot y + \frac{1}{2}(N(\theta)z, z)_{L^2(\mathbb{T})}, \quad (4.11)$$

where

$$\begin{aligned} \frac{1}{2}(N(\theta)z, z)_{L^2(\mathbb{T})} &:= \frac{1}{2}((\partial_z \nabla H_\varepsilon)(\theta, 0, 0)[z], z)_{L^2(\mathbb{T})} = \frac{1}{2} \int_{\mathbb{T}} z_x^2 dx + 3\varepsilon \int_{\mathbb{T}} v_\varepsilon(\theta, 0) z^2 dx \\ &+ \varepsilon^2 \left\{ 6 \int_{\mathbb{T}} v_\varepsilon(\theta, 0) z \Pi_S((\partial_x^{-1} v_\varepsilon(\theta, 0))(\partial_x^{-1} z)) dx + 3 \int_{\mathbb{T}} z^2 \pi_0(\partial_x^{-1} v_\varepsilon(\theta, 0))^2 dx \right\} + \dots \end{aligned} \quad (4.12)$$

and  $P := H_\varepsilon - \mathcal{N}$ .

## 5 The nonlinear functional setting

We look for an embedded invariant torus

$$i : \mathbb{T}^\nu \rightarrow \mathbb{T}^\nu \times \mathbb{R}^\nu \times H_S^\perp, \quad \varphi \mapsto i(\varphi) := (\theta(\varphi), y(\varphi), z(\varphi)) \quad (5.1)$$

of the Hamiltonian vector field  $X_{H_\varepsilon}$  filled by quasi-periodic solutions with diophantine frequency  $\omega$ . We require that  $\omega$  belongs to the set

$$\Omega_\varepsilon := \alpha([1, 2]^\nu) = \{\alpha(\xi) : \xi \in [1, 2]^\nu\} \quad (5.2)$$

where  $\alpha$  is the diffeomorphism (4.10), and, in the Hamiltonian  $H_\varepsilon$  in (4.11), we choose

$$\xi = \alpha^{-1}(\omega) = \varepsilon^{-2} \mathbb{A}^{-1}(\omega - \bar{\omega}). \quad (5.3)$$

Since any  $\omega \in \Omega_\varepsilon$  is  $\varepsilon^2$ -close to the integer vector  $\bar{\omega}$  (see (2.10)), we require that the constant  $\gamma$  in the diophantine inequality

$$|\omega \cdot l| \geq \gamma \langle l \rangle^{-\tau}, \quad \forall l \in \mathbb{Z}^\nu \setminus \{0\}, \quad \text{satisfies } \gamma = \varepsilon^{2+a} \quad \text{for some } a > 0. \quad (5.4)$$

We remark that the definition of  $\gamma$  in (5.4) is slightly stronger than the minimal condition, which is  $\gamma \leq c\varepsilon^2$  with  $c$  small enough. In addition to (5.4) we shall also require that  $\omega$  satisfies the first and second order Melnikov-non-resonance conditions (8.120).

We look for an embedded invariant torus of the modified Hamiltonian vector field  $X_{H_{\varepsilon, \zeta}} = X_{H_\varepsilon} + (0, \zeta, 0)$  which is generated by the Hamiltonian

$$H_{\varepsilon, \zeta}(\theta, y, z) := H_\varepsilon(\theta, y, z) + \zeta \cdot \theta, \quad \zeta \in \mathbb{R}^\nu. \quad (5.5)$$

Note that  $X_{H_{\varepsilon, \zeta}}$  is periodic in  $\theta$  (unlike  $H_{\varepsilon, \zeta}$ ). It turns out that an invariant torus for  $X_{H_{\varepsilon, \zeta}}$  is actually invariant for  $X_{H_\varepsilon}$ , see Lemma 6.1. We introduce the parameter  $\zeta \in \mathbb{R}^\nu$  in order to control the average in the  $y$ -component of the linearized equations. Thus we look for zeros of the nonlinear operator

$$\begin{aligned} \mathcal{F}(i, \zeta) &:= \mathcal{F}(i, \zeta, \omega, \varepsilon) := \mathcal{D}_\omega i(\varphi) - X_{H_{\varepsilon, \zeta}}(i(\varphi)) = \mathcal{D}_\omega i(\varphi) - X_{\mathcal{N}}(i(\varphi)) - X_P(i(\varphi)) + (0, \zeta, 0) \\ &:= \begin{pmatrix} \mathcal{D}_\omega \theta(\varphi) - \partial_y H_\varepsilon(i(\varphi)) \\ \mathcal{D}_\omega y(\varphi) + \partial_\theta H_\varepsilon(i(\varphi)) + \zeta \\ \mathcal{D}_\omega z(\varphi) - \partial_x \nabla_z H_\varepsilon(i(\varphi)) \end{pmatrix} = \begin{pmatrix} \mathcal{D}_\omega \Theta(\varphi) - \partial_y P(i(\varphi)) \\ \mathcal{D}_\omega y(\varphi) + \frac{1}{2} \partial_\theta (N(\theta(\varphi))z(\varphi), z(\varphi))_{L^2(\mathbb{T})} + \partial_\theta P(i(\varphi)) + \zeta \\ \mathcal{D}_\omega z(\varphi) - \partial_x N(\theta(\varphi))z(\varphi) - \partial_x \nabla_z P(i(\varphi)) \end{pmatrix} \end{aligned} \quad (5.6)$$

where  $\Theta(\varphi) := \theta(\varphi) - \varphi$  is  $(2\pi)^\nu$ -periodic and we use the short notation

$$\mathcal{D}_\omega := \omega \cdot \partial_\varphi. \quad (5.7)$$

The Sobolev norm of the periodic component of the embedded torus

$$\mathfrak{I}(\varphi) := i(\varphi) - (\varphi, 0, 0) := (\Theta(\varphi), y(\varphi), z(\varphi)), \quad \Theta(\varphi) := \theta(\varphi) - \varphi, \quad (5.8)$$

is

$$\|\mathcal{J}\|_s := \|\Theta\|_{H_\varphi^s} + \|y\|_{H_\varphi^s} + \|z\|_s \quad (5.9)$$

where  $\|z\|_s := \|z\|_{H_{\varphi,x}^s}$  is defined in (2.11). We link the rescaling (4.5) with the diophantine constant  $\gamma = \varepsilon^{2+a}$  by choosing

$$\gamma = \varepsilon^{2b}, \quad b = 1 + (a/2). \quad (5.10)$$

Other choices are possible, see Remark 5.2.

**Theorem 5.1.** *Let the tangential sites  $S$  in (1.8) satisfy (S1), (S2). Then, for all  $\varepsilon \in (0, \varepsilon_0)$ , where  $\varepsilon_0$  is small enough, there exists a Cantor-like set  $\mathcal{C}_\varepsilon \subset \Omega_\varepsilon$ , with asymptotically full measure as  $\varepsilon \rightarrow 0$ , namely*

$$\lim_{\varepsilon \rightarrow 0} \frac{|\mathcal{C}_\varepsilon|}{|\Omega_\varepsilon|} = 1, \quad (5.11)$$

such that, for all  $\omega \in \mathcal{C}_\varepsilon$ , there exists a solution  $i_\infty(\varphi) := i_\infty(\omega, \varepsilon)(\varphi)$  of  $\mathcal{D}_\omega i_\infty(\varphi) - X_{H_\varepsilon}(i_\infty(\varphi)) = 0$ . Hence the embedded torus  $\varphi \mapsto i_\infty(\varphi)$  is invariant for the Hamiltonian vector field  $X_{H_\varepsilon(\cdot, \varepsilon)}$  with  $\xi$  as in (5.3), and it is filled by quasi-periodic solutions with frequency  $\omega$ . The torus  $i_\infty$  satisfies

$$\|i_\infty(\varphi) - (\varphi, 0, 0)\|_{s_0+\mu}^{\text{Lip}(\gamma)} = O(\varepsilon^{6-2b}\gamma^{-1}) \quad (5.12)$$

for some  $\mu := \mu(\nu) > 0$ . Moreover, the torus  $i_\infty$  is LINEARLY STABLE.

Theorem 5.1 is proved in sections 6-9. It implies Theorem 1.1 where the  $\xi_j$  in (1.9) are  $\varepsilon^2 \xi_j$ ,  $\xi_j \in [1, 2]$ , in (5.3). By (5.12), going back to the variables before the rescaling (4.5), we get  $\Theta_\infty = O(\varepsilon^{6-2b}\gamma^{-1})$ ,  $y_\infty = O(\varepsilon^6\gamma^{-1})$ ,  $z_\infty = O(\varepsilon^{6-b}\gamma^{-1})$ , which, as  $b \rightarrow 1^+$ , tend to the expected optimal estimates.

**Remark 5.2.** There are other possible ways to link the rescaling (4.5) with the diophantine constant  $\gamma = \varepsilon^{2+a}$ . The choice  $\gamma > \varepsilon^{2b}$  reduces to study perturbations of an isochronous system (as in [22], [24], [26]), and it is convenient to introduce  $\xi(\omega)$  as a variable. The case  $\varepsilon^{2b} > \gamma$ , in particular  $b = 1$ , has to be dealt with a perturbation approach of a non-isochronous system à la Arnold-Kolmogorov.  $\square$

We now give the tame estimates for the composition operator induced by the Hamiltonian vector fields  $X_{\mathcal{N}}$  and  $X_P$  in (5.6), that we shall use in the next sections.

We first estimate the tame composition operator induced by  $v_\varepsilon(\theta, y)$  defined in (4.7). Since the functions  $y \mapsto \sqrt{\xi + \varepsilon^{2(b-1)}|j|y}$ ,  $\theta \mapsto e^{i\theta}$  are analytic for  $\varepsilon$  small enough and  $|y| \leq C$ , the composition Lemma 2.2 implies that, for all  $\Theta, y \in H^s(\mathbb{T}^\nu, \mathbb{R}^\nu)$ ,  $\|\Theta\|_{s_0}, \|y\|_{s_0} \leq 1$ , setting  $\theta(\varphi) := \varphi + \Theta(\varphi)$ ,  $\|v_\varepsilon(\theta(\varphi), y(\varphi))\|_s \leq_s 1 + \|\Theta\|_s + \|y\|_s$ . Hence, using also (5.3), the map  $A_\varepsilon$  in (4.7) satisfies, for all  $\|\mathcal{J}\|_{s_0}^{\text{Lip}(\gamma)} \leq 1$  (see (5.8))

$$\|A_\varepsilon(\theta(\varphi), y(\varphi), z(\varphi))\|_s^{\text{Lip}(\gamma)} \leq_s \varepsilon(1 + \|\mathcal{J}\|_s^{\text{Lip}(\gamma)}). \quad (5.13)$$

We now give tame estimates for the Hamiltonian vector fields  $X_{\mathcal{N}}, X_P, X_{H_\varepsilon}$ , see (4.11)-(4.12).

**Lemma 5.3.** *Let  $\mathcal{J}(\varphi)$  in (5.8) satisfy  $\|\mathcal{J}\|_{s_0+3}^{\text{Lip}(\gamma)} \leq C\varepsilon^{6-2b}\gamma^{-1}$ . Then*

$$\|\partial_y P(i)\|_s^{\text{Lip}(\gamma)} \leq_s \varepsilon^4 + \varepsilon^{2b}\|\mathcal{J}\|_{s+1}^{\text{Lip}(\gamma)}, \quad \|\partial_\theta P(i)\|_s^{\text{Lip}(\gamma)} \leq_s \varepsilon^{6-2b}(1 + \|\mathcal{J}\|_{s+1}^{\text{Lip}(\gamma)}) \quad (5.14)$$

$$\|\nabla_z P(i)\|_s^{\text{Lip}(\gamma)} \leq_s \varepsilon^{5-b} + \varepsilon^{6-b}\gamma^{-1}\|\mathcal{J}\|_{s+1}^{\text{Lip}(\gamma)}, \quad \|X_P(i)\|_s^{\text{Lip}(\gamma)} \leq_s \varepsilon^{6-2b} + \varepsilon^{2b}\|\mathcal{J}\|_{s+3}^{\text{Lip}(\gamma)} \quad (5.15)$$

$$\|\partial_\theta \partial_y P(i)\|_s^{\text{Lip}(\gamma)} \leq_s \varepsilon^4 + \varepsilon^5\gamma^{-1}\|\mathcal{J}\|_{s+2}^{\text{Lip}(\gamma)}, \quad \|\partial_y \nabla_z P(i)\|_s^{\text{Lip}(\gamma)} \leq_s \varepsilon^{b+3} + \varepsilon^{2b-1}\|\mathcal{J}\|_{s+2}^{\text{Lip}(\gamma)} \quad (5.16)$$

$$\|\partial_{yy} P(i) + 3\varepsilon^{2b} I_{\mathbb{R}^\nu}\|_s^{\text{Lip}(\gamma)} \leq_s \varepsilon^{2+2b} + \varepsilon^{2b+3}\gamma^{-1}\|\mathcal{J}\|_{s+2}^{\text{Lip}(\gamma)} \quad (5.17)$$

and, for all  $\widehat{v} := (\widehat{\Theta}, \widehat{y}, \widehat{z})$ ,

$$\|\partial_y d_i X_P(i)[\widehat{v}]\|_s^{\text{Lip}(\gamma)} \leq_s \varepsilon^{2b-1}(\|\widehat{v}\|_{s+3}^{\text{Lip}(\gamma)} + \|\mathcal{J}\|_{s+3}^{\text{Lip}(\gamma)}\|\widehat{v}\|_{s_0+3}^{\text{Lip}(\gamma)}) \quad (5.18)$$

$$\|d_i X_{H_\varepsilon}(i)[\widehat{v}] + (0, 0, \partial_{xxx}\widehat{z})\|_s^{\text{Lip}(\gamma)} \leq_s \varepsilon(\|\widehat{v}\|_{s+3}^{\text{Lip}(\gamma)} + \|\mathcal{J}\|_{s+3}^{\text{Lip}(\gamma)}\|\widehat{v}\|_{s_0+3}^{\text{Lip}(\gamma)}) \quad (5.19)$$

$$\|d_i^2 X_{H_\varepsilon}(i)[\widehat{v}, \widehat{v}]\|_s^{\text{Lip}(\gamma)} \leq_s \varepsilon\left(\|\widehat{v}\|_{s+3}^{\text{Lip}(\gamma)}\|\widehat{v}\|_{s_0+3}^{\text{Lip}(\gamma)} + \|\mathcal{J}\|_{s+3}^{\text{Lip}(\gamma)}(\|\widehat{v}\|_{s_0+3}^{\text{Lip}(\gamma)})^2\right). \quad (5.20)$$

In the sequel we will also use that, by the diophantine condition (5.4), the operator  $\mathcal{D}_\omega^{-1}$  (see (5.7)) is defined for all functions  $u$  with zero  $\varphi$ -average, and satisfies

$$\|\mathcal{D}_\omega^{-1}u\|_s \leq C\gamma^{-1}\|u\|_{s+\tau}, \quad \|\mathcal{D}_\omega^{-1}u\|_s^{\text{Lip}(\gamma)} \leq C\gamma^{-1}\|u\|_{s+2\tau+1}^{\text{Lip}(\gamma)}. \quad (5.21)$$

## 6 Approximate inverse

In order to implement a convergent Nash-Moser scheme that leads to a solution of  $\mathcal{F}(i, \zeta) = 0$  our aim is to construct an *approximate right inverse* (which satisfies tame estimates) of the linearized operator

$$d_{i, \zeta} \mathcal{F}(i_0, \zeta_0)[\widehat{v}, \widehat{\zeta}] = d_{i, \zeta} \mathcal{F}(i_0)[\widehat{v}, \widehat{\zeta}] = \mathcal{D}_\omega \widehat{v} - d_i X_{H_\varepsilon}(i_0(\varphi))[\widehat{v}] + (0, \widehat{\zeta}, 0), \quad (6.1)$$

see Theorem 6.10. Note that  $d_{i, \zeta} \mathcal{F}(i_0, \zeta_0) = d_{i, \zeta} \mathcal{F}(i_0)$  is independent of  $\zeta_0$  (see (5.6)).

The notion of approximate right inverse is introduced in [33]. It denotes a linear operator which is an *exact* right inverse at a solution  $(i_0, \zeta_0)$  of  $\mathcal{F}(i_0, \zeta_0) = 0$ . We want to implement the general strategy in [6]-[7] which reduces the search of an approximate right inverse of (6.1) to the search of an approximate inverse on the normal directions only.

It is well known that an invariant torus  $i_0$  with diophantine flow is isotropic (see e.g. [6]), namely the pull-back 1-form  $i_0^* \Lambda$  is closed, where  $\Lambda$  is the contact 1-form in (4.4). This is tantamount to say that the 2-form  $\mathcal{W}$  (see (4.3)) vanishes on the torus  $i_0(\mathbb{T}^\nu)$  (i.e.  $\mathcal{W}$  vanishes on the tangent space at each point  $i_0(\varphi)$  of the manifold  $i_0(\mathbb{T}^\nu)$ ), because  $i_0^* \mathcal{W} = i_0^* d\Lambda = di_0^* \Lambda$ . For an ‘‘approximately invariant’’ torus  $i_0$  the 1-form  $i_0^* \Lambda$  is only ‘‘approximately closed’’. In order to make this statement quantitative we consider

$$i_0^* \Lambda = \sum_{k=1}^\nu a_k(\varphi) d\varphi_k, \quad a_k(\varphi) := -([\partial_\varphi \theta_0(\varphi)]^T y_0(\varphi))_k + \frac{1}{2}(\partial_{\varphi_k} z_0(\varphi), \partial_x^{-1} z_0(\varphi))_{L^2(\mathbb{T})} \quad (6.2)$$

and we quantify how small is

$$i_0^* \mathcal{W} = di_0^* \Lambda = \sum_{1 \leq k < j \leq \nu} A_{kj}(\varphi) d\varphi_k \wedge d\varphi_j, \quad A_{kj}(\varphi) := \partial_{\varphi_k} a_j(\varphi) - \partial_{\varphi_j} a_k(\varphi). \quad (6.3)$$

Along this section we will always assume the following hypothesis (which will be verified at each step of the Nash-Moser iteration):

- ASSUMPTION. The map  $\omega \mapsto i_0(\omega)$  is a Lipschitz function defined on some subset  $\Omega_o \subset \Omega_\varepsilon$ , where  $\Omega_\varepsilon$  is defined in (5.2), and, for some  $\mu := \mu(\tau, \nu) > 0$ ,

$$\|\mathfrak{J}_0\|_{s_0+\mu}^{\text{Lip}(\gamma)} \leq C\varepsilon^{6-2b}\gamma^{-1}, \quad \|Z\|_{s_0+\mu}^{\text{Lip}(\gamma)} \leq C\varepsilon^{6-2b}, \quad \gamma = \varepsilon^{2+a}, \quad b := 1 + (a/2), \quad a \in (0, 1/6), \quad (6.4)$$

where  $\mathfrak{J}_0(\varphi) := i_0(\varphi) - (\varphi, 0, 0)$ , and

$$Z(\varphi) := (Z_1, Z_2, Z_3)(\varphi) := \mathcal{F}(i_0, \zeta_0)(\varphi) = \omega \cdot \partial_\varphi i_0(\varphi) - X_{H_{\varepsilon, \zeta_0}}(i_0(\varphi)). \quad (6.5)$$

**Lemma 6.1.**  $\|\zeta_0\|_{s_0}^{\text{Lip}(\gamma)} \leq C\|Z\|_{s_0}^{\text{Lip}(\gamma)}$ . If  $\mathcal{F}(i_0, \zeta_0) = 0$  then  $\zeta_0 = 0$ , namely the torus  $i_0$  is invariant for  $X_{H_\varepsilon}$ .

*Proof.* It is proved in [6] the formula

$$\zeta_0 = \int_{\mathbb{T}^\nu} -[\partial_\varphi y_0(\varphi)]^T Z_1(\varphi) + [\partial_\varphi \theta_0(\varphi)]^T Z_2(\varphi) - [\partial_\varphi z_0(\varphi)]^T \partial_x^{-1} Z_3(\varphi) d\varphi.$$

Hence the lemma follows by (6.4) and usual algebra estimate.  $\square$

We now quantify the size of  $i_0^* \mathcal{W}$  in terms of  $Z$ .

**Lemma 6.2.** *The coefficients  $A_{kj}(\varphi)$  in (6.3) satisfy*

$$\|A_{kj}\|_s^{\text{Lip}(\gamma)} \leq_s \gamma^{-1} (\|Z\|_{s+2\tau+2}^{\text{Lip}(\gamma)} \|\mathfrak{J}_0\|_{s_0+1}^{\text{Lip}(\gamma)} + \|Z\|_{s_0+1}^{\text{Lip}(\gamma)} \|\mathfrak{J}_0\|_{s+2\tau+2}^{\text{Lip}(\gamma)}). \quad (6.6)$$

*Proof.* We estimate the coefficients of the Lie derivative  $L_\omega(i_0^* \mathcal{W}) := \sum_{k < j} \mathcal{D}_\omega A_{kj}(\varphi) d\varphi_k \wedge d\varphi_j$ . Denoting by  $\underline{e}_k$  the  $k$ -th versor of  $\mathbb{R}^\nu$  we have

$$\mathcal{D}_\omega A_{kj} = L_\omega(i_0^* \mathcal{W})(\varphi)[\underline{e}_k, \underline{e}_j] = \mathcal{W}(\partial_\varphi Z(\varphi)\underline{e}_k, \partial_\varphi i_0(\varphi)\underline{e}_j) + \mathcal{W}(\partial_\varphi i_0(\varphi)\underline{e}_k, \partial_\varphi Z(\varphi)\underline{e}_j)$$

(see [6]). Hence

$$\|\mathcal{D}_\omega A_{kj}\|_s^{\text{Lip}(\gamma)} \leq_s \|Z\|_{s+1}^{\text{Lip}(\gamma)} \|\mathfrak{J}_0\|_{s_0+1}^{\text{Lip}(\gamma)} + \|Z\|_{s_0+1}^{\text{Lip}(\gamma)} \|\mathfrak{J}_0\|_{s+1}^{\text{Lip}(\gamma)}. \quad (6.7)$$

The bound (6.6) follows applying  $\mathcal{D}_\omega^{-1}$  and using (6.3), (5.21).  $\square$

As in [6] we first modify the approximate torus  $i_0$  to obtain an isotropic torus  $i_\delta$  which is still approximately invariant. We denote the Laplacian  $\Delta_\varphi := \sum_{k=1}^\nu \partial_{\varphi_k}^2$ .

**Lemma 6.3. (Isotropic torus)** *The torus  $i_\delta(\varphi) := (\theta_0(\varphi), y_\delta(\varphi), z_0(\varphi))$  defined by*

$$y_\delta := y_0 + [\partial_\varphi \theta_0(\varphi)]^{-T} \rho(\varphi), \quad \rho_j(\varphi) := \Delta_\varphi^{-1} \sum_{k=1}^\nu \partial_{\varphi_j} A_{kj}(\varphi) \quad (6.8)$$

*is isotropic. If (6.4) holds, then, for some  $\sigma := \sigma(\nu, \tau)$ ,*

$$\|y_\delta - y_0\|_s^{\text{Lip}(\gamma)} \leq_s \gamma^{-1} (\|Z\|_{s+\sigma}^{\text{Lip}(\gamma)} \|\mathfrak{J}_0\|_{s_0+\sigma}^{\text{Lip}(\gamma)} + \|Z\|_{s_0+\sigma}^{\text{Lip}(\gamma)} \|\mathfrak{J}_0\|_{s+\sigma}^{\text{Lip}(\gamma)}), \quad (6.9)$$

$$\|\mathcal{F}(i_\delta, \zeta_0)\|_s^{\text{Lip}(\gamma)} \leq_s \|Z\|_{s+\sigma}^{\text{Lip}(\gamma)} + \|Z\|_{s_0+\sigma}^{\text{Lip}(\gamma)} \|\mathfrak{J}_0\|_{s+\sigma}^{\text{Lip}(\gamma)} \quad (6.10)$$

$$\|\partial_i [i_\delta][\hat{v}]\|_s \leq_s \|\hat{v}\|_s + \|\mathfrak{J}_0\|_{s+\sigma} \|\hat{v}\|_s. \quad (6.11)$$

In the paper we denote equivalently the differential by  $\partial_i$  or  $d_i$ . Moreover we denote by  $\sigma := \sigma(\nu, \tau)$  possibly different (larger) “loss of derivatives” constants.

*Proof.* In this proof we write  $\|\cdot\|_s$  to denote  $\|\cdot\|_s^{\text{Lip}(\gamma)}$ . The proof of the isotropy of  $i_\delta$  is in [6]. The estimate (6.9) follows by (6.8), (6.6), (6.4) and the tame bound for the inverse  $\|[\partial_\varphi \theta_0]^{-T}\|_s \leq_s 1 + \|\mathfrak{J}_0\|_{s+1}$ . It remains to estimate the difference (see (5.6) and note that  $X_{\mathcal{N}}$  does not depend on  $y$ )

$$\mathcal{F}(i_\delta, \zeta_0) - \mathcal{F}(i_0, \zeta_0) = \begin{pmatrix} 0 \\ \mathcal{D}_\omega(y_\delta - y_0) \\ 0 \end{pmatrix} + X_P(i_\delta) - X_P(i_0). \quad (6.12)$$

Using (5.16), (5.17), we get  $\|\partial_y X_P(i)\|_s \leq_s \varepsilon^{2b} + \varepsilon^{2b-1} \|\mathfrak{J}\|_{s+3}$ . Hence (6.9), (6.4) imply

$$\|X_P(i_\delta) - X_P(i_0)\|_s \leq_s \|\mathfrak{J}_0\|_{s_0+\sigma} \|Z\|_{s+\sigma} + \|\mathfrak{J}_0\|_{s+\sigma} \|Z\|_{s_0+\sigma}. \quad (6.13)$$

Differentiating (6.8) we have

$$\mathcal{D}_\omega(y_\delta - y_0) = [\partial_\varphi \theta_0(\varphi)]^{-T} \mathcal{D}_\omega \rho(\varphi) + (\mathcal{D}_\omega [\partial_\varphi \theta_0(\varphi)]^{-T}) \rho(\varphi) \quad (6.14)$$

and  $\mathcal{D}_\omega \rho_j(\varphi) = \Delta_\varphi^{-1} \sum_{k=1}^\nu \partial_{\varphi_j} \mathcal{D}_\omega A_{kj}(\varphi)$ . Using (6.7), we deduce that

$$\|[\partial_\varphi \theta_0]^{-T} \mathcal{D}_\omega \rho\|_s \leq_s \|Z\|_{s+1} \|\mathfrak{J}_0\|_{s_0+1} + \|Z\|_{s_0+1} \|\mathfrak{J}_0\|_{s+1}. \quad (6.15)$$

To estimate the second term in (6.14), we differentiate  $Z_1(\varphi) = \mathcal{D}_\omega \theta_0(\varphi) - \omega - (\partial_y P)(i_0(\varphi))$  (which is the first component in (5.6)) with respect to  $\varphi$ . We get  $\mathcal{D}_\omega \partial_\varphi \theta_0(\varphi) = \partial_\varphi (\partial_y P)(i_0(\varphi)) + \partial_\varphi Z_1(\varphi)$ . Then, by (5.14),

$$\|\mathcal{D}_\omega [\partial_\varphi \theta_0]^T\|_s \leq_s \varepsilon^4 + \varepsilon^{2b} \|\mathfrak{J}_0\|_{s+2} + \|Z\|_{s+1}. \quad (6.16)$$

Since  $\mathcal{D}_\omega [\partial_\varphi \theta_0(\varphi)]^{-T} = -[\partial_\varphi \theta_0(\varphi)]^{-T} (\mathcal{D}_\omega [\partial_\varphi \theta_0(\varphi)]^T) [\partial_\varphi \theta_0(\varphi)]^{-T}$ , the bounds (6.16), (6.6), (6.4) imply

$$\|(\mathcal{D}_\omega [\partial_\varphi \theta_0]^{-T}) \rho\|_s \leq_s \varepsilon^{6-2b} \gamma^{-1} (\|Z\|_{s+\sigma} \|\mathfrak{J}_0\|_{s_0+\sigma} + \|Z\|_{s_0+\sigma} \|\mathfrak{J}_0\|_{s+\sigma}). \quad (6.17)$$

In conclusion (6.12), (6.13), (6.14), (6.15), (6.17) imply (6.10). The bound (6.11) follows by (6.8), (6.3), (6.2), (6.4).  $\square$

Note that there is no  $\gamma^{-1}$  in the right hand side of (6.10). It turns out that an approximate inverse of  $d_{i,\zeta} \mathcal{F}(i_\delta)$  is an approximate inverse of  $d_{i,\zeta} \mathcal{F}(i_0)$  as well. In order to find an approximate inverse of the linearized operator  $d_{i,\zeta} \mathcal{F}(i_\delta)$  we introduce a suitable set of symplectic coordinates nearby the isotropic torus  $i_\delta$ . We consider the map  $G_\delta : (\psi, \eta, w) \rightarrow (\theta, y, z)$  of the phase space  $\mathbb{T}^\nu \times \mathbb{R}^\nu \times H_{\frac{1}{2}}^\nu$  defined by

$$\begin{pmatrix} \theta \\ y \\ z \end{pmatrix} := G_\delta \begin{pmatrix} \psi \\ \eta \\ w \end{pmatrix} := \begin{pmatrix} \theta_0(\psi) \\ y_\delta(\psi) + [\partial_\psi \theta_0(\psi)]^{-T} \eta + [(\partial_\theta \tilde{z}_0)(\theta_0(\psi))]^T \partial_x^{-1} w \\ z_0(\psi) + w \end{pmatrix} \quad (6.18)$$

where  $\tilde{z}_0(\theta) := z_0(\theta_0^{-1}(\theta))$ . It is proved in [6] that  $G_\delta$  is symplectic, using that the torus  $i_\delta$  is isotropic (Lemma 6.3). In the new coordinates,  $i_\delta$  is the trivial embedded torus  $(\psi, \eta, w) = (\psi, 0, 0)$ . The transformed Hamiltonian  $K := K(\psi, \eta, w, \zeta_0)$  is (recall (5.5))

$$\begin{aligned} K := H_{\varepsilon, \zeta_0} \circ G_\delta &= \theta_0(\psi) \cdot \zeta_0 + K_{00}(\psi) + K_{10}(\psi) \cdot \eta + (K_{01}(\psi), w)_{L^2(\mathbb{T})} + \frac{1}{2} K_{20}(\psi) \eta \cdot \eta \\ &\quad + (K_{11}(\psi) \eta, w)_{L^2(\mathbb{T})} + \frac{1}{2} (K_{02}(\psi) w, w)_{L^2(\mathbb{T})} + K_{\geq 3}(\psi, \eta, w) \end{aligned} \quad (6.19)$$

where  $K_{\geq 3}$  collects the terms at least cubic in the variables  $(\eta, w)$ . At any fixed  $\psi$ , the Taylor coefficient  $K_{00}(\psi) \in \mathbb{R}$ ,  $K_{10}(\psi) \in \mathbb{R}^\nu$ ,  $K_{01}(\psi) \in H_S^\perp$  (it is a function of  $x \in \mathbb{T}$ ),  $K_{20}(\psi)$  is a  $\nu \times \nu$  real matrix,  $K_{02}(\psi)$  is a linear self-adjoint operator of  $H_S^\perp$  and  $K_{11}(\psi) : \mathbb{R}^\nu \rightarrow H_S^\perp$ . Note that the above Taylor coefficients do not depend on the parameter  $\zeta_0$ .

The Hamilton equations associated to (6.19) are

$$\begin{cases} \dot{\psi} = K_{10}(\psi) + K_{20}(\psi) \eta + K_{11}^T(\psi) w + \partial_\eta K_{\geq 3}(\psi, \eta, w) \\ \dot{\eta} = -[\partial_\psi \theta_0(\psi)]^T \zeta_0 - \partial_\psi K_{00}(\psi) - [\partial_\psi K_{10}(\psi)]^T \eta - [\partial_\psi K_{01}(\psi)]^T w \\ \quad - \partial_\psi \left( \frac{1}{2} K_{20}(\psi) \eta \cdot \eta + (K_{11}(\psi) \eta, w)_{L^2(\mathbb{T})} + \frac{1}{2} (K_{02}(\psi) w, w)_{L^2(\mathbb{T})} + K_{\geq 3}(\psi, \eta, w) \right) \\ \dot{w} = \partial_x (K_{01}(\psi) + K_{11}(\psi) \eta + K_{02}(\psi) w) + \nabla_w K_{\geq 3}(\psi, \eta, w) \end{cases} \quad (6.20)$$

where  $[\partial_\psi K_{10}(\psi)]^T$  is the  $\nu \times \nu$  transposed matrix and  $[\partial_\psi K_{01}(\psi)]^T$ ,  $K_{11}^T(\psi) : H_S^\perp \rightarrow \mathbb{R}^\nu$  are defined by the duality relation  $(\partial_\psi K_{01}(\psi)[\hat{\psi}], w)_{L^2} = \hat{\psi} \cdot [\partial_\psi K_{01}(\psi)]^T w$ ,  $\forall \hat{\psi} \in \mathbb{R}^\nu, w \in H_S^\perp$ , and similarly for  $K_{11}$ . Explicitly, for all  $w \in H_S^\perp$ , and denoting  $\underline{e}_k$  the  $k$ -th versor of  $\mathbb{R}^\nu$ ,

$$K_{11}^T(\psi) w = \sum_{k=1}^\nu (K_{11}^T(\psi) w \cdot \underline{e}_k) \underline{e}_k = \sum_{k=1}^\nu (w, K_{11}(\psi) \underline{e}_k)_{L^2(\mathbb{T})} \underline{e}_k \in \mathbb{R}^\nu. \quad (6.21)$$

In the next lemma we estimate the coefficients  $K_{00}$ ,  $K_{10}$ ,  $K_{01}$  in the Taylor expansion (6.19). Note that on an exact solution we have  $Z = 0$  and therefore  $K_{00}(\psi) = \text{const}$ ,  $K_{10} = \omega$  and  $K_{01} = 0$ .

**Lemma 6.4.** *Assume (6.4). Then there is  $\sigma := \sigma(\tau, \nu)$  such that*

$$\|\partial_\psi K_{00}\|_s^{\text{Lip}(\gamma)} + \|K_{10} - \omega\|_s^{\text{Lip}(\gamma)} + \|K_{01}\|_s^{\text{Lip}(\gamma)} \leq_s \|Z\|_{s+\sigma}^{\text{Lip}(\gamma)} + \|Z\|_{s_0+\sigma}^{\text{Lip}(\gamma)} \|\mathfrak{J}_0\|_{s+\sigma}^{\text{Lip}(\gamma)}.$$

*Proof.* Let  $\mathcal{F}(i_\delta, \zeta_0) := Z_\delta := (Z_{1,\delta}, Z_{2,\delta}, Z_{3,\delta})$ . By a direct calculation as in [6] (using (6.19), (5.6))

$$\begin{aligned} \partial_\psi K_{00}(\psi) &= -[\partial_\psi \theta_0(\psi)]^T (-Z_{2,\delta} - [\partial_\psi y_\delta] [\partial_\psi \theta_0]^{-1} Z_{1,\delta} + [(\partial_\theta \tilde{z}_0)(\theta_0(\psi))]^T \partial_x^{-1} Z_{3,\delta} \\ &\quad + [(\partial_\theta \tilde{z}_0)(\theta_0(\psi))]^T \partial_x^{-1} \partial_\psi z_0(\psi) [\partial_\psi \theta_0(\psi)]^{-1} Z_{1,\delta}), \\ K_{10}(\psi) &= \omega - [\partial_\psi \theta_0(\psi)]^{-1} Z_{1,\delta}(\psi), \\ K_{01}(\psi) &= -\partial_x^{-1} Z_{3,\delta} + \partial_x^{-1} \partial_\psi z_0(\psi) [\partial_\psi \theta_0(\psi)]^{-1} Z_{1,\delta}(\psi). \end{aligned}$$

Then (6.4), (6.9), (6.10) (using Lemma 2.4) imply the lemma.  $\square$

**Remark 6.5.** If  $\mathcal{F}(i_0, \zeta_0) = 0$  then  $\zeta_0 = 0$  by Lemma 6.1, and Lemma 6.4 implies that (6.19) simplifies to  $K = \text{const} + \omega \cdot \eta + \frac{1}{2} K_{20}(\psi) \eta \cdot \eta + (K_{11}(\psi) \eta, w)_{L^2(\mathbb{T})} + \frac{1}{2} (K_{02}(\psi) w, w)_{L^2(\mathbb{T})} + K_{\geq 3}$ .  $\square$

We now estimate  $K_{20}$ ,  $K_{11}$  in (6.19). The norm of  $K_{20}$  is the sum of the norms of its matrix entries.

**Lemma 6.6.** *Assume (6.4). Then*

$$\|K_{20} + 3\varepsilon^{2b} I\|_s^{\text{Lip}(\gamma)} \leq_s \varepsilon^{2b+2} + \varepsilon^{2b} \|\mathfrak{J}_0\|_{s+\sigma}^{\text{Lip}(\gamma)} + \varepsilon^3 \gamma^{-1} \|\mathfrak{J}_0\|_{s_0+\sigma}^{\text{Lip}(\gamma)} \|Z\|_{s+\sigma}^{\text{Lip}(\gamma)} \quad (6.22)$$

$$\|K_{11} \eta\|_s^{\text{Lip}(\gamma)} \leq_s \varepsilon^5 \gamma^{-1} \|\eta\|_s^{\text{Lip}(\gamma)} + \varepsilon^{2b-1} (\|\mathfrak{J}_0\|_{s+\sigma}^{\text{Lip}(\gamma)} + \gamma^{-1} \|\mathfrak{J}_0\|_{s_0+\sigma}^{\text{Lip}(\gamma)} \|Z\|_{s+\sigma}^{\text{Lip}(\gamma)}) \|\eta\|_{s_0}^{\text{Lip}(\gamma)} \quad (6.23)$$

$$\|K_{11}^T w\|_s^{\text{Lip}(\gamma)} \leq_s \varepsilon^5 \gamma^{-1} \|w\|_{s+2}^{\text{Lip}(\gamma)} + \varepsilon^{2b-1} (\|\mathfrak{J}_0\|_{s+\sigma}^{\text{Lip}(\gamma)} + \gamma^{-1} \|\mathfrak{J}_0\|_{s_0+\sigma}^{\text{Lip}(\gamma)} \|Z\|_{s+\sigma}^{\text{Lip}(\gamma)}) \|w\|_{s_0+2}^{\text{Lip}(\gamma)}. \quad (6.24)$$

In particular  $\|K_{20} + 3\varepsilon^{2b} I\|_{s_0}^{\text{Lip}(\gamma)} \leq C \varepsilon^6 \gamma^{-1}$ , and

$$\|K_{11} \eta\|_{s_0}^{\text{Lip}(\gamma)} \leq C \varepsilon^5 \gamma^{-1} \|\eta\|_{s_0}^{\text{Lip}(\gamma)}, \quad \|K_{11}^T w\|_{s_0}^{\text{Lip}(\gamma)} \leq C \varepsilon^5 \gamma^{-1} \|w\|_{s_0}^{\text{Lip}(\gamma)}.$$

*Proof.* To shorten the notation, in this proof we write  $\|\cdot\|_s$  for  $\|\cdot\|_s^{\text{Lip}(\gamma)}$ . We have

$$K_{20}(\varphi) = [\partial_\varphi \theta_0(\varphi)]^{-1} \partial_{yy} H_\varepsilon(i_\delta(\varphi)) [\partial_\varphi \theta_0(\varphi)]^{-T} = [\partial_\varphi \theta_0(\varphi)]^{-1} \partial_{yy} P(i_\delta(\varphi)) [\partial_\varphi \theta_0(\varphi)]^{-T}.$$

Then (5.17), (6.4), (6.9) imply (6.22). Now (see also [6])

$$K_{11}(\varphi) = \partial_y \nabla_z H_\varepsilon(i_\delta(\varphi)) [\partial_\varphi \theta_0(\varphi)]^{-T} - \partial_x^{-1} (\partial_\theta \tilde{z}_0)(\theta_0(\varphi)) (\partial_{yy} H_\varepsilon)(i_\delta(\varphi)) [\partial_\varphi \theta_0(\varphi)]^{-T} \\ \stackrel{(4.11)}{=} \partial_y \nabla_z P(i_\delta(\varphi)) [\partial_\varphi \theta_0(\varphi)]^{-T} - \partial_x^{-1} (\partial_\theta \tilde{z}_0)(\theta_0(\varphi)) (\partial_{yy} P)(i_\delta(\varphi)) [\partial_\varphi \theta_0(\varphi)]^{-T},$$

therefore, using (5.16), (5.17), (6.4), we deduce (6.23). The bound (6.24) for  $K_{11}^T$  follows by (6.21).  $\square$

Under the linear change of variables

$$DG_\delta(\varphi, 0, 0) \begin{pmatrix} \hat{\psi} \\ \hat{\eta} \\ \hat{w} \end{pmatrix} := \begin{pmatrix} \partial_\psi \theta_0(\varphi) & 0 & 0 \\ \partial_\psi y_\delta(\varphi) & [\partial_\psi \theta_0(\varphi)]^{-T} & -[(\partial_\theta \tilde{z}_0)(\theta_0(\varphi))]^T \partial_x^{-1} \\ \partial_\psi z_0(\varphi) & 0 & I \end{pmatrix} \begin{pmatrix} \hat{\psi} \\ \hat{\eta} \\ \hat{w} \end{pmatrix} \quad (6.25)$$

the linearized operator  $d_{i,\zeta} \mathcal{F}(i_\delta)$  transforms (approximately, see (6.45)) into the operator obtained linearizing (6.20) at  $(\psi, \eta, w, \zeta) = (\varphi, 0, 0, \zeta_0)$  (with  $\partial_t \rightsquigarrow \mathcal{D}_\omega$ ), namely

$$\begin{pmatrix} \mathcal{D}_\omega \hat{\psi} - \partial_\psi K_{10}(\varphi) [\hat{\psi}] - K_{20}(\varphi) \hat{\eta} - K_{11}^T(\varphi) \hat{w} \\ \mathcal{D}_\omega \hat{\eta} + [\partial_\psi \theta_0(\varphi)]^T \hat{\zeta} + \partial_\psi [\partial_\psi \theta_0(\varphi)]^T [\hat{\psi}, \zeta_0] + \partial_{\psi\psi} K_{00}(\varphi) [\hat{\psi}] + [\partial_\psi K_{10}(\varphi)]^T \hat{\eta} + [\partial_\psi K_{01}(\varphi)]^T \hat{w} \\ \mathcal{D}_\omega \hat{w} - \partial_x \{ \partial_\psi K_{01}(\varphi) [\hat{\psi}] + K_{11}(\varphi) \hat{\eta} + K_{02}(\varphi) \hat{w} \} \end{pmatrix}. \quad (6.26)$$

We now estimate the induced composition operator.

**Lemma 6.7.** *Assume (6.4) and let  $\hat{v} := (\hat{\psi}, \hat{\eta}, \hat{w})$ . Then*

$$\|DG_\delta(\varphi, 0, 0) [\hat{v}]\|_s + \|DG_\delta(\varphi, 0, 0)^{-1} [\hat{v}]\|_s \leq_s \|\hat{v}\|_s + (\|\mathcal{J}_0\|_{s+\sigma} + \gamma^{-1} \|\mathcal{J}_0\|_{s_0+\sigma} \|Z\|_{s+\sigma}) \|\hat{v}\|_{s_0}, \quad (6.27) \\ \|D^2 G_\delta(\varphi, 0, 0) [\hat{v}_1, \hat{v}_2]\|_s \leq_s \|\hat{v}_1\|_s \|\hat{v}_2\|_{s_0} + \|\hat{v}_1\|_{s_0} \|\hat{v}_2\|_s + (\|\mathcal{J}_0\|_{s+\sigma} + \gamma^{-1} \|\mathcal{J}_0\|_{s_0+\sigma} \|Z\|_{s+\sigma}) \|\hat{v}_1\|_{s_0} \|\hat{v}_2\|_{s_0}$$

for some  $\sigma := \sigma(\nu, \tau)$ . Moreover the same estimates hold if we replace the norm  $\|\cdot\|_s$  with  $\|\cdot\|_s^{\text{Lip}(\gamma)}$ .

*Proof.* The estimate (6.27) for  $DG_\delta(\varphi, 0, 0)$  follows by (6.25) and (6.9). By (6.4),  $\|(DG_\delta(\varphi, 0, 0) - I) \hat{v}\|_{s_0} \leq C \varepsilon^{6-2b} \gamma^{-1} \|\hat{v}\|_{s_0} \leq \|\hat{v}\|_{s_0}/2$ . Therefore  $DG_\delta(\varphi, 0, 0)$  is invertible and, by Neumann series, the inverse satisfies (6.27). The bound for  $D^2 G_\delta$  follows by differentiating  $DG_\delta$ .  $\square$

In order to construct an approximate inverse of (6.26) it is sufficient to solve the equation

$$\mathbb{D}[\hat{\psi}, \hat{\eta}, \hat{w}, \hat{\zeta}] := \begin{pmatrix} \mathcal{D}_\omega \hat{\psi} - K_{20}(\varphi) \hat{\eta} - K_{11}^T(\varphi) \hat{w} \\ \mathcal{D}_\omega \hat{\eta} + [\partial_\psi \theta_0(\varphi)]^T \hat{\zeta} \\ \mathcal{D}_\omega \hat{w} - \partial_x K_{11}(\varphi) \hat{\eta} - \partial_x K_{02}(\varphi) \hat{w} \end{pmatrix} = \begin{pmatrix} g_1 \\ g_2 \\ g_3 \end{pmatrix} \quad (6.28)$$

which is obtained by neglecting in (6.26) the terms  $\partial_\psi K_{10}$ ,  $\partial_{\psi\psi} K_{00}$ ,  $\partial_\psi K_{00}$ ,  $\partial_\psi K_{01}$  and  $\partial_\psi [\partial_\psi \theta_0(\varphi)]^T [\cdot, \zeta_0]$  (which are naught at a solution by Lemmata 6.4 and 6.1).

First we solve the second equation in (6.28), namely  $\mathcal{D}_\omega \hat{\eta} = g_2 - [\partial_\psi \theta_0(\varphi)]^T \hat{\zeta}$ . We choose  $\hat{\zeta}$  so that the  $\varphi$ -average of the right hand side is zero, namely

$$\hat{\zeta} = \langle g_2 \rangle \quad (6.29)$$

(we denote  $\langle g \rangle := (2\pi)^{-\nu} \int_{\mathbb{T}^\nu} g(\varphi) d\varphi$ ). Note that the  $\varphi$ -averaged matrix  $\langle [\partial_\psi \theta_0]^T \rangle = \langle I + [\partial_\psi \Theta_0]^T \rangle = I$  because  $\theta_0(\varphi) = \varphi + \Theta_0(\varphi)$  and  $\Theta_0(\varphi)$  is a periodic function. Therefore

$$\hat{\eta} := \mathcal{D}_\omega^{-1} (g_2 - [\partial_\psi \theta_0(\varphi)]^T \langle g_2 \rangle) + \langle \hat{\eta} \rangle, \quad \langle \hat{\eta} \rangle \in \mathbb{R}^\nu, \quad (6.30)$$

where the average  $\langle \hat{\eta} \rangle$  will be fixed below. Then we consider the third equation

$$\mathcal{L}_\omega \hat{w} = g_3 + \partial_x K_{11}(\varphi) \hat{\eta}, \quad \mathcal{L}_\omega := \omega \cdot \partial_\varphi - \partial_x K_{02}(\varphi). \quad (6.31)$$



- **INVERSION ASSUMPTION.** *There exists a set  $\Omega_\infty \subset \Omega_o$  such that for all  $\omega \in \Omega_\infty$ , for every function  $g \in H_{S^\perp}^{s+\mu}(\mathbb{T}^{\nu+1})$  there exists a solution  $h := \mathcal{L}_\omega^{-1}g \in H_{S^\perp}^s(\mathbb{T}^{\nu+1})$  of the linear equation  $\mathcal{L}_\omega h = g$  which satisfies*

$$\|\mathcal{L}_\omega^{-1}g\|_s^{\text{Lip}(\gamma)} \leq C(s)\gamma^{-1}(\|g\|_{s+\mu}^{\text{Lip}(\gamma)} + \varepsilon\gamma^{-1}\{\|\mathfrak{J}_0\|_{s+\mu}^{\text{Lip}(\gamma)} + \gamma^{-1}\|\mathfrak{J}_0\|_{s_0+\mu}^{\text{Lip}(\gamma)}\|Z\|_{s+\mu}^{\text{Lip}(\gamma)}\})\|g\|_{s_0}^{\text{Lip}(\gamma)} \quad (6.32)$$

for some  $\mu := \mu(\tau, \nu) > 0$ .

**Remark 6.8.** The term  $\varepsilon\gamma^{-1}\{\|\mathfrak{J}_0\|_{s+\mu}^{\text{Lip}(\gamma)} + \gamma^{-1}\|\mathfrak{J}_0\|_{s_0+\mu}^{\text{Lip}(\gamma)}\|Z\|_{s+\mu}^{\text{Lip}(\gamma)}\}$  arises because the remainder  $R_6$  in section 8.6 contains the term  $\varepsilon(\|\Theta_0\|_{s+\mu}^{\text{Lip}(\gamma)} + \|y_\delta\|_{s+\mu}^{\text{Lip}(\gamma)}) \leq \varepsilon\|\mathfrak{J}_\delta\|_{s+\mu}^{\text{Lip}(\gamma)}$ , see Lemma 8.24.  $\square$

By the above assumption there exists a solution

$$\widehat{w} := \mathcal{L}_\omega^{-1}[g_3 + \partial_x K_{11}(\varphi)\widehat{\eta}] \quad (6.33)$$

of (6.31). Finally, we solve the first equation in (6.28), which, substituting (6.30), (6.33), becomes

$$\mathcal{D}_\omega \widehat{\psi} = g_1 + M_1(\varphi)\langle \widehat{\eta} \rangle + M_2(\varphi)g_2 + M_3(\varphi)g_3 - M_2(\varphi)[\partial_\psi \theta_0]^T \langle g_2 \rangle, \quad (6.34)$$

where

$$M_1(\varphi) := K_{20}(\varphi) + K_{11}^T(\varphi)\mathcal{L}_\omega^{-1}\partial_x K_{11}(\varphi), \quad M_2(\varphi) := M_1(\varphi)\mathcal{D}_\omega^{-1}, \quad M_3(\varphi) := K_{11}^T(\varphi)\mathcal{L}_\omega^{-1}. \quad (6.35)$$

In order to solve the equation (6.34) we have to choose  $\langle \widehat{\eta} \rangle$  such that the right hand side in (6.34) has zero average. By Lemma 6.6 and (6.4), the  $\varphi$ -averaged matrix  $\langle M_1 \rangle = -3\varepsilon^{2b}I + O(\varepsilon^{10}\gamma^{-3})$ . Therefore, for  $\varepsilon$  small,  $\langle M_1 \rangle$  is invertible and  $\langle M_1 \rangle^{-1} = O(\varepsilon^{-2b}) = O(\gamma^{-1})$  (recall (5.10)). Thus we define

$$\langle \widehat{\eta} \rangle := -\langle M_1 \rangle^{-1}[\langle g_1 \rangle + \langle M_2 g_2 \rangle + \langle M_3 g_3 \rangle - \langle M_2 [\partial_\psi \theta_0]^T \rangle \langle g_2 \rangle]. \quad (6.36)$$

With this choice of  $\langle \widehat{\eta} \rangle$  the equation (6.34) has the solution

$$\widehat{\psi} := \mathcal{D}_\omega^{-1}[g_1 + M_1(\varphi)\langle \widehat{\eta} \rangle + M_2(\varphi)g_2 + M_3(\varphi)g_3 - M_2(\varphi)[\partial_\psi \theta_0]^T \langle g_2 \rangle]. \quad (6.37)$$

In conclusion, we have constructed a solution  $(\widehat{\psi}, \widehat{\eta}, \widehat{w}, \widehat{\zeta})$  of the linear system (6.28).

**Proposition 6.9.** *Assume (6.4) and (6.32). Then,  $\forall \omega \in \Omega_\infty$ ,  $\forall g := (g_1, g_2, g_3)$ , the system (6.28) has a solution  $\mathbb{D}^{-1}g := (\widehat{\psi}, \widehat{\eta}, \widehat{w}, \widehat{\zeta})$  where  $(\widehat{\psi}, \widehat{\eta}, \widehat{w}, \widehat{\zeta})$  are defined in (6.37), (6.30), (6.36), (6.33), (6.29) satisfying*

$$\|\mathbb{D}^{-1}g\|_s^{\text{Lip}(\gamma)} \leq_s \gamma^{-1}(\|g\|_{s+\mu}^{\text{Lip}(\gamma)} + \varepsilon\gamma^{-1}\{\|\mathfrak{J}_0\|_{s+\mu}^{\text{Lip}(\gamma)} + \gamma^{-1}\|\mathfrak{J}_0\|_{s_0+\mu}^{\text{Lip}(\gamma)}\|\mathcal{F}(i_0, \zeta_0)\|_{s+\mu}^{\text{Lip}(\gamma)}\})\|g\|_{s_0+\mu}^{\text{Lip}(\gamma)}. \quad (6.38)$$

*Proof.* Recalling (6.35), by Lemma 6.6, (6.32), (6.4) we get  $\|M_2 h\|_{s_0} + \|M_3 h\|_{s_0} \leq C\|h\|_{s_0+\sigma}$ . Then, by (6.36) and  $\langle M_1 \rangle^{-1} = O(\varepsilon^{-2b}) = O(\gamma^{-1})$ , we deduce  $\|\langle \widehat{\eta} \rangle\|_s^{\text{Lip}(\gamma)} \leq C\gamma^{-1}\|g\|_{s_0+\sigma}^{\text{Lip}(\gamma)}$  and (6.30), (5.21) imply  $\|\widehat{\eta}\|_s^{\text{Lip}(\gamma)} \leq_s \gamma^{-1}(\|g\|_{s+\sigma}^{\text{Lip}(\gamma)} + \|\mathfrak{J}_0\|_{s+\sigma}\|g\|_{s_0}^{\text{Lip}(\gamma)})$ . The bound (6.38) is sharp for  $\widehat{w}$  because  $\mathcal{L}_\omega^{-1}g_3$  in (6.33) is estimated using (6.32). Finally  $\widehat{\psi}$  satisfies (6.38) using (6.37), (6.35), (6.32), (5.21) and Lemma 6.6.  $\square$

Finally we prove that the operator

$$\mathbf{T}_0 := (D\widetilde{G}_\delta)(\varphi, 0, 0) \circ \mathbb{D}^{-1} \circ (DG_\delta)(\varphi, 0, 0)^{-1} \quad (6.39)$$

is an approximate right inverse for  $d_{i,\zeta}\mathcal{F}(i_0)$  where  $\widetilde{G}_\delta(\psi, \eta, w, \zeta) := (G_\delta(\psi, \eta, w), \zeta)$  is the identity on the  $\zeta$ -component. We denote the norm  $\|(\psi, \eta, w, \zeta)\|_s^{\text{Lip}(\gamma)} := \max\{\|(\psi, \eta, w)\|_s^{\text{Lip}(\gamma)}, |\zeta|^{\text{Lip}(\gamma)}\}$ .

**Theorem 6.10. (Approximate inverse)** *Assume (6.4) and the inversion assumption (6.32). Then there exists  $\mu := \mu(\tau, \nu) > 0$  such that, for all  $\omega \in \Omega_\infty$ , for all  $g := (g_1, g_2, g_3)$ , the operator  $\mathbf{T}_0$  defined in (6.39) satisfies*

$$\|\mathbf{T}_0 g\|_s^{\text{Lip}(\gamma)} \leq_s \gamma^{-1} (\|g\|_{s+\mu}^{\text{Lip}(\gamma)} + \varepsilon \gamma^{-1} \{\|\mathfrak{J}_0\|_{s+\mu}^{\text{Lip}(\gamma)} + \gamma^{-1} \|\mathfrak{J}_0\|_{s_0+\mu}^{\text{Lip}(\gamma)} \|\mathcal{F}(i_0, \zeta_0)\|_{s+\mu}^{\text{Lip}(\gamma)}\} \|g\|_{s_0+\mu}^{\text{Lip}(\gamma)}). \quad (6.40)$$

It is an approximate inverse of  $d_{i,\zeta} \mathcal{F}(i_0)$ , namely

$$\begin{aligned} & \| (d_{i,\zeta} \mathcal{F}(i_0) \circ \mathbf{T}_0 - I) g \|_s^{\text{Lip}(\gamma)} \\ & \leq_s \gamma^{-1} \left( \|\mathcal{F}(i_0, \zeta_0)\|_{s_0+\mu}^{\text{Lip}(\gamma)} \|g\|_{s+\mu}^{\text{Lip}(\gamma)} + \{ \|\mathcal{F}(i_0, \zeta_0)\|_{s+\mu}^{\text{Lip}(\gamma)} + \varepsilon \gamma^{-1} \|\mathcal{F}(i_0, \zeta_0)\|_{s_0+\mu}^{\text{Lip}(\gamma)} \|\mathfrak{J}_0\|_{s+\mu}^{\text{Lip}(\gamma)} \} \|g\|_{s_0+\mu}^{\text{Lip}(\gamma)} \right). \end{aligned} \quad (6.41)$$

*Proof.* We denote  $\|\cdot\|_s$  instead of  $\|\cdot\|_s^{\text{Lip}(\gamma)}$ . The bound (6.40) follows from (6.39), (6.38), (6.27). By (5.6), since  $X_{\mathcal{N}}$  does not depend on  $y$ , and  $i_\delta$  differs from  $i_0$  only for the  $y$  component, we have

$$\begin{aligned} d_{i,\zeta} \mathcal{F}(i_0)[\widehat{v}, \widehat{\zeta}] - d_{i,\zeta} \mathcal{F}(i_\delta)[\widehat{v}, \widehat{\zeta}] &= d_i X_P(i_\delta)[\widehat{v}] - d_i X_P(i_0)[\widehat{v}] \\ &= \int_0^1 \partial_y d_i X_P(\theta_0, y_0 + s(y_\delta - y_0), z_0)[y_\delta - y_0, \widehat{v}] ds =: \mathcal{E}_0[\widehat{v}, \widehat{\zeta}]. \end{aligned} \quad (6.42)$$

By (5.18), (6.9), (6.4), we estimate

$$\|\mathcal{E}_0[\widehat{v}, \widehat{\zeta}]\|_s \leq_s \|Z\|_{s_0+\sigma} \|\widehat{v}\|_{s+\sigma} + \|Z\|_{s+\sigma} \|\widehat{v}\|_{s_0+\sigma} + \varepsilon^{2b-1} \gamma^{-1} \|Z\|_{s_0+\sigma} \|\widehat{v}\|_{s_0+\sigma} \|\mathfrak{J}_0\|_{s+\sigma} \quad (6.43)$$

where  $Z := \mathcal{F}(i_0, \zeta_0)$  (recall (6.5)). Note that  $\mathcal{E}_0[\widehat{v}, \widehat{\zeta}]$  is, in fact, independent of  $\widehat{\zeta}$ . Denote the set of variables  $(\psi, \eta, w) =: \mathbf{u}$ . Under the transformation  $G_\delta$ , the nonlinear operator  $\mathcal{F}$  in (5.6) transforms into

$$\mathcal{F}(G_\delta(\mathbf{u}(\varphi)), \zeta) = DG_\delta(\mathbf{u}(\varphi)) (\mathcal{D}_\omega \mathbf{u}(\varphi) - X_K(\mathbf{u}(\varphi), \zeta)), \quad K = H_{\varepsilon, \zeta} \circ G_\delta, \quad (6.44)$$

see (6.20). Differentiating (6.44) at the trivial torus  $\mathbf{u}_\delta(\varphi) = G_\delta^{-1}(i_\delta)(\varphi) = (\varphi, 0, 0)$ , at  $\zeta = \zeta_0$ , in the directions  $(\widehat{\mathbf{u}}, \widehat{\zeta}) = (DG_\delta(\mathbf{u}_\delta)^{-1}[\widehat{v}], \widehat{\zeta}) = D\widetilde{G}_\delta(\mathbf{u}_\delta)^{-1}[\widehat{v}, \widehat{\zeta}]$ , we get

$$d_{i,\zeta} \mathcal{F}(i_\delta)[\widehat{v}, \widehat{\zeta}] = DG_\delta(\mathbf{u}_\delta) (\mathcal{D}_\omega \widehat{\mathbf{u}} - d_{\mathbf{u}, \zeta} X_K(\mathbf{u}_\delta, \zeta_0)[\widehat{\mathbf{u}}, \widehat{\zeta}]) + \mathcal{E}_1[\widehat{v}, \widehat{\zeta}], \quad (6.45)$$

$$\mathcal{E}_1[\widehat{v}, \widehat{\zeta}] := D^2 G_\delta(\mathbf{u}_\delta) [DG_\delta(\mathbf{u}_\delta)^{-1} \mathcal{F}(i_\delta, \zeta_0), DG_\delta(\mathbf{u}_\delta)^{-1}[\widehat{v}]], \quad (6.46)$$

where  $d_{\mathbf{u}, \zeta} X_K(\mathbf{u}_\delta, \zeta_0)$  is expanded in (6.26). In fact,  $\mathcal{E}_1$  is independent of  $\widehat{\zeta}$ . We split

$$\mathcal{D}_\omega \widehat{\mathbf{u}} - d_{\mathbf{u}, \zeta} X_K(\mathbf{u}_\delta, \zeta_0)[\widehat{\mathbf{u}}, \widehat{\zeta}] = \mathbb{D}[\widehat{\mathbf{u}}, \widehat{\zeta}] + R_Z[\widehat{\mathbf{u}}, \widehat{\zeta}],$$

where  $\mathbb{D}[\widehat{\mathbf{u}}, \widehat{\zeta}]$  is defined in (6.28) and

$$R_Z[\widehat{\psi}, \widehat{\eta}, \widehat{w}, \widehat{\zeta}] := \begin{pmatrix} -\partial_\psi K_{10}(\varphi)[\widehat{\psi}] \\ \partial_\psi [\partial_\psi \theta_0(\varphi)]^T [\widehat{\psi}, \zeta_0] + \partial_{\psi\psi} K_{00}(\varphi)[\widehat{\psi}] + [\partial_\psi K_{10}(\varphi)]^T \widehat{\eta} + [\partial_\psi K_{01}(\varphi)]^T \widehat{w} \\ -\partial_x \{ \partial_\psi K_{01}(\varphi)[\widehat{\psi}] \} \end{pmatrix} \quad (6.47)$$

( $R_Z$  is independent of  $\widehat{\zeta}$ ). By (6.42) and (6.45),

$$d_{i,\zeta} \mathcal{F}(i_0) = DG_\delta(\mathbf{u}_\delta) \circ \mathbb{D} \circ D\widetilde{G}_\delta(\mathbf{u}_\delta)^{-1} + \mathcal{E}_0 + \mathcal{E}_1 + \mathcal{E}_2, \quad \mathcal{E}_2 := DG_\delta(\mathbf{u}_\delta) \circ R_Z \circ D\widetilde{G}_\delta(\mathbf{u}_\delta)^{-1}. \quad (6.48)$$

By Lemmata 6.4, 6.7, 6.1, and (6.10), (6.4), the terms  $\mathcal{E}_1, \mathcal{E}_2$  (see (6.46), (6.48), (6.47)) satisfy the same bound (6.43) as  $\mathcal{E}_0$  (in fact even better). Thus the sum  $\mathcal{E} := \mathcal{E}_0 + \mathcal{E}_1 + \mathcal{E}_2$  satisfies (6.43). Applying  $\mathbf{T}_0$  defined in (6.39) to the right in (6.48), since  $\mathbb{D} \circ \mathbb{D}^{-1} = I$  (see Proposition 6.9), we get  $d_{i,\zeta} \mathcal{F}(i_0) \circ \mathbf{T}_0 - I = \mathcal{E} \circ \mathbf{T}_0$ . Then (6.41) follows from (6.40) and the bound (6.43) for  $\mathcal{E}$ .  $\square$

## 7 The linearized operator in the normal directions

The goal of this section is to write an explicit expression of the linearized operator  $\mathcal{L}_\omega$  defined in (6.31), see Proposition 7.6. To this aim, we compute  $\frac{1}{2}(K_{02}(\psi)w, w)_{L^2(\mathbb{T})}$ ,  $w \in H_S^\perp$ , which collects all the components of  $(H_\varepsilon \circ G_\delta)(\psi, 0, w)$  that are quadratic in  $w$ , see (6.19).

We first prove some preliminary lemmata.

**Lemma 7.1.** *Let  $H$  be a Hamiltonian of class  $C^2(H_0^1(\mathbb{T}_x), \mathbb{R})$  and consider a map  $\Phi(u) := u + \Psi(u)$  satisfying  $\Psi(u) = \Pi_E \Psi(\Pi_E u)$ , for all  $u$ , where  $E$  is a finite dimensional subspace as in (3.3). Then*

$$\partial_u [\nabla(H \circ \Phi)](u)[h] = (\partial_u \nabla H)(\Phi(u))[h] + \mathcal{R}(u)[h], \quad (7.1)$$

where  $\mathcal{R}(u)$  has the “finite dimensional” form

$$\mathcal{R}(u)[h] = \sum_{|j| \leq C} (h, g_j(u))_{L^2(\mathbb{T})} \chi_j(u) \quad (7.2)$$

with  $\chi_j(u) = e^{ijx}$  or  $g_j(u) = e^{ijx}$ . The remainder  $\mathcal{R}(u) = \mathcal{R}_0(u) + \mathcal{R}_1(u) + \mathcal{R}_2(u)$  with

$$\begin{aligned} \mathcal{R}_0(u) &:= (\partial_u \nabla H)(\Phi(u)) \partial_u \Psi(u), & \mathcal{R}_1(u) &:= [\partial_u \{\Psi'(u)^T\}][\cdot, \nabla H(\Phi(u))], \\ \mathcal{R}_2(u) &:= [\partial_u \Psi(u)]^T (\partial_u \nabla H)(\Phi(u)) \partial_u \Phi(u). \end{aligned} \quad (7.3)$$

*Proof.* By a direct calculation,

$$\nabla(H \circ \Phi)(u) = [\Phi'(u)]^T \nabla H(\Phi(u)) = \nabla H(\Phi(u)) + [\Psi'(u)]^T \nabla H(\Phi(u)) \quad (7.4)$$

where  $\Phi'(u) := (\partial_u \Phi)(u)$  and  $[\cdot]^T$  denotes the transpose with respect to the  $L^2$  scalar product. Differentiating (7.4), we get (7.1) and (7.3).

Let us show that each  $\mathcal{R}_m$  has the form (7.2). We have

$$\Psi'(u) = \Pi_E \Psi'(\Pi_E u) \Pi_E, \quad [\Psi'(u)]^T = \Pi_E [\Psi'(\Pi_E u)]^T \Pi_E. \quad (7.5)$$

Hence, setting  $A := (\partial_u \nabla H)(\Phi(u)) \Pi_E \Psi'(\Pi_E u)$ , we get

$$\mathcal{R}_0(u)[h] = A[\Pi_E h] = \sum_{|j| \leq C} h_j A(e^{ijx}) = \sum_{|j| \leq C} (h, g_j)_{L^2(\mathbb{T})} \chi_j$$

with  $g_j := e^{ijx}$ ,  $\chi_j := A(e^{ijx})$ . Similarly, using (7.5), and setting  $A := [\Psi'(\Pi_E u)]^T \Pi_E (\partial_u \nabla H)(\Phi(u)) \Phi'(u)$ , we get

$$\mathcal{R}_2(u)[h] = \Pi_E [Ah] = \sum_{|j| \leq C} (Ah, e^{ijx})_{L^2(\mathbb{T})} e^{ijx} = \sum_{|j| \leq C} (h, A^T e^{ijx})_{L^2(\mathbb{T})} e^{ijx},$$

which has the form (7.2) with  $g_j := A^T(e^{ijx})$  and  $\chi_j := e^{ijx}$ . Differentiating the second equality in (7.5), we see that

$$\mathcal{R}_1(u)[h] = \Pi_E [Ah], \quad Ah := \partial_u \{\Psi'(\Pi_E u)^T\} [\Pi_E h, \Pi_E (\nabla H)(\Phi(u))],$$

which has the same form of  $\mathcal{R}_2$  and so (7.2).  $\square$

**Lemma 7.2.** *Let  $H(u) := \int_{\mathbb{T}} f(u) X(u) dx$  where  $X(u) = \Pi_E X(\Pi_E u)$  and  $f(u)(x) := f(u(x))$  is the composition operator for a function of class  $C^2$ . Then*

$$(\partial_u \nabla H)(u)[h] = f''(u) X(u) h + \mathcal{R}(u)[h] \quad (7.6)$$

where  $\mathcal{R}(u)$  has the form (7.2) with  $\chi_j(u) = e^{ijx}$  or  $g_j(u) = e^{ijx}$ .

*Proof.* A direct calculation proves that  $\nabla H(u) = f'(u) X(u) + X'(u)^T [f(u)]$ , and (7.6) follows with  $\mathcal{R}(u)[h] = f'(u) X'(u)[h] + \partial_u \{X'(u)^T\} [h, f(u)] + X'(u)^T [f'(u) h]$ , which has the form (7.2).  $\square$

We conclude this section with a technical lemma used from the end of section 8.3 about the decay norms of “finite dimensional operators”. Note that operators of the form (7.7) (that will appear in section 8.1) reduce to those in (7.2) when the functions  $g_j(\tau)$ ,  $\chi_j(\tau)$  are independent of  $\tau$

**Lemma 7.3.** *Let  $\mathcal{R}$  be an operator of the form*

$$\mathcal{R}h = \sum_{|j| \leq C} \int_0^1 (h, g_j(\tau))_{L^2(\mathbb{T})} \chi_j(\tau) d\tau, \quad (7.7)$$

where the functions  $g_j(\tau), \chi_j(\tau) \in H^s, \tau \in [0, 1]$  depend in a Lipschitz way on the parameter  $\omega$ . Then its matrix  $s$ -decay norm (see (2.16)-(2.17)) satisfies

$$|\mathcal{R}|_s^{\text{Lip}(\gamma)} \leq_s \sum_{|j| \leq C} \sup_{\tau \in [0, 1]} \{ \|\chi_j(\tau)\|_s^{\text{Lip}(\gamma)} \|g_j(\tau)\|_{s_0}^{\text{Lip}(\gamma)} + \|\chi_j(\tau)\|_{s_0}^{\text{Lip}(\gamma)} \|g_j(\tau)\|_s^{\text{Lip}(\gamma)} \}.$$

*Proof.* For each  $\tau \in [0, 1]$ , the operator  $h \mapsto (h, g_j(\tau)) \chi_j(\tau)$  is the composition  $\chi_j(\tau) \circ \Pi_0 \circ g_j(\tau)$  of the multiplication operators for  $g_j(\tau), \chi_j(\tau)$  and  $h \mapsto \Pi_0 h := \int_{\mathbb{T}} h dx$ . Hence the lemma follows by the interpolation estimate (2.20) and (2.18).  $\square$

## 7.1 Composition with the map $G_\delta$

In the sequel we shall use that  $\mathfrak{I}_\delta := \mathfrak{I}_\delta(\varphi; \omega) := i_\delta(\varphi; \omega) - (\varphi, 0, 0)$  satisfies, by Lemma 6.3 and (6.4),

$$\|\mathfrak{I}_\delta\|_{s_0 + \mu}^{\text{Lip}(\gamma)} \leq C \varepsilon^{6-2b} \gamma^{-1}. \quad (7.8)$$

We now study the Hamiltonian  $K := H_\varepsilon \circ G_\delta = \varepsilon^{-2b} \mathcal{H} \circ A_\varepsilon \circ G_\delta$  defined in (6.19), (4.6).

Recalling (4.7) and (6.18) the map  $A_\varepsilon \circ G_\delta$  has the form

$$A_\varepsilon \circ G_\delta(\psi, \eta, w) = \varepsilon \sum_{j \in S} \sqrt{\xi_j + \varepsilon^{2(b-1)} |j| [y_\delta(\psi) + L_1(\psi)\eta + L_2(\psi)w]_j} e^{i[\theta_0(\psi)]_j} e^{ijx} + \varepsilon^b (z_0(\psi) + w) \quad (7.9)$$

where

$$L_1(\psi) := [\partial_\psi \theta_0(\psi)]^{-T}, \quad L_2(\psi) := [(\partial_\theta \tilde{z}_0)(\theta_0(\psi))]^T \partial_x^{-1}. \quad (7.10)$$

By Taylor's formula, we develop (7.9) in  $w$  at  $\eta = 0, w = 0$ , and we get  $A_\varepsilon \circ G_\delta(\psi, 0, w) = T_\delta(\psi) + T_1(\psi)w + T_2(\psi)[w, w] + T_{\geq 3}(\psi, w)$ , where

$$T_\delta(\psi) := (A_\varepsilon \circ G_\delta)(\psi, 0, 0) = \varepsilon v_\delta(\psi) + \varepsilon^b z_0(\psi), \quad v_\delta(\psi) := \sum_{j \in S} \sqrt{\xi_j + \varepsilon^{2(b-1)} |j| [y_\delta(\psi)]_j} e^{i[\theta_0(\psi)]_j} e^{ijx} \quad (7.11)$$

is the approximate isotropic torus in phase space (it corresponds to  $i_\delta$  in Lemma 6.3),

$$T_1(\psi)w = \varepsilon \sum_{j \in S} \frac{\varepsilon^{2(b-1)} |j| [L_2(\psi)w]_j e^{i[\theta_0(\psi)]_j}}{2\sqrt{\xi_j + \varepsilon^{2(b-1)} |j| [y_\delta(\psi)]_j}} e^{ijx} + \varepsilon^b w =: \varepsilon^{2b-1} U_1(\psi)w + \varepsilon^b w \quad (7.12)$$

$$T_2(\psi)[w, w] = -\varepsilon \sum_{j \in S} \frac{\varepsilon^{4(b-1)} j^2 [L_2(\psi)w]_j^2 e^{i[\theta_0(\psi)]_j}}{8\{\xi_j + \varepsilon^{2(b-1)} |j| [y_\delta(\psi)]_j\}^{3/2}} e^{ijx} =: \varepsilon^{4b-3} U_2(\psi)[w, w] \quad (7.13)$$

and  $T_{\geq 3}(\psi, w)$  collects all the terms of order at least cubic in  $w$ . In the notation of (4.7), the function  $v_\delta(\psi)$  in (7.11) is  $v_\delta(\psi) = v_\varepsilon(\theta_0(\psi), y_\delta(\psi))$ . The terms  $U_1, U_2 = O(1)$  in  $\varepsilon$ . Moreover, using that  $L_2(\psi)$  in (7.10) vanishes as  $z_0 = 0$ , they satisfy

$$\|U_1 w\|_s \leq \|\mathfrak{I}_\delta\|_s \|w\|_{s_0} + \|\mathfrak{I}_\delta\|_{s_0} \|w\|_s, \quad \|U_2[w, w]\|_s \leq \|\mathfrak{I}_\delta\|_s \|\mathfrak{I}_\delta\|_{s_0} \|w\|_{s_0}^2 + \|\mathfrak{I}_\delta\|_{s_0}^2 \|w\|_{s_0} \|w\|_s \quad (7.14)$$

and also in the  $\|\cdot\|_s^{\text{Lip}(\gamma)}$ -norm.

By Taylor's formula  $\mathcal{H}(u+h) = \mathcal{H}(u) + ((\nabla \mathcal{H})(u), h)_{L^2(\mathbb{T})} + \frac{1}{2}((\partial_u \nabla \mathcal{H})(u)[h], h)_{L^2(\mathbb{T})} + O(h^3)$ . Specifying at  $u = T_\delta(\psi)$  and  $h = T_1(\psi)w + T_2(\psi)[w, w] + T_{\geq 3}(\psi, w)$ , we obtain that the sum of all the components of  $K = \varepsilon^{-2b}(\mathcal{H} \circ A_\varepsilon \circ G_\delta)(\psi, 0, w)$  that are quadratic in  $w$  is

$$\frac{1}{2}(K_{02}w, w)_{L^2(\mathbb{T})} = \varepsilon^{-2b}((\nabla \mathcal{H})(T_\delta), T_2[w, w])_{L^2(\mathbb{T})} + \varepsilon^{-2b} \frac{1}{2}((\partial_u \nabla \mathcal{H})(T_\delta)[T_1 w], T_1 w)_{L^2(\mathbb{T})}.$$

Inserting the expressions (7.12), (7.13) we get

$$\begin{aligned} K_{02}(\psi)w &= (\partial_u \nabla \mathcal{H})(T_\delta)[w] + 2\varepsilon^{b-1}(\partial_u \nabla \mathcal{H})(T_\delta)[U_1 w] + \varepsilon^{2(b-1)}U_1^T (\partial_u \nabla \mathcal{H})(T_\delta)[U_1 w] \\ &\quad + 2\varepsilon^{2b-3}U_2[w, \cdot]^T (\nabla \mathcal{H})(T_\delta). \end{aligned} \quad (7.15)$$

**Lemma 7.4.**

$$(K_{02}(\psi)w, w)_{L^2(\mathbb{T})} = ((\partial_u \nabla \mathcal{H})(T_\delta)[w], w)_{L^2(\mathbb{T})} + (R(\psi)w, w)_{L^2(\mathbb{T})} \quad (7.16)$$

where  $R(\psi)w$  has the “finite dimensional” form

$$R(\psi)w = \sum_{|j| \leq C} (w, g_j(\psi))_{L^2(\mathbb{T})} \chi_j(\psi) \quad (7.17)$$

where, for some  $\sigma := \sigma(\nu, \tau) > 0$ ,

$$\|g_j\|_s^{\text{Lip}(\gamma)} \|\chi_j\|_{s_0}^{\text{Lip}(\gamma)} + \|g_j\|_{s_0}^{\text{Lip}(\gamma)} \|\chi_j\|_s^{\text{Lip}(\gamma)} \leq_s \varepsilon^{b+1} \|\mathfrak{J}_\delta\|_{s+\sigma}^{\text{Lip}(\gamma)} \quad (7.18)$$

$$\begin{aligned} \|\partial_i g_j[\hat{v}]\|_s \|\chi_j\|_{s_0} + \|\partial_i g_j[\hat{v}]\|_{s_0} \|\chi_j\|_s + \|g_j\|_{s_0} \|\partial_i \chi_j[\hat{v}]\|_s + \|g_j\|_s \|\partial_i \chi_j[\hat{v}]\|_{s_0} &\leq_s \varepsilon^{b+1} \|\hat{v}\|_{s+\sigma} \\ &\quad + \varepsilon^{2b-1} \|\mathfrak{J}_\delta\|_{s+\sigma} \|\hat{v}\|_{s_0+\sigma}, \end{aligned} \quad (7.19)$$

and, as usual,  $i = (\theta, y, z)$  (see (5.1)),  $\hat{v} = (\hat{\theta}, \hat{y}, \hat{z})$ .

*Proof.* Since  $U_1 = \Pi_S U_1$  and  $U_2 = \Pi_S U_2$ , the last three terms in (7.15) have all the form (7.17) (argue as in Lemma 7.1). We now prove that they are also small in size.

The contributions in (7.15) from  $H_2$  are better analyzed by the expression

$$\varepsilon^{-2b} H_2 \circ A_\varepsilon \circ G_\delta(\psi, \eta, w) = \text{const} + \sum_{j \in S^+} j^3 [y_\delta(\psi) + L_1(\psi)\eta + L_2(\psi)w]_j + \frac{1}{2} \int_{\mathbb{T}} (z_0(\psi) + w)_x^2 dx$$

which follows by (4.8), (6.18), (7.10). Hence the only contribution to  $(K_{02}w, w)$  is  $\int_{\mathbb{T}} w_x^2 dx$ . Now we consider the cubic term  $\mathcal{H}_3$  in (3.6). A direct calculation shows that for  $u = v + z$ ,  $\nabla \mathcal{H}_3(u) = 3z^2 + 6\Pi_S^\perp(vz)$ , and  $\partial_u \nabla \mathcal{H}_3(u)[U_1 w] = 6\Pi_S^\perp(zU_1 w)$  (since  $U_1 w \in H_S$ ). Therefore

$$\nabla \mathcal{H}_3(T_\delta) = 3\varepsilon^{2b} z_0^2 + 6\varepsilon^{b+1} \Pi_S^\perp(v_\delta z_0), \quad \partial_u \nabla \mathcal{H}_3(T_\delta)[U_1 w] = 6\varepsilon^b \Pi_S^\perp(z_0 U_1 w). \quad (7.20)$$

By (7.20) one has  $((\partial_u \nabla \mathcal{H}_3)(T_\delta)[U_1 w], U_1 w)_{L^2(\mathbb{T})} = 0$ , and since also  $U_2 = \Pi_S U_2$ ,

$$\varepsilon^{b-1} \partial_u \nabla \mathcal{H}_3(T_\delta)[U_1 w] + \varepsilon^{2b-3} U_2[w, \cdot]^T \nabla \mathcal{H}_3(T_\delta) = 6\varepsilon^{2b-1} \Pi_S^\perp(z_0 U_1 w) + 3\varepsilon^{4b-3} U_2[w, \cdot]^T z_0^2. \quad (7.21)$$

These terms have the form (7.17) and, using (7.14), (6.4), they satisfy (7.18).

Finally we consider all the terms which arise from  $\mathcal{H}_{\geq 4} = O(u^4)$ . The operators  $\varepsilon^{b-1} \partial_u \nabla \mathcal{H}_{\geq 4}(T_\delta) U_1$ ,  $\varepsilon^{2(b-1)} U_1^T (\partial_u \nabla \mathcal{H}_{\geq 4})(T_\delta) U_1$ ,  $\varepsilon^{2b-3} U_2^T \nabla \mathcal{H}_{\geq 4}(T_\delta)$  have the form (7.17) and, using  $\|T_\delta\|_s^{\text{Lip}(\gamma)} \leq \varepsilon(1 + \|\mathfrak{J}_\delta\|_s^{\text{Lip}(\gamma)})$ , (7.14), (6.4), the bound (7.18) holds. Notice that the biggest term is  $\varepsilon^{b-1} \partial_u \nabla \mathcal{H}_{\geq 4}(T_\delta) U_1$ .

By (6.11) and using explicit formulae (7.10)-(7.13) we get estimate (7.19).  $\square$

The conclusion of this section is that, after the composition with the action-angle variables, the rescaling (4.5), and the transformation  $G_\delta$ , the linearized operator to analyze is  $H_S^\perp \ni w \mapsto (\partial_u \nabla \mathcal{H})(T_\delta)[w]$ , up to finite dimensional operators which have the form (7.17) and size (7.18).

## 7.2 The linearized operator in the normal directions

In view of (7.16) we now compute  $((\partial_u \nabla \mathcal{H})(T_\delta)[w], w)_{L^2(\mathbb{T})}$ ,  $w \in H_S^\perp$ , where  $\mathcal{H} = H \circ \Phi_B$  and  $\Phi_B$  is the Birkhoff map of Proposition 3.1. It is convenient to estimate separately the terms in

$$\mathcal{H} = H \circ \Phi_B = (H_2 + H_3) \circ \Phi_B + H_{\geq 5} \circ \Phi_B \quad (7.22)$$

where  $H_2, H_3, H_{\geq 5}$  are defined in (3.1).

We first consider  $H_{\geq 5} \circ \Phi_B$ . By (3.1) we get  $\nabla H_{\geq 5}(u) = \pi_0[(\partial_u f)(x, u, u_x)] - \partial_x\{(\partial_{u_x} f)(x, u, u_x)\}$ , see (2.2). Since the Birkhoff transformation  $\Phi_B$  has the form (3.4), Lemma 7.1 (at  $u = T_\delta$ , see (7.11)) implies that

$$\begin{aligned} \partial_u \nabla(H_{\geq 5} \circ \Phi_B)(T_\delta)[h] &= (\partial_u \nabla H_{\geq 5})(\Phi_B(T_\delta))[h] + \mathcal{R}_{H_{\geq 5}}(T_\delta)[h] \\ &= \partial_x(r_1(T_\delta)\partial_x h) + r_0(T_\delta)h + \mathcal{R}_{H_{\geq 5}}(T_\delta)[h] \end{aligned} \quad (7.23)$$

where the multiplicative functions  $r_0(T_\delta)$ ,  $r_1(T_\delta)$  are

$$r_0(T_\delta) := \sigma_0(\Phi_B(T_\delta)), \quad \sigma_0(u) := (\partial_{uu} f)(x, u, u_x) - \partial_x\{(\partial_{u_x u} f)(x, u, u_x)\}, \quad (7.24)$$

$$r_1(T_\delta) := \sigma_1(\Phi_B(T_\delta)), \quad \sigma_1(u) := -(\partial_{u_x u_x} f)(x, u, u_x), \quad (7.25)$$

the remainder  $\mathcal{R}_{H_{\geq 5}}(u)$  has the form (7.2) with  $\chi_j = e^{ijx}$  or  $g_j = e^{ijx}$  and, using (7.3), it satisfies, for some  $\sigma := \sigma(\nu, \tau) > 0$ ,

$$\begin{aligned} &\|g_j\|_s^{\text{Lip}(\gamma)} \|\chi_j\|_{s_0}^{\text{Lip}(\gamma)} + \|g_j\|_{s_0}^{\text{Lip}(\gamma)} \|\chi_j\|_s^{\text{Lip}(\gamma)} \leq \varepsilon^4 (1 + \|\mathcal{J}_\delta\|_{s+2}^{\text{Lip}(\gamma)}) \\ \|\partial_i g_j[\tilde{v}]\|_s \|\chi_j\|_{s_0} + \|\partial_i g_j[\tilde{v}]\|_{s_0} \|\chi_j\|_s + \|g_j\|_{s_0} \|\partial_i \chi_j[\tilde{v}]\|_s + \|g_j\|_s \|\partial_i \chi_j[\tilde{v}]\|_{s_0} &\leq \varepsilon^4 (\|\tilde{v}\|_{s+\sigma} + \|\mathcal{J}_\delta\|_{s+2} \|\tilde{v}\|_{s_0+2}). \end{aligned}$$

Now we consider the contributions from  $(H_2 + H_3) \circ \Phi_B$ . By Lemma 7.1 and the expressions of  $H_2, H_3$  in (3.1) we deduce that

$$\partial_u \nabla(H_2 \circ \Phi_B)(T_\delta)[h] = -\partial_{xx} h + \mathcal{R}_{H_2}(T_\delta)[h], \quad \partial_u \nabla(H_3 \circ \Phi_B)(T_\delta)[h] = 6\Phi_B(T_\delta)h + \mathcal{R}_{H_3}(T_\delta)[h],$$

where  $\Phi_B(T_\delta)$  is a function with zero space average, because  $\Phi_B : H_0^1(\mathbb{T}_x) \rightarrow H_0^1(\mathbb{T}_x)$  (Proposition 3.1) and  $\mathcal{R}_{H_2}(u), \mathcal{R}_{H_3}(u)$  have the form (7.2). By (7.3), the size  $(\mathcal{R}_{H_2} + \mathcal{R}_{H_3})(T_\delta) = O(\varepsilon)$ . We expand

$$(\mathcal{R}_{H_2} + \mathcal{R}_{H_3})(T_\delta) = \varepsilon \mathcal{R}_1 + \varepsilon^2 \mathcal{R}_2 + \tilde{\mathcal{R}}_{>2},$$

where  $\tilde{\mathcal{R}}_{>2}$  has size  $o(\varepsilon^2)$ , and we get,  $\forall h \in H_S^\perp$ ,

$$\Pi_S^\perp \partial_u \nabla((H_2 + H_3) \circ \Phi_B)(T_\delta)[h] = -\partial_{xx} h + \Pi_S^\perp (6\Phi_B(T_\delta)h) + \Pi_S^\perp (\varepsilon \mathcal{R}_1 + \varepsilon^2 \mathcal{R}_2 + \tilde{\mathcal{R}}_{>2})[h]. \quad (7.26)$$

We also develop the function  $\Phi_B(T_\delta)$  in powers of  $\varepsilon$ . Expand  $\Phi_B(u) = u + \Psi_2(u) + \Psi_{\geq 3}(u)$ , where  $\Psi_2(u)$  is quadratic,  $\Psi_{\geq 3}(u) = O(u^3)$ , and both map  $H_0^1(\mathbb{T}_x) \rightarrow H_0^1(\mathbb{T}_x)$ . At  $u = T_\delta = \varepsilon v_\delta + \varepsilon^b z_0$  we get

$$\Phi_B(T_\delta) = T_\delta + \Psi_2(T_\delta) + \Psi_{\geq 3}(T_\delta) = \varepsilon v_\delta + \varepsilon^2 \Psi_2(v_\delta) + \tilde{q} \quad (7.27)$$

where  $\tilde{q} := \varepsilon^b z_0 + \Psi_2(T_\delta) - \varepsilon^2 \Psi_2(v_\delta) + \Psi_{\geq 3}(T_\delta)$  has zero space average and it satisfies

$$\|\tilde{q}\|_s^{\text{Lip}(\gamma)} \leq \varepsilon^3 + \varepsilon^b \|\mathcal{J}_\delta\|_s^{\text{Lip}(\gamma)}, \quad \|\partial_i \tilde{q}[\tilde{v}]\|_s \leq \varepsilon^b (\|\tilde{v}\|_s + \|\mathcal{J}_\delta\|_s \|\tilde{v}\|_{s_0}).$$

In particular, its low norm  $\|\tilde{q}\|_{s_0}^{\text{Lip}(\gamma)} \leq \varepsilon^{6-b} \gamma^{-1} = o(\varepsilon^2)$ .

We need an exact expression of the terms of order  $\varepsilon$  and  $\varepsilon^2$  in (7.26). We compare the Hamiltonian (3.5) with (7.22), noting that  $(H_{\geq 5} \circ \Phi_B)(u) = O(u^5)$  because  $f$  satisfies (1.5) and  $\Phi_B(u) = O(u)$ . Therefore

$$(H_2 + H_3) \circ \Phi_B = H_2 + \mathcal{H}_3 + \mathcal{H}_4 + O(u^5),$$

and the homogeneous terms of  $(H_2 + H_3) \circ \Phi_B$  of degree 2, 3, 4 in  $u$  are  $H_2, \mathcal{H}_3, \mathcal{H}_4$  respectively. As a consequence, the terms of order  $\varepsilon$  and  $\varepsilon^2$  in (7.26) (both in the function  $\Phi_B(T_\delta)$  and in the remainders  $\mathcal{R}_1, \mathcal{R}_2$ ) come only from  $H_2 + \mathcal{H}_3 + \mathcal{H}_4$ . Actually they come from  $H_2, \mathcal{H}_3$  and  $\mathcal{H}_{4,2}$  (see (3.6), (3.7)) because, at  $u = T_\delta = \varepsilon v_\delta + \varepsilon^b z_0$ , for all  $h \in H_S^\perp$ ,

$$\Pi_S^\perp (\partial_u \nabla \mathcal{H}_4)(T_\delta)[h] = \Pi_S^\perp (\partial_u \nabla \mathcal{H}_{4,2})(T_\delta)[h] + o(\varepsilon^2).$$

A direct calculation based on the expressions (3.6), (3.7) shows that, for all  $h \in H_S^\perp$ ,

$$\begin{aligned} \Pi_S^\perp (\partial_u \nabla (H_2 + \mathcal{H}_3 + \mathcal{H}_4))(T_\delta)[h] &= -\partial_{xx} h + 6\varepsilon \Pi_S^\perp (v_\delta h) + 6\varepsilon^b \Pi_S^\perp (z_0 h) + \varepsilon^2 \Pi_S^\perp \{6\pi_0 [(\partial_x^{-1} v_\delta)^2] h \\ &\quad + 6v_\delta \Pi_S [(\partial_x^{-1} v_\delta)(\partial_x^{-1} h)] - 6\partial_x^{-1} \{(\partial_x^{-1} v_\delta) \Pi_S [v_\delta h]\}\} + o(\varepsilon^2). \end{aligned} \quad (7.28)$$



Thus, comparing the terms of order  $\varepsilon, \varepsilon^2$  in (7.26) (using (7.27)) with those in (7.28) we deduce that the operators  $\mathcal{R}_1, \mathcal{R}_2$  and the function  $\Psi_2(v_\delta)$  are

$$\mathcal{R}_1 = 0, \quad \mathcal{R}_2[h] = 6v_\delta \Pi_S [(\partial_x^{-1} v_\delta)(\partial_x^{-1} h)] - 6\partial_x^{-1} \{(\partial_x^{-1} v_\delta) \Pi_S [v_\delta h]\}, \quad \Psi_2(v_\delta) = \pi_0 [(\partial_x^{-1} v_\delta)^2]. \quad (7.29)$$

In conclusion, by (7.22), (7.26), (7.23), (7.27), (7.29), we get, for all  $h \in H_{S^\perp}$ ,

$$\begin{aligned} \Pi_S^\perp \partial_u \nabla \mathcal{H}(T_\delta)[h] &= -\partial_{xx} h + \Pi_S^\perp [(\varepsilon 6v_\delta + \varepsilon^2 6\pi_0 [(\partial_x^{-1} v_\delta)^2] + q_{>2} + p_{\geq 4})h] \\ &\quad + \Pi_S^\perp \partial_x (r_1(T_\delta) \partial_x h) + \varepsilon^2 \Pi_S^\perp \mathcal{R}_2[h] + \Pi_S^\perp \mathcal{R}_{>2}[h] \end{aligned} \quad (7.30)$$

where  $r_1$  is defined in (7.24),  $\mathcal{R}_2$  in (7.29), the remainder  $\mathcal{R}_{>2} := \tilde{\mathcal{R}}_{>2} + \mathcal{R}_{H_{\geq 5}^\perp}(T_\delta)$  and the functions (using also (7.24), (7.25), (1.5)),

$$q_{>2} := 6\tilde{q} + \varepsilon^3 ((\partial_{uu} f_5)(v_\delta, (v_\delta)_x) - \partial_x \{(\partial_{uu_x} f_5)(v_\delta, (v_\delta)_x)\}) \quad (7.31)$$

$$p_{\geq 4} := r_0(T_\delta) - \varepsilon^3 [(\partial_{uu} f_5)(v_\delta, (v_\delta)_x) - \partial_x \{(\partial_{uu_x} f_5)(v_\delta, (v_\delta)_x)\}]. \quad (7.32)$$

**Lemma 7.5.**  $\int_{\mathbb{T}} q_{>2} dx = 0$ .

*Proof.* We already observed that  $\tilde{q}$  has zero  $x$ -average as well as the derivative  $\partial_x \{(\partial_{uu_x} f_5)(v, v_x)\}$ . Finally

$$(\partial_{uu} f_5)(v, v_x) = \sum_{j_1, j_2, j_3 \in S} c_{j_1 j_2 j_3} v_{j_1} v_{j_2} v_{j_3} e^{i(j_1 + j_2 + j_3)x}, \quad v := \sum_{j \in S} v_j e^{ijx} \quad (7.33)$$

for some coefficient  $c_{j_1 j_2 j_3}$ , and therefore it has zero average by hypothesis (S1).  $\square$

By Lemma 7.4 and the results of this section (in particular (7.30)) we deduce:

**Proposition 7.6.** *Assume (7.8). Then the Hamiltonian operator  $\mathcal{L}_\omega$  has the form,  $\forall h \in H_{S^\perp}^s(\mathbb{T}^{\nu+1})$ ,*

$$\mathcal{L}_\omega h := \omega \cdot \partial_\varphi h - \partial_x K_{02} h = \Pi_S^\perp (\omega \cdot \partial_\varphi h + \partial_{xx} (a_1 \partial_x h) + \partial_x (a_0 h) - \varepsilon^2 \partial_x \mathcal{R}_2 h - \partial_x \mathcal{R}_* h) \quad (7.34)$$

where  $\mathcal{R}_2$  is defined in (7.29),  $\mathcal{R}_* := \mathcal{R}_{>2} + R(\psi)$  (with  $R(\psi)$  defined in Lemma 7.4), the functions

$$a_1 := 1 - r_1(T_\delta), \quad a_0 := -(\varepsilon p_1 + \varepsilon^2 p_2 + q_{>2} + p_{\geq 4}), \quad p_1 := 6v_\delta, \quad p_2 := 6\pi_0 [(\partial_x^{-1} v_\delta)^2], \quad (7.35)$$

the function  $q_{>2}$  is defined in (7.31) and satisfies  $\int_{\mathbb{T}} q_{>2} dx = 0$ , the function  $p_{\geq 4}$  is defined in (7.32),  $r_1$  in (7.25),  $T_\delta$  and  $v_\delta$  in (7.11). For  $p_k = p_1, p_2$ ,

$$\|p_k\|_s^{\text{Lip}(\gamma)} \leq_s 1 + \|\mathfrak{J}_\delta\|_s^{\text{Lip}(\gamma)}, \quad \|\partial_i p_k[\tilde{v}]\|_s \leq_s \|\tilde{v}\|_{s+1} + \|\mathfrak{J}_\delta\|_{s+1} \|\tilde{v}\|_{s_0+1}, \quad (7.36)$$

$$\|q_{>2}\|_s^{\text{Lip}(\gamma)} \leq_s \varepsilon^3 + \varepsilon^b \|\mathfrak{J}_\delta\|_s^{\text{Lip}(\gamma)}, \quad \|\partial_i q_{>2}[\tilde{v}]\|_s \leq_s \varepsilon^b (\|\tilde{v}\|_{s+1} + \|\mathfrak{J}_\delta\|_{s+1} \|\tilde{v}\|_{s_0+1}), \quad (7.37)$$

$$\|a_1 - 1\|_s^{\text{Lip}(\gamma)} \leq_s \varepsilon^3 (1 + \|\mathfrak{J}_\delta\|_{s+1}^{\text{Lip}(\gamma)}), \quad \|\partial_i a_1[\tilde{v}]\|_s \leq_s \varepsilon^3 (\|\tilde{v}\|_{s+1} + \|\mathfrak{J}_\delta\|_{s+1} \|\tilde{v}\|_{s_0+1}) \quad (7.38)$$

$$\|p_{\geq 4}\|_s^{\text{Lip}(\gamma)} \leq_s \varepsilon^4 + \varepsilon^{b+2} \|\mathfrak{J}_\delta\|_{s+2}^{\text{Lip}(\gamma)}, \quad \|\partial_i p_{\geq 4}[\tilde{v}]\|_s \leq_s \varepsilon^{b+2} (\|\tilde{v}\|_{s+2} + \|\mathfrak{J}_\delta\|_{s+2} \|\tilde{v}\|_{s_0+2}) \quad (7.39)$$

where  $\mathfrak{J}_\delta(\varphi) := (\theta_0(\varphi) - \varphi, y_\delta(\varphi), z_0(\varphi))$  corresponds to  $T_\delta$ . The remainder  $\mathcal{R}_2$  has the form (7.2) with

$$\|g_j\|_s^{\text{Lip}(\gamma)} + \|\chi_j\|_s^{\text{Lip}(\gamma)} \leq_s 1 + \|\mathfrak{J}_\delta\|_{s+\sigma}^{\text{Lip}(\gamma)}, \quad \|\partial_i g_j[\tilde{v}]\|_s + \|\partial_i \chi_j[\tilde{v}]\|_s \leq_s \|\tilde{v}\|_{s+\sigma} + \|\mathfrak{J}_\delta\|_{s+\sigma} \|\tilde{v}\|_{s_0+\sigma} \quad (7.40)$$

and also  $\mathcal{R}_*$  has the form (7.2) with

$$\|g_j^*\|_s^{\text{Lip}(\gamma)} \|\chi_j^*\|_{s_0}^{\text{Lip}(\gamma)} + \|g_j^*\|_{s_0}^{\text{Lip}(\gamma)} \|\chi_j^*\|_s^{\text{Lip}(\gamma)} \leq_s \varepsilon^3 + \varepsilon^{b+1} \|\mathfrak{J}_\delta\|_{s+\sigma}^{\text{Lip}(\gamma)} \quad (7.41)$$

$$\begin{aligned} \|\partial_i g_j^*[\tilde{v}]\|_s \|\chi_j^*\|_{s_0} + \|\partial_i g_j^*[\tilde{v}]\|_{s_0} \|\chi_j^*\|_s + \|g_j^*\|_{s_0} \|\partial_i \chi_j^*[\tilde{v}]\|_s + \|g_j^*\|_s \|\partial_i \chi_j^*[\tilde{v}]\|_{s_0} &\leq_s \varepsilon^{b+1} \|\tilde{v}\|_{s+\sigma} \\ &\quad + \varepsilon^{2b-1} \|\mathfrak{J}_\delta\|_{s+\sigma} \|\tilde{v}\|_{s_0+\sigma}. \end{aligned} \quad (7.42)$$

The bounds (7.40), (7.41) imply, by Lemma 7.3, estimates for the  $s$ -decay norms of  $\mathcal{R}_2$  and  $\mathcal{R}_*$ . The linearized operator  $\mathcal{L}_\omega := \mathcal{L}_\omega(\omega, i_\delta(\omega))$  depends on the parameter  $\omega$  both directly and also through the dependence on the torus  $i_\delta(\omega)$ . We have estimated also the partial derivative  $\partial_i$  with respect to the variables  $i$  (see (5.1)) in order to control, along the nonlinear Nash-Moser iteration, the Lipschitz variation of the eigenvalues of  $\mathcal{L}_\omega$  with respect to  $\omega$  and the approximate solution  $i_\delta$ .

## 8 Reduction of the linearized operator in the normal directions

The goal of this section is to conjugate the Hamiltonian operator  $\mathcal{L}_\omega$  in (7.34) to the diagonal operator  $\mathcal{L}_\infty$  defined in (8.121). The proof is obtained applying different kind of symplectic transformations. We shall always assume (7.8).

### 8.1 Change of the space variable

The first task is to conjugate  $\mathcal{L}_\omega$  in (7.34) to  $\mathcal{L}_1$  in (8.31), which has the coefficient of  $\partial_{xxx}$  independent on the space variable. We look for a  $\varphi$ -dependent family of *symplectic* diffeomorphisms  $\Phi(\varphi)$  of  $H_S^\perp$  which differ from

$$\mathcal{A}_\perp := \Pi_S^\perp \mathcal{A} \Pi_S^\perp, \quad (\mathcal{A}h)(\varphi, x) := (1 + \beta_x(\varphi, x))h(\varphi, x + \beta(\varphi, x)), \quad (8.1)$$

up to a small “finite dimensional” remainder, see (8.6). Each  $\mathcal{A}(\varphi)$  is a symplectic map of the phase space, see [2]-Remark 3.3. If  $\|\beta\|_{W^{1,\infty}} < 1/2$  then  $\mathcal{A}$  is invertible, see Lemma 2.4, and its inverse and adjoint maps are

$$(\mathcal{A}^{-1}h)(\varphi, y) := (1 + \tilde{\beta}_y(\varphi, y))h(\varphi, y + \tilde{\beta}(\varphi, y)), \quad (\mathcal{A}^T h)(\varphi, y) = h(\varphi, y + \tilde{\beta}(\varphi, y)) \quad (8.2)$$

where  $x = y + \tilde{\beta}(\varphi, y)$  is the inverse diffeomorphism (of  $\mathbb{T}$ ) of  $y = x + \beta(\varphi, x)$ .

The restricted maps  $\mathcal{A}_\perp(\varphi) : H_S^\perp \rightarrow H_S^\perp$  are not symplectic. In order to find a symplectic diffeomorphism near  $\mathcal{A}_\perp(\varphi)$ , the first observation is that each  $\mathcal{A}(\varphi)$  can be seen as the time 1-flow of a time dependent Hamiltonian PDE. Indeed  $\mathcal{A}(\varphi)$  (for simplicity we skip the dependence on  $\varphi$ ) is homotopic to the identity via the path of symplectic diffeomorphisms

$$u \mapsto (1 + \tau\beta_x)u(x + \tau\beta(x)), \quad \tau \in [0, 1],$$

which is the trajectory solution of the time dependent, linear Hamiltonian PDE

$$\partial_\tau u = \partial_x(b(\tau, x)u), \quad b(\tau, x) := \frac{\beta(x)}{1 + \tau\beta_x(x)}, \quad (8.3)$$

with value  $u(x)$  at  $\tau = 0$  and  $\mathcal{A}u = (1 + \beta_x(x))u(x + \beta(x))$  at  $\tau = 1$ . The equation (8.3) is a *transport* equation. Its associated characteristic ODE is

$$\frac{d}{d\tau}x = -b(\tau, x). \quad (8.4)$$

We denote its flow by  $\gamma^{\tau_0, \tau}$ , namely  $\gamma^{\tau_0, \tau}(y)$  is the solution of (8.4) with  $\gamma^{\tau_0, \tau_0}(y) = y$ . Each  $\gamma^{\tau_0, \tau}$  is a diffeomorphism of the torus  $\mathbb{T}_x$ .

**Remark 8.1.** Let  $y \mapsto y + \tilde{\beta}(\tau, y)$  be the inverse diffeomorphism of  $x \mapsto x + \tau\beta(x)$ . Differentiating the identity  $\tilde{\beta}(\tau, y) + \tau\beta(y + \tilde{\beta}(\tau, y)) = 0$  with respect to  $\tau$  it results that  $\gamma^\tau(y) := \gamma^{0, \tau}(y) = y + \tilde{\beta}(\tau, y)$ .  $\square$

Then we define a symplectic map  $\Phi$  of  $H_S^\perp$  as the time-1 flow of the Hamiltonian PDE

$$\partial_\tau u = \Pi_S^\perp \partial_x(b(\tau, x)u) = \partial_x(b(\tau, x)u) - \Pi_S \partial_x(b(\tau, x)u), \quad u \in H_S^\perp. \quad (8.5)$$

Note that  $\Pi_S^\perp \partial_x(b(\tau, x)u)$  is the Hamiltonian vector field generated by  $\frac{1}{2} \int_{\mathbb{T}} b(\tau, x)u^2 dx$  restricted to  $H_S^\perp$ . We denote by  $\Phi^{\tau_0, \tau}$  the flow of (8.5), namely  $\Phi^{\tau_0, \tau}(u_0)$  is the solution of (8.5) with initial condition  $\Phi^{\tau_0, \tau_0}(u_0) = u_0$ . The flow is well defined in Sobolev spaces  $H_{S^\perp}^s(\mathbb{T}_x)$  for  $b(\tau, x)$  is smooth enough (standard theory of linear hyperbolic PDEs, see e.g. section 0.8 in [29]). It is natural to expect that the difference between the flow map  $\Phi := \Phi^{0, 1}$  and  $\mathcal{A}_\perp$  is a “finite-dimensional” remainder of the size of  $\beta$ .

**Lemma 8.2.** *For  $\|\beta\|_{W^{s_0+1, \infty}}$  small, there exists an invertible symplectic transformation  $\Phi = \mathcal{A}_\perp + \mathcal{R}_\Phi$  of  $H_{S^\perp}^s$ , where  $\mathcal{A}_\perp$  is defined in (8.1) and  $\mathcal{R}_\Phi$  is a “finite-dimensional” remainder*

$$\mathcal{R}_\Phi h = \sum_{j \in S} \int_0^1 (h, g_j(\tau))_{L^2(\mathbb{T})} \chi_j(\tau) d\tau + \sum_{j \in S} (h, \psi_j)_{L^2(\mathbb{T})} e^{ijx} \quad (8.6)$$

for some functions  $\chi_j(\tau), g_j(\tau), \psi_j \in H^s$  satisfying

$$\|\psi_j\|_s, \|g_j(\tau)\|_s \leq_s \|\beta\|_{W^{s+2,\infty}}, \quad \|\chi_j(\tau)\|_s \leq_s 1 + \|\beta\|_{W^{s+1,\infty}}, \quad \forall \tau \in [0, 1]. \quad (8.7)$$

Furthermore, the following tame estimates holds

$$\|\Phi^{\pm 1} h\|_s \leq_s \|h\|_s + \|\beta\|_{W^{s+2,\infty}} \|h\|_{s_0}, \quad \forall h \in H_{S^\perp}^s. \quad (8.8)$$

*Proof.* Let  $w(\tau, x) := (\Phi^\tau u_0)(x)$  denote the solution of (8.5) with initial condition  $\Phi^0(w) = u_0 \in H_S^\perp$ . The difference

$$(\mathcal{A}_\perp - \Phi)u_0 = \Pi_S^\perp \mathcal{A}u_0 - w(1, \cdot) = \mathcal{A}u_0 - w(1, \cdot) - \Pi_S \mathcal{A}u_0, \quad \forall u_0 \in H_S^\perp, \quad (8.9)$$

and

$$\Pi_S \mathcal{A}u_0 = \Pi_S (\mathcal{A} - I) \Pi_S^\perp u_0 = \sum_{j \in S} (u_0, \psi_j)_{L^2(\mathbb{T})} e^{ijx}, \quad \psi_j := (\mathcal{A}^T - I) e^{ijx}. \quad (8.10)$$

We claim that the difference

$$\mathcal{A}u_0 - w(1, x) = (1 + \beta_x(x)) \int_0^1 (1 + \tau \beta_x(x))^{-1} [\Pi_S \partial_x (b(\tau) w(\tau))] (\gamma^\tau(x + \beta(x))) d\tau \quad (8.11)$$

where  $\gamma^\tau(y) := \gamma^{0,\tau}(y)$  is the flow of (8.4). Indeed the solution  $w(\tau, x)$  of (8.5) satisfies

$$\partial_\tau \{w(\tau, \gamma^\tau(y))\} = b_x(\tau, \gamma^\tau(y)) w(\tau, \gamma^\tau(y)) - [\Pi_S \partial_x (b(\tau) w(\tau))] (\gamma^\tau(y)).$$

Then, by the variation of constant formula, we find

$$w(\tau, \gamma^\tau(y)) = e^{\int_0^\tau b_x(s, \gamma^s(y)) ds} \left( u_0(y) - \int_0^\tau e^{-\int_0^s b_x(\zeta, \gamma^\zeta(y)) d\zeta} [\Pi_S \partial_x (b(s) w(s))] (\gamma^s(y)) ds \right).$$

Since  $\partial_y \gamma^\tau(y)$  solves the variational equation  $\partial_\tau (\partial_y \gamma^\tau(y)) = -b_x(\tau, \gamma^\tau(y)) (\partial_y \gamma^\tau(y))$  with  $\partial_y \gamma^0(y) = 1$  we have that

$$e^{\int_0^\tau b_x(s, \gamma^s(y)) ds} = (\partial_y \gamma^\tau(y))^{-1} = 1 + \tau \beta_x(x) \quad (8.12)$$

by remark 8.1, and so we derive the expression

$$w(\tau, x) = (1 + \tau \beta_x(x)) \left\{ u_0(x + \tau \beta(x)) - \int_0^\tau (1 + s \beta_x(x))^{-1} [\Pi_S \partial_x (b(s) w(s))] (\gamma^s(x + \tau \beta(x))) ds \right\}.$$

Evaluating at  $\tau = 1$ , formula (8.11) follows. Next, we develop (recall  $w(\tau) = \Phi^\tau(u_0)$ )

$$[\Pi_S \partial_x (b(\tau) w(\tau))](x) = \sum_{j \in S} (u_0, g_j(\tau))_{L^2(\mathbb{T})} e^{ijx}, \quad g_j(\tau) := -(\Phi^\tau)^T [b(\tau) \partial_x e^{ijx}], \quad (8.13)$$

and (8.11) becomes

$$\mathcal{A}u_0 - w(1, \cdot) = - \int_0^1 \sum_{j \in S} (u_0, g_j(\tau))_{L^2(\mathbb{T})} \chi_j(\tau, \cdot) d\tau, \quad (8.14)$$

where

$$\chi_j(\tau, x) := -(1 + \beta_x(x)) (1 + \tau \beta_x(x))^{-1} e^{ij\gamma^\tau(x + \beta(x))}. \quad (8.15)$$

By (8.9), (8.10), (8.11), (8.14) we deduce that  $\Phi = \mathcal{A}_\perp + \mathcal{R}_\Phi$  as in (8.6).

We now prove the estimates (8.7). Each function  $\psi_j$  in (8.10) satisfies  $\|\psi_j\|_s \leq_s \|\beta\|_{W^{s,\infty}}$ , see (8.2). The bound  $\|\chi_j(\tau)\|_s \leq_s 1 + \|\beta\|_{W^{s+1,\infty}}$  follows by (8.15). The tame estimates for  $g_j(\tau)$  defined in (8.13) are more difficult because require tame estimates for the adjoint  $(\Phi^\tau)^T$ ,  $\forall \tau \in [0, 1]$ . The adjoint of the flow map can be represented as the flow map of the ‘‘adjoint’’ PDE

$$\partial_\tau z = \Pi_S^\perp \{b(\tau, x) \partial_x \Pi_S^\perp z\} = b(\tau, x) \partial_x z - \Pi_S (b(\tau, x) \partial_x z), \quad z \in H_S^\perp, \quad (8.16)$$

where  $-\Pi_S^\perp b(\tau, x) \partial_x$  is the  $L^2$ -adjoint of the Hamiltonian vector field in (8.5). We denote by  $\Psi^{\tau_0, \tau}$  the flow of (8.16), namely  $\Psi^{\tau_0, \tau}(v)$  is the solution of (8.16) with  $\Psi^{\tau_0, \tau_0}(v) = v$ . Since the derivative  $\partial_\tau(\Phi^\tau(u_0), \Psi^{\tau_0, \tau}(v))_{L^2(\mathbb{T})} = 0, \forall \tau$ , we deduce that  $(\Phi^{\tau_0}(u_0), \Psi^{\tau_0, \tau_0}(v))_{L^2(\mathbb{T})} = (\Phi^0(u_0), \Psi^{\tau_0, 0}(v))_{L^2(\mathbb{T})}$ , namely (recall that  $\Psi^{\tau_0, \tau_0}(v) = v$ ) the adjoint

$$(\Phi^{\tau_0})^T = \Psi^{\tau_0, 0}, \quad \forall \tau_0 \in [0, 1]. \quad (8.17)$$

Thus it is sufficient to prove tame estimates for the flow  $\Psi^{\tau_0, \tau}$ . We first provide a useful expression for the solution  $z(\tau, x) := \Psi^{\tau_0, \tau}(v)$  of (8.16), obtained by the methods of characteristics. Let  $\gamma^{\tau_0, \tau}(y)$  be the flow of (8.4). Since  $\partial_\tau z(\tau, \gamma^{\tau_0, \tau}(y)) = -[\Pi_S(b(\tau) \partial_x z(\tau))](\gamma^{\tau_0, \tau}(y))$  we get

$$z(\tau, \gamma^{\tau_0, \tau}(y)) = v(y) + \int_\tau^{\tau_0} [\Pi_S(b(s) \partial_x z(s))](\gamma^{\tau_0, s}(y)) ds, \quad \forall \tau \in [0, 1].$$

Denoting by  $y = x + \sigma(\tau, x)$  the inverse diffeomorphism of  $x = \gamma^{\tau_0, \tau}(y) = y + \tilde{\sigma}(\tau, y)$ , we get

$$\begin{aligned} \Psi^{\tau_0, \tau}(v) &= z(\tau, x) = v(x + \sigma(\tau, x)) + \int_\tau^{\tau_0} [\Pi_S(b(s) \partial_x z(s))](\gamma^{\tau_0, s}(x + \sigma(\tau, x))) ds \\ &= v(x + \sigma(\tau, x)) + \int_\tau^{\tau_0} \sum_{j \in S} (z(s), p_j(s)) \kappa_j(s, x) ds = v(x + \sigma(\tau, x)) + \mathcal{R}_\tau v, \end{aligned} \quad (8.18)$$

where  $p_j(s) := -\partial_x(b(s)e^{ijx})$ ,  $\kappa_j(s, x) := e^{ij\gamma^{\tau_0, s}(x + \sigma(\tau, x))}$  and

$$(\mathcal{R}_\tau v)(x) := \int_\tau^{\tau_0} \sum_{j \in S} (\Psi^{\tau_0, s}(v), p_j(s))_{L^2(\mathbb{T})} \kappa_j(s, x) ds.$$

Since  $\|\sigma(\tau, \cdot)\|_{W^{s, \infty}}, \|\tilde{\sigma}(\tau, \cdot)\|_{W^{s, \infty}} \leq_s \|\beta\|_{W^{s+1, \infty}}$  (recall also (8.3)), we derive  $\|p_j\|_s \leq_s \|\beta\|_{W^{s+2, \infty}}, \|\kappa_j\|_s \leq_s 1 + \|\beta\|_{W^{s+1, \infty}}$  and  $\|v(x + \sigma(\tau, x))\|_s \leq_s \|v\|_s + \|\beta\|_{W^{s+1, \infty}} \|v\|_{s_0}, \forall \tau \in [0, 1]$ . Moreover

$$\|\mathcal{R}_\tau v\|_s \leq_s \sup_{\tau \in [0, 1]} \|\Psi^{\tau_0, \tau}(v)\|_s \|\beta\|_{W^{s_0+2, \infty}} + \sup_{\tau \in [0, 1]} \|\Psi^{\tau_0, \tau}(v)\|_{s_0} \|\beta\|_{W^{s+2, \infty}}.$$

Therefore, for all  $\tau \in [0, 1]$ ,

$$\|\Psi^{\tau_0, \tau} v\|_s \leq_s \|v\|_s + \|\beta\|_{W^{s+1, \infty}} \|v\|_{s_0} + \sup_{\tau \in [0, 1]} \{ \|\Psi^{\tau_0, \tau} v\|_s \|\beta\|_{W^{s_0+2, \infty}} + \|\Psi^{\tau_0, \tau} v\|_{s_0} \|\beta\|_{W^{s+2, \infty}} \}. \quad (8.19)$$

For  $s = s_0$  it implies

$$\sup_{\tau \in [0, 1]} \|\Psi^{\tau_0, \tau}(v)\|_{s_0} \leq_{s_0} \|v\|_{s_0} (1 + \|\beta\|_{W^{s_0+1, \infty}}) + \sup_{\tau \in [0, 1]} \|\Psi^{\tau_0, \tau}(v)\|_{s_0} \|\beta\|_{W^{s_0+2, \infty}}$$

and so, for  $\|\beta\|_{W^{s_0+2, \infty}} \leq c(s_0)$  small enough,

$$\sup_{\tau \in [0, 1]} \|\Psi^{\tau_0, \tau}(v)\|_{s_0} \leq_{s_0} \|v\|_{s_0}. \quad (8.20)$$

Finally (8.19), (8.20) imply the tame estimate

$$\sup_{\tau \in [0, 1]} \|\Psi^{\tau_0, \tau}(v)\|_s \leq_s \|v\|_s + \|\beta\|_{W^{s+2, \infty}} \|v\|_{s_0}. \quad (8.21)$$

By (8.17) and (8.21) we deduce the bound (8.7) for  $g_j$  defined in (8.13). The tame estimate (8.8) for  $\Phi$  follows by that of  $\mathcal{A}$  and (8.7) (use Lemma 2.4). The estimate for  $\Phi^{-1}$  follows in the same way because  $\Phi^{-1} = \Phi^{1, 0}$  is the backward flow.  $\square$

We conjugate  $\mathcal{L}_\omega$  in (7.34) via the symplectic map  $\Phi = \mathcal{A}_\perp + \mathcal{R}_\Phi$  of Lemma 8.2. We compute (split  $\Pi_S^\perp = I - \Pi_S$ )

$$\mathcal{L}_\omega \Phi = \Phi \mathcal{D}_\omega + \Pi_S^\perp \mathcal{A}(b_3 \partial_{yyy} + b_2 \partial_{yy} + b_1 \partial_y + b_0) \Pi_S^\perp + \mathcal{R}_I, \quad (8.22)$$

where the coefficients are

$$b_3(\varphi, y) := \mathcal{A}^T [a_1(1 + \beta_x)^3] \quad b_2(\varphi, y) := \mathcal{A}^T [2(a_1)_x(1 + \beta_x)^2 + 6a_1\beta_{xx}(1 + \beta_x)] \quad (8.23)$$

$$b_1(\varphi, y) := \mathcal{A}^T \left[ (\mathcal{D}_\omega \beta) + 3a_1 \frac{\beta_{xx}^2}{1 + \beta_x} + 4a_1\beta_{xxx} + 6(a_1)_x\beta_{xx} + (a_1)_{xx}(1 + \beta_x) + a_0(1 + \beta_x) \right] \quad (8.24)$$

$$b_0(\varphi, y) := \mathcal{A}^T \left[ \frac{(\mathcal{D}_\omega \beta_x)}{1 + \beta_x} + a_1 \frac{\beta_{xxxx}}{1 + \beta_x} + 2(a_1)_x \frac{\beta_{xxx}}{1 + \beta_x} + (a_1)_{xx} \frac{\beta_{xx}}{1 + \beta_x} + a_0 \frac{\beta_{xx}}{1 + \beta_x} + (a_0)_x \right] \quad (8.25)$$

and the remainder

$$\begin{aligned} \mathcal{R}_I := & -\Pi_S^\perp \partial_x (\varepsilon^2 \mathcal{R}_2 + \mathcal{R}_*) \mathcal{A}_\perp - \Pi_S^\perp (a_1 \partial_{xxx} + 2(a_1)_x \partial_{xx} + ((a_1)_{xx} + a_0) \partial_x + (a_0)_x) \Pi_S \mathcal{A} \Pi_S^\perp \\ & + [\mathcal{D}_\omega, \mathcal{R}_\Phi] + (\mathcal{L}_\omega - \mathcal{D}_\omega) \mathcal{R}_\Phi. \end{aligned} \quad (8.26)$$

The commutator  $[\mathcal{D}_\omega, \mathcal{R}_\Phi]$  has the form (8.6) with  $\mathcal{D}_\omega g_j$  or  $\mathcal{D}_\omega \chi_j$ ,  $\mathcal{D}_\omega \psi_j$  instead of  $\chi_j$ ,  $g_j$ ,  $\psi_j$  respectively. Also the last term  $(\mathcal{L}_\omega - \mathcal{D}_\omega) \mathcal{R}_\Phi$  in (8.26) has the form (8.6) (note that  $\mathcal{L}_\omega - \mathcal{D}_\omega$  does not contain derivatives with respect to  $\varphi$ ). By (8.22), and decomposing  $I = \Pi_S + \Pi_S^\perp$ , we get

$$\mathcal{L}_\omega \Phi = \Phi (\mathcal{D}_\omega + b_3 \partial_{yyy} + b_2 \partial_{yy} + b_1 \partial_y + b_0) \Pi_S^\perp + \mathcal{R}_{II}, \quad (8.27)$$

$$\mathcal{R}_{II} := \{ \Pi_S^\perp (\mathcal{A} - I) \Pi_S - \mathcal{R}_\Phi \} (b_3 \partial_{yyy} + b_2 \partial_{yy} + b_1 \partial_y + b_0) \Pi_S^\perp + \mathcal{R}_I. \quad (8.28)$$

Now we choose the function  $\beta = \beta(\varphi, x)$  such that

$$a_1(\varphi, x)(1 + \beta_x(\varphi, x))^3 = b_3(\varphi) \quad (8.29)$$

so that the coefficient  $b_3$  in (8.23) depends only on  $\varphi$  (note that  $\mathcal{A}^T [b_3(\varphi)] = b_3(\varphi)$ ). The only solution of (8.29) with zero space average is (see e.g. [2]-section 3.1)

$$\beta := \partial_x^{-1} \rho_0, \quad \rho_0 := b_3(\varphi)^{1/3} (a_1(\varphi, x))^{-1/3} - 1, \quad b_3(\varphi) := \left( \frac{1}{2\pi} \int_{\mathbb{T}} (a_1(\varphi, x))^{-1/3} dx \right)^{-3}. \quad (8.30)$$

Applying the symplectic map  $\Phi^{-1}$  in (8.27) we obtain the Hamiltonian operator (see Definition 2.2)

$$\mathcal{L}_1 := \Phi^{-1} \mathcal{L}_\omega \Phi = \Pi_S^\perp (\omega \cdot \partial_\varphi + b_3(\varphi) \partial_{yyy} + b_1 \partial_y + b_0) \Pi_S^\perp + \mathfrak{R}_1 \quad (8.31)$$

where  $\mathfrak{R}_1 := \Phi^{-1} \mathcal{R}_{II}$ . We used that, by the Hamiltonian nature of  $\mathcal{L}_1$ , the coefficient  $b_2 = 2(b_3)_y$  (see [2]-Remark 3.5) and so, by the choice (8.30), we have  $b_2 = 2(b_3)_y = 0$ . In the next Lemma we analyse the structure of the remainder  $\mathfrak{R}_1$ .

**Lemma 8.3.** *The operator  $\mathfrak{R}_1$  has the form (7.7).*

*Proof.* The remainders  $\mathcal{R}_I$  and  $\mathcal{R}_{II}$  have the form (7.7). Indeed  $\mathcal{R}_2, \mathcal{R}_*$  in (8.26) have the form (7.2) (see Proposition 7.6) and the term  $\Pi_S \mathcal{A} w = \sum_{j \in S} (\mathcal{A}^T e^{ijx}, w)_{L^2(\mathbb{T})} e^{ijx}$  has the same form. By (8.6), the terms of  $\mathcal{R}_I, \mathcal{R}_{II}$  which involves the operator  $\mathcal{R}_\Phi$  have the form (7.7). All the operations involved preserve this structure: if  $R_\tau w = \chi(\tau)(w, g(\tau))_{L^2(\mathbb{T})}$ ,  $\tau \in [0, 1]$ , then

$$\begin{aligned} R_\tau \Pi_S^\perp w &= \chi(\tau) (\Pi_S^\perp g(\tau), w)_{L^2(\mathbb{T})}, \quad R_\tau \mathcal{A} w = \chi(\tau) (\mathcal{A}^T g(\tau), w)_{L^2(\mathbb{T})}, \quad \partial_x R_\tau w = \chi_x(\tau) (g(\tau), w)_{L^2(\mathbb{T})}, \\ \Pi_S^\perp R_\tau w &= (\Pi_S^\perp \chi(\tau)) (g(\tau), w)_{L^2(\mathbb{T})}, \quad \mathcal{A} R_\tau w = (\mathcal{A} \chi(\tau)) (g(\tau), w)_{L^2(\mathbb{T})}, \quad \Phi^{-1} R_\tau w = (\Phi^{-1} \chi(\tau)) (g(\tau), w)_{L^2(\mathbb{T})} \end{aligned}$$

(the last equality holds because  $\Phi^{-1}(f(\varphi)w) = f(\varphi)\Phi^{-1}(w)$  for all function  $f(\varphi)$ ). Hence  $\mathfrak{R}_1$  has the form (7.7) where  $\chi_j(\tau) \in H_S^\perp$  for all  $\tau \in [0, 1]$ .  $\square$

We now put in evidence the terms of order  $\varepsilon, \varepsilon^2, \dots$ , in  $b_1, b_0, \mathfrak{R}_1$ , recalling that  $a_1 - 1 = O(\varepsilon^3)$  (see (7.38)),  $a_0 = O(\varepsilon)$  (see (7.35)-(7.39)), and  $\beta = O(\varepsilon^3)$  (proved below in (8.35)). We expand  $b_1$  in (8.24) as

$$b_1 = -\varepsilon p_1 - \varepsilon^2 p_2 - q_{>2} + \mathcal{D}_\omega \beta + 4\beta_{xxx} + (a_1)_{xx} + b_{1, \geq 4} \quad (8.32)$$

where  $b_{1, \geq 4} = O(\varepsilon^4)$  is defined by difference (the precise estimate is in Lemma 8.5).

**Remark 8.4.** The function  $\mathcal{D}_\omega \beta$  has zero average in  $x$  by (8.30) as well as  $(a_1)_{xx}, \beta_{xxx}$ .  $\square$

Similarly, we expand  $b_0$  in (8.25) as

$$b_0 = -\varepsilon(p_1)_x - \varepsilon^2(p_2)_x - (q_{>2})_x + \mathcal{D}_\omega \beta_x + \beta_{xxxx} + b_{0,\geq 4} \quad (8.33)$$

where  $b_{0,\geq 4} = O(\varepsilon^4)$  is defined by difference.

Using the equalities (8.28), (8.26) and  $\Pi_S \mathcal{A} \Pi_S^\perp = \Pi_S (\mathcal{A} - I) \Pi_S^\perp$  we get

$$\mathfrak{R}_1 := \Phi^{-1} \mathcal{R}_{II} = -\varepsilon^2 \Pi_S^\perp \partial_x \mathcal{R}_2 + \mathcal{R}_* \quad (8.34)$$

where  $\mathcal{R}_2$  is defined in (7.29) and we have renamed  $\mathcal{R}_*$  the term of order  $o(\varepsilon^2)$  in  $\mathfrak{R}_1$ . The remainder  $\mathcal{R}_*$  in (8.34) has the form (7.7).

**Lemma 8.5.** *There is  $\sigma = \sigma(\tau, \nu) > 0$  such that*

$$\|\beta\|_s^{\text{Lip}(\gamma)} \leq_s \varepsilon^3 (1 + \|\mathfrak{J}_\delta\|_{s+1}^{\text{Lip}(\gamma)}), \quad \|\partial_i \beta[\tilde{v}]\|_s \leq_s \varepsilon^3 (\|\tilde{v}\|_{s+\sigma} + \|\mathfrak{J}_\delta\|_{s+\sigma} \|\tilde{v}\|_{s_0+\sigma}), \quad (8.35)$$

$$\|b_3 - 1\|_s^{\text{Lip}(\gamma)} \leq_s \varepsilon^4 + \varepsilon^{b+2} \|\mathfrak{J}_\delta\|_{s+\sigma}^{\text{Lip}(\gamma)}, \quad \|\partial_i b_3[\tilde{v}]\|_s \leq_s \varepsilon^{b+2} (\|\tilde{v}\|_{s+\sigma} + \|\mathfrak{J}_\delta\|_{s+\sigma} \|\tilde{v}\|_{s_0+\sigma}) \quad (8.36)$$

$$\|b_{1,\geq 4}\|_s^{\text{Lip}(\gamma)} + \|b_{0,\geq 4}\|_s^{\text{Lip}(\gamma)} \leq_s \varepsilon^4 + \varepsilon^{b+2} \|\mathfrak{J}_\delta\|_{s+\sigma}^{\text{Lip}(\gamma)} \quad (8.37)$$

$$\|\partial_i b_{1,\geq 4}[\tilde{v}]\|_s + \|\partial_i b_{0,\geq 4}[\tilde{v}]\|_s \leq_s \varepsilon^{b+2} (\|\tilde{v}\|_{s+\sigma} + \|\mathfrak{J}_\delta\|_{s+\sigma} \|\tilde{v}\|_{s_0+\sigma}). \quad (8.38)$$

The transformations  $\Phi, \Phi^{-1}$  satisfy

$$\|\Phi^{\pm 1} h\|_s^{\text{Lip}(\gamma)} \leq_s \|h\|_{s+1}^{\text{Lip}(\gamma)} + \|\mathfrak{J}_\delta\|_{s+\sigma}^{\text{Lip}(\gamma)} \|h\|_{s_0+1}^{\text{Lip}(\gamma)} \quad (8.39)$$

$$\|\partial_i(\Phi^{\pm 1} h)[\tilde{v}]\|_s \leq_s \|h\|_{s+\sigma} \|\tilde{v}\|_{s_0+\sigma} + \|h\|_{s_0+\sigma} \|\tilde{v}\|_{s+\sigma} + \|\mathfrak{J}_\delta\|_{s+\sigma} \|h\|_{s_0+\sigma} \|\tilde{v}\|_{s_0+\sigma}. \quad (8.40)$$

Moreover the remainder  $\mathcal{R}_*$  has the form (7.7), where the functions  $\chi_j(\tau), g_j(\tau)$  satisfy the estimates (7.41)-(7.42) uniformly in  $\tau \in [0, 1]$ .

*Proof.* The estimates (8.35) follow by (8.30), (7.38), and the usual interpolation and tame estimates in Lemmata 2.2-2.4 (and Lemma 5.13) and (7.8). For the estimates of  $b_3$ , by (8.30) and (7.35) we consider the function  $r_1$  defined in (7.25). Recalling also (3.4) and (7.11), the function

$$r_1(T_\delta) = \varepsilon^3 (\partial_{u_x u_x} f_5)(v_\delta, (v_\delta)_x) + r_{1,\geq 4}, \quad r_{1,\geq 4} := r_1(T_\delta) - \varepsilon^3 (\partial_{u_x u_x} f_5)(v_\delta, (v_\delta)_x).$$

Hypothesis (S1) implies, as in the proof of Lemma 7.5, that the space average  $\int_{\mathbb{T}} (\partial_{u_x u_x} f_5)(v_\delta, (v_\delta)_x) dx = 0$ . Hence the bound (8.36) for  $b_3 - 1$  follows. For the estimates on  $\Phi, \Phi^{-1}$  we apply Lemma 8.2 and the estimate (8.35) for  $\beta$ . We estimate the remainder  $\mathcal{R}_*$  in (8.34), using (8.26), (8.28) and (7.41)-(7.42).  $\square$

## 8.2 Reparametrization of time

The goal of this section is to make constant the coefficient of the highest order spatial derivative operator  $\partial_{yyy}$ , by a quasi-periodic reparametrization of time. We consider the change of variable

$$(Bw)(\varphi, y) := w(\varphi + \omega\alpha(\varphi), y), \quad (B^{-1}h)(\vartheta, y) := h(\vartheta + \omega\tilde{\alpha}(\vartheta), y),$$

where  $\varphi = \vartheta + \omega\tilde{\alpha}(\vartheta)$  is the inverse diffeomorphism of  $\vartheta = \varphi + \omega\alpha(\varphi)$  in  $\mathbb{T}^\nu$ . By conjugation, the differential operators become

$$B^{-1}\omega \cdot \partial_\varphi B = \rho(\vartheta)\omega \cdot \partial_\vartheta, \quad B^{-1}\partial_y B = \partial_y, \quad \rho := B^{-1}(1 + \omega \cdot \partial_\varphi \alpha). \quad (8.41)$$

By (8.31), using also that  $B$  and  $B^{-1}$  commute with  $\Pi_S^\perp$ , we get

$$B^{-1}\mathcal{L}_1 B = \Pi_S^\perp [\rho\omega \cdot \partial_\vartheta + (B^{-1}b_3)\partial_{yyy} + (B^{-1}b_1)\partial_y + (B^{-1}b_0)]\Pi_S^\perp + B^{-1}\mathfrak{R}_1 B. \quad (8.42)$$

We choose  $\alpha$  such that

$$(B^{-1}b_3)(\vartheta) = m_3\rho(\vartheta), \quad m_3 \in \mathbb{R}, \quad \text{namely } b_3(\varphi) = m_3(1 + \omega \cdot \partial_\varphi \alpha(\varphi)) \quad (8.43)$$

(recall (8.41)). The unique solution with zero average of (8.43) is

$$\alpha(\varphi) := \frac{1}{m_3}(\omega \cdot \partial_\varphi)^{-1}(b_3 - m_3)(\varphi), \quad m_3 := \frac{1}{(2\pi)^\nu} \int_{\mathbb{T}^\nu} b_3(\varphi) d\varphi. \quad (8.44)$$

Hence, by (8.42),

$$B^{-1}\mathcal{L}_1 B = \rho\mathcal{L}_2, \quad \mathcal{L}_2 := \Pi_S^\perp(\omega \cdot \partial_\vartheta + m_3\partial_{yyy} + c_1\partial_y + c_0)\Pi_S^\perp + \mathfrak{R}_2 \quad (8.45)$$

$$c_1 := \rho^{-1}(B^{-1}b_1), \quad c_0 := \rho^{-1}(B^{-1}b_0), \quad \mathfrak{R}_2 := \rho^{-1}B^{-1}\mathfrak{R}_1 B. \quad (8.46)$$

The transformed operator  $\mathcal{L}_2$  in (8.45) is still Hamiltonian, since the reparametrization of time preserves the Hamiltonian structure (see Section 2.2 and Remark 3.7 in [2]).

We now put in evidence the terms of order  $\varepsilon, \varepsilon^2, \dots$  in  $c_1, c_0$ . To this aim, we anticipate the following estimates:  $\rho(\vartheta) = 1 + O(\varepsilon^4)$ ,  $\alpha = O(\varepsilon^4\gamma^{-1})$ ,  $m_3 = 1 + O(\varepsilon^4)$ ,  $B^{-1} - I = O(\alpha)$  (in low norm), which are proved in Lemma 8.7 below. Then, by (8.32)-(8.33), we expand the functions  $c_1, c_0$  in (8.46) as

$$c_1 = -\varepsilon p_1 - \varepsilon^2 p_2 - B^{-1}q_{>2} + \varepsilon(p_1 - B^{-1}p_1) + \varepsilon^2(p_2 - B^{-1}p_2) + \mathcal{D}_\omega\beta + 4\beta_{xxx} + (a_1)_{xx} + c_{1,\geq 4}, \quad (8.47)$$

$$c_0 = -\varepsilon(p_1)_x - \varepsilon^2(p_2)_x - (B^{-1}q_{>2})_x + \varepsilon(p_1 - B^{-1}p_1)_x + \varepsilon^2(p_2 - B^{-1}p_2)_x + (\mathcal{D}_\omega\beta)_x + \beta_{xxx} + c_{0,\geq 4}, \quad (8.48)$$

where  $c_{1,\geq 4}, c_{0,\geq 4} = O(\varepsilon^4)$  are defined by difference.

**Remark 8.6.** The functions  $\varepsilon(p_1 - B^{-1}p_1) = O(\varepsilon^5\gamma^{-1})$  and  $\varepsilon^2(p_2 - B^{-1}p_2) = O(\varepsilon^6\gamma^{-1})$ , see (8.53). For the reducibility scheme, the terms of order  $\partial_x^0$  with size  $O(\varepsilon^5\gamma^{-1})$  are perturbative, since  $\varepsilon^5\gamma^{-2} \ll 1$ .  $\square$

The remainder  $\mathfrak{R}_2$  in (8.46) has still the form (7.7) and, by (8.34),

$$\mathfrak{R}_2 := -\rho^{-1}B^{-1}\mathfrak{R}_1 B = -\varepsilon^2\Pi_S^\perp\partial_x\mathcal{R}_2 + \mathcal{R}_* \quad (8.49)$$

where  $\mathcal{R}_2$  is defined in (7.29) and we have renamed  $\mathcal{R}_*$  the term of order  $o(\varepsilon^2)$  in  $\mathfrak{R}_2$ .

**Lemma 8.7.** *There is  $\sigma = \sigma(\nu, \tau) > 0$  (possibly larger than  $\sigma$  in Lemma 8.5) such that*

$$|m_3 - 1|^{\text{Lip}(\gamma)} \leq C\varepsilon^4, \quad |\partial_i m_3[\hat{\eta}]| \leq C\varepsilon^{b+2}\|\hat{\eta}\|_{s_0+\sigma} \quad (8.50)$$

$$\|\alpha\|_s^{\text{Lip}(\gamma)} \leq_s \varepsilon^4\gamma^{-1} + \varepsilon^{b+2}\gamma^{-1}\|\mathfrak{J}_\delta\|_{s+\sigma}^{\text{Lip}(\gamma)}, \quad \|\partial_i\alpha[\hat{\eta}]\|_s \leq_s \varepsilon^{b+2}\gamma^{-1}(\|\hat{\eta}\|_{s+\sigma} + \|\mathfrak{J}_\delta\|_{s+\sigma}\|\hat{\eta}\|_{s_0+\sigma}), \quad (8.51)$$

$$\|\rho - 1\|_s^{\text{Lip}(\gamma)} \leq_s \varepsilon^4 + \varepsilon^{b+2}\|\mathfrak{J}_\delta\|_{s+\sigma}^{\text{Lip}(\gamma)}, \quad \|\partial_i\rho[\hat{\eta}]\|_s \leq_s \varepsilon^{b+2}(\|\hat{\eta}\|_{s+\sigma} + \|\mathfrak{J}_\delta\|_{s+\sigma}\|\hat{\eta}\|_{s_0+\sigma}) \quad (8.52)$$

$$\|p_k - B^{-1}p_k\|_s^{\text{Lip}(\gamma)} \leq_s \varepsilon^4\gamma^{-1} + \varepsilon^{b+2}\gamma^{-1}\|\mathfrak{J}_\delta\|_{s+\sigma}^{\text{Lip}(\gamma)}, \quad k = 1, 2 \quad (8.53)$$

$$\|\partial_i(p_k - B^{-1}p_k)[\hat{\eta}]\|_s \leq_s \varepsilon^{b+2}\gamma^{-1}(\|\hat{\eta}\|_{s+\sigma} + \|\mathfrak{J}_\delta\|_{s+\sigma}\|\hat{\eta}\|_{s_0+\sigma}) \quad (8.54)$$

$$\|B^{-1}q_{>2}\|_s^{\text{Lip}(\gamma)} \leq_s \varepsilon^3 + \varepsilon^b\|\mathfrak{J}_\delta\|_{s+\sigma}^{\text{Lip}(\gamma)}, \quad (8.55)$$

$$\|\partial_i(B^{-1}q_{>2})[\hat{\eta}]\|_s \leq_s \varepsilon^b(\|\hat{\eta}\|_{s+\sigma} + \|\mathfrak{J}_\delta\|_{s+\sigma}\|\hat{\eta}\|_{s_0+\sigma}). \quad (8.56)$$

The terms  $c_{1,\geq 4}, c_{0,\geq 4}$  satisfy the bounds (8.37)-(8.38). The transformations  $B, B^{-1}$  satisfy the estimates (8.39), (8.40). The remainder  $\mathcal{R}_*$  has the form (7.7), and the functions  $g_j(\tau), \chi_j(\tau)$  satisfy the estimates (7.41)-(7.42) for all  $\tau \in [0, 1]$ .

*Proof.* (8.50) follows from (8.44), (8.36). The estimate  $\|\alpha\|_s \leq_s \varepsilon^4\gamma^{-1} + \varepsilon^{b+2}\gamma^{-1}\|\mathfrak{J}_\delta\|_{s+\sigma}$  and the inequality for  $\partial_i\alpha$  in (8.51) follow by (8.44), (8.36), (8.50). For the first bound in (8.51) we also differentiate (8.44) with respect to the parameter  $\omega$ . The estimates for  $\rho$  follow from  $\rho - 1 = B^{-1}(b_3 - m_3)/m_3$ .  $\square$



### 8.3 Translation of the space variable

In view of the next linear Birkhoff normal form steps (whose goal is to eliminate the terms of size  $\varepsilon$  and  $\varepsilon^2$ ), in the expressions (8.47), (8.48) we split  $p_1 = \bar{p}_1 + (p_1 - \bar{p}_1)$ ,  $p_2 = \bar{p}_2 + (p_2 - \bar{p}_2)$  (see (7.35)), where

$$\bar{p}_1 := 6\bar{v}, \quad \bar{p}_2 := 6\pi_0[(\partial_x^{-1}\bar{v})^2], \quad \bar{v}(\varphi, x) := \sum_{j \in S} \sqrt{\xi_j} e^{i\ell(j) \cdot \varphi} e^{ijx}, \quad (8.57)$$

and  $\ell : S \rightarrow \mathbb{Z}^\nu$  is the odd injective map (see (1.8))

$$\ell : S \rightarrow \mathbb{Z}^\nu, \quad \ell(\bar{j}_i) := e_i, \quad \ell(-\bar{j}_i) := -\ell(\bar{j}_i) = -e_i, \quad i = 1, \dots, \nu, \quad (8.58)$$

denoting by  $e_i = (0, \dots, 1, \dots, 0)$  the  $i$ -th vector of the canonical basis of  $\mathbb{R}^\nu$ .

**Remark 8.8.** All the functions  $\bar{p}_1, \bar{p}_2, p_1 - \bar{p}_1, p_2 - \bar{p}_2$  have zero average in  $x$ .  $\square$

We write the variable coefficients  $c_1, c_0$  of the operator  $\mathcal{L}_2$  in (8.45) (see (8.47), (8.48)) as

$$c_1 = -\varepsilon\bar{p}_1 - \varepsilon^2\bar{p}_2 + q_{c_1} + c_{1, \geq 4}, \quad c_0 = -\varepsilon(\bar{p}_1)_x - \varepsilon^2(\bar{p}_2)_x + q_{c_0} + c_{0, \geq 4}, \quad (8.59)$$

where we define

$$q_{c_1} := q + 4\beta_{xxx} + (a_1)_{xx}, \quad q_{c_0} := q_x + \beta_{xxxx}, \quad (8.60)$$

$$q := \varepsilon(p_1 - B^{-1}p_1) + \varepsilon(\bar{p}_1 - p_1) + \varepsilon^2(p_2 - B^{-1}p_2) + \varepsilon^2(\bar{p}_2 - p_2) - B^{-1}q_{>2} + \mathcal{D}_\omega\beta. \quad (8.61)$$

**Remark 8.9.** The functions  $q_{c_1}, q_{c_0}$  have zero average in  $x$  (see Remarks 8.8, 8.4 and Lemma 7.5).  $\square$

**Lemma 8.10.** *The functions  $\bar{p}_k - p_k$ ,  $k = 1, 2$  and  $q_{c_m}$ ,  $m = 0, 1$ , satisfy*

$$\|\bar{p}_k - p_k\|_s^{\text{Lip}(\gamma)} \leq_s \|\mathfrak{J}_\delta\|_s^{\text{Lip}(\gamma)}, \quad \|\partial_i(\bar{p}_k - p_k)[\hat{v}]\|_s \leq_s \|\hat{v}\|_s + \|\mathfrak{J}_\delta\|_s \|\hat{v}\|_{s_0}, \quad (8.62)$$

$$\|q_{c_m}\|_s^{\text{Lip}(\gamma)} \leq_s \varepsilon^5 \gamma^{-1} + \varepsilon \|\mathfrak{J}_\delta\|_{s+\sigma}^{\text{Lip}(\gamma)}, \quad \|\partial_i q_{c_m}[\hat{v}]\|_s^{\text{Lip}(\gamma)} \leq_s \varepsilon (\|\hat{v}\|_{s+\sigma} + \|\mathfrak{J}_\delta\|_{s+\sigma} \|\hat{v}\|_{s_0+\sigma}). \quad (8.63)$$

*Proof.* The bound (8.62) follows from (8.57), (7.35), (7.11), (7.8). Then use (8.62), (8.53)-(8.56), (8.35), (7.38) to prove (8.63). The biggest term comes from  $\varepsilon(\bar{p}_1 - p_1)$ .  $\square$

We now apply the transformation  $\mathcal{T}$  defined in (8.64) whose goal is to remove the space average from the coefficient in front of  $\partial_y$ .

Consider the change of the space variable  $z = y + p(\vartheta)$  which induces on  $H_{S^\perp}^s(\mathbb{T}^{\nu+1})$  the operators

$$(\mathcal{T}w)(\vartheta, y) := w(\vartheta, y + p(\vartheta)), \quad (\mathcal{T}^{-1}h)(\vartheta, z) = h(\vartheta, z - p(\vartheta)) \quad (8.64)$$

(which are a particular case of those used in section 8.1). The differential operator becomes  $\mathcal{T}^{-1}\omega \cdot \partial_\vartheta \mathcal{T} = \omega \cdot \partial_\vartheta + \{\omega \cdot \partial_\vartheta p(\vartheta)\} \partial_z$ ,  $\mathcal{T}^{-1} \partial_y \mathcal{T} = \partial_z$ . Since  $\mathcal{T}, \mathcal{T}^{-1}$  commute with  $\Pi_S^\perp$ , we get

$$\mathcal{L}_3 := \mathcal{T}^{-1} \mathcal{L}_2 \mathcal{T} = \Pi_S^\perp (\omega \cdot \partial_\vartheta + m_3 \partial_{zzz} + d_1 \partial_z + d_0) \Pi_S^\perp + \mathfrak{R}_3, \quad (8.65)$$

$$d_1 := (\mathcal{T}^{-1} c_1) + \omega \cdot \partial_\vartheta p, \quad d_0 := \mathcal{T}^{-1} c_0, \quad \mathfrak{R}_3 := \mathcal{T}^{-1} \mathfrak{R}_2 \mathcal{T}. \quad (8.66)$$

We choose

$$m_1 := \frac{1}{(2\pi)^{\nu+1}} \int_{\mathbb{T}^{\nu+1}} c_1 d\vartheta dy, \quad p := (\omega \cdot \partial_\vartheta)^{-1} \left( m_1 - \frac{1}{2\pi} \int_{\mathbb{T}} c_1 dy \right), \quad (8.67)$$

so that  $\frac{1}{2\pi} \int_{\mathbb{T}} d_1(\vartheta, z) dz = m_1$  for all  $\vartheta \in \mathbb{T}^\nu$ . Note that, by (8.59),

$$\int_{\mathbb{T}} c_1(\vartheta, y) dy = \int_{\mathbb{T}} c_{1, \geq 4}(\vartheta, y) dy, \quad \omega \cdot \partial_\vartheta p(\vartheta) = m_1 - \frac{1}{2\pi} \int_{\mathbb{T}} c_{1, \geq 4}(\vartheta, y) dy \quad (8.68)$$

because  $\bar{p}_1, \bar{p}_2, q_{c_1}$  have all zero space-average. Also note that  $\mathfrak{R}_3$  has the form (7.7). Since  $\mathcal{T}$  is symplectic, the operator  $\mathcal{L}_3$  in (8.65) is Hamiltonian.

**Remark 8.11.** We require Hypothesis (S1) so that the function  $q_{>2}$  has zero space average (see Lemma 7.5). If  $q_{>2}$  did not have zero average, then  $p$  in (8.67) would have size  $O(\varepsilon^3\gamma^{-1})$  (see (7.31)) and, since  $\mathcal{T}^{-1} - I = O(\varepsilon^3\gamma^{-1})$ , the function  $\tilde{d}_0$  in (8.71) would satisfy  $\tilde{d}_0 = O(\varepsilon^4\gamma^{-1})$ . Therefore it would remain a term of order  $\partial_x^0$  which is not perturbative for the reducibility scheme of section 8.7.  $\square$

We put in evidence the terms of size  $\varepsilon, \varepsilon^2$  in  $d_0, d_1, \mathfrak{R}_3$ . Recalling (8.66), (8.59), we split

$$d_1 = -\varepsilon\bar{p}_1 - \varepsilon^2\bar{p}_2 + \tilde{d}_1, \quad d_0 = -\varepsilon(\bar{p}_1)_x - \varepsilon^2(\bar{p}_2)_x + \tilde{d}_0, \quad \mathfrak{R}_3 = -\varepsilon^2\Pi_S^\perp\partial_x\bar{\mathcal{R}}_2 + \tilde{\mathcal{R}}_* \quad (8.69)$$

where  $\bar{\mathcal{R}}_2$  is obtained replacing  $v_\delta$  with  $\bar{v}$  in  $\mathcal{R}_2$  (see (7.29)), and

$$\tilde{d}_1 := \varepsilon(\bar{p}_1 - \mathcal{T}^{-1}\bar{p}_1) + \varepsilon^2(\bar{p}_2 - \mathcal{T}^{-1}\bar{p}_2) + \mathcal{T}^{-1}(q_{c_1} + c_{1,\geq 4}) + \omega \cdot \partial_\vartheta p, \quad (8.70)$$

$$\tilde{d}_0 := \varepsilon(\bar{p}_1 - \mathcal{T}^{-1}\bar{p}_1)_x + \varepsilon^2(\bar{p}_2 - \mathcal{T}^{-1}\bar{p}_2)_x + \mathcal{T}^{-1}(q_{c_0} + c_{0,\geq 4}), \quad (8.71)$$

$$\tilde{\mathcal{R}}_* := \mathcal{T}^{-1}\mathcal{R}_*\mathcal{T} + \varepsilon^2\Pi_S^\perp\partial_x(\mathcal{R}_2 - \mathcal{T}^{-1}\mathcal{R}_2\mathcal{T}) + \varepsilon^2\Pi_S^\perp\partial_x(\bar{\mathcal{R}}_2 - \mathcal{R}_2), \quad (8.72)$$

and  $\mathcal{R}_*$  is defined in (8.49). We have also used that  $\mathcal{T}^{-1}$  commutes with  $\partial_x$  and with  $\Pi_S^\perp$ .

**Remark 8.12.** The space average  $\frac{1}{2\pi}\int_{\mathbb{T}}\tilde{d}_1(\vartheta, z)dz = \frac{1}{2\pi}\int_{\mathbb{T}}d_1(\vartheta, z)dz = m_1$  for all  $\vartheta \in \mathbb{T}^\nu$ .  $\square$

**Lemma 8.13.** *There is  $\sigma := \sigma(\nu, \tau) > 0$  (possibly larger than in Lemma 8.7) such that*

$$|m_1|^{\text{Lip}(\gamma)} \leq C\varepsilon^4, \quad |\partial_i m_1[\hat{v}]| \leq C\varepsilon^{b+2}\|\hat{v}\|_{s_0+\sigma} \quad (8.73)$$

$$\|p\|_s^{\text{Lip}(\gamma)} \leq_s \varepsilon^4\gamma^{-1} + \varepsilon^{b+2}\gamma^{-1}\|\mathfrak{J}_\delta\|_{s+\sigma}^{\text{Lip}(\gamma)}, \quad \|\partial_i p[\hat{v}]\|_s \leq_s \varepsilon^{b+2}\gamma^{-1}(\|\hat{v}\|_{s+\sigma} + \|\mathfrak{J}_\delta\|_{s+\sigma}\|\hat{v}\|_{s_0+\sigma}), \quad (8.74)$$

$$\|\tilde{d}_k\|_s^{\text{Lip}(\gamma)} \leq_s \varepsilon^5\gamma^{-1} + \varepsilon\|\mathfrak{J}_\delta\|_{s+\sigma}^{\text{Lip}(\gamma)}, \quad \|\partial_i \tilde{d}_k[\hat{v}]\|_s \leq_s \varepsilon(\|\hat{v}\|_{s+\sigma} + \|\mathfrak{J}_\delta\|_{s+\sigma}\|\hat{v}\|_{s_0+\sigma}) \quad (8.75)$$

for  $k = 0, 1$ . Moreover the matrix  $s$ -decay norm (see (2.16))

$$|\tilde{\mathcal{R}}_*|_s^{\text{Lip}(\gamma)} \leq_s \varepsilon^3 + \varepsilon^2\|\mathfrak{J}_\delta\|_{s+\sigma}^{\text{Lip}(\gamma)}, \quad |\partial_i \tilde{\mathcal{R}}_*[\hat{v}]|_s \leq_s \varepsilon^2\|\hat{v}\|_{s+\sigma} + \varepsilon^{2b-1}\|\mathfrak{J}_\delta\|_{s+\sigma}\|\hat{v}\|_{s_0+\sigma}. \quad (8.76)$$

The transformations  $\mathcal{T}, \mathcal{T}^{-1}$  satisfy (8.39), (8.40).

*Proof.* The estimates (8.73), (8.74) follow by (8.67), (8.59), (8.68), and the bounds for  $c_{1,\geq 4}, c_{0,\geq 4}$  in Lemma 8.7. The estimates (8.75) follow similarly by (8.63), (8.68), (8.74). The estimates (8.76) follow because  $\mathcal{T}^{-1}\mathcal{R}_*\mathcal{T}$  satisfies the bounds (7.41) like  $\mathcal{R}_*$  does (use Lemma 7.3 and (8.74)) and  $|\varepsilon^2\Pi_S^\perp\partial_x(\bar{\mathcal{R}}_2 - \mathcal{R}_2)|_s^{\text{Lip}(\gamma)} \leq_s \varepsilon^2\|\mathfrak{J}_\delta\|_{s+\sigma}^{\text{Lip}(\gamma)}$ .  $\square$

It is sufficient to estimate  $\tilde{\mathcal{R}}_*$  (which has the form (7.7)) only in the  $s$ -decay norm (see (8.76)) because the next transformations will preserve it. Such norms are used in the reducibility scheme of section 8.7.

## 8.4 Linear Birkhoff normal form. Step 1

Now we eliminate the terms of order  $\varepsilon$  and  $\varepsilon^2$  of  $\mathcal{L}_3$ . This step is different from the reducibility steps that we shall perform in section 8.7, because the diophantine constant  $\gamma = o(\varepsilon^2)$  (see (5.4)) and so terms  $O(\varepsilon), O(\varepsilon^2)$  are not perturbative. This reduction is possible thanks to the special form of the terms  $\varepsilon\mathcal{B}_1, \varepsilon^2\mathcal{B}_2$  defined in (8.77): the harmonics of  $\varepsilon\mathcal{B}_1$ , and  $\varepsilon^2T$  in (8.93), which correspond to a possible small divisor are naught, see Corollary 8.17, and Lemma 8.21. In this section we eliminate the term  $\varepsilon\mathcal{B}_1$ . In section 8.5 we eliminate the terms of order  $\varepsilon^2$ .

Note that, since the previous transformations  $\Phi, B, \mathcal{T}$  are  $O(\varepsilon^4\gamma^{-1})$ -close to the identity, the terms of order  $\varepsilon$  and  $\varepsilon^2$  in  $\mathcal{L}_3$  are the same as in the original linearized operator.

We first collect all the terms of order  $\varepsilon$  and  $\varepsilon^2$  in the operator  $\mathcal{L}_3$  defined in (8.65). By (8.69), (7.29), (8.57) we have, renaming  $\vartheta = \varphi, z = x$ ,

$$\mathcal{L}_3 = \Pi_S^\perp(\omega \cdot \partial_\varphi + m_3\partial_{xxx} + \varepsilon\mathcal{B}_1 + \varepsilon^2\mathcal{B}_2 + \tilde{d}_1\partial_x + \tilde{d}_0)\Pi_S^\perp + \tilde{\mathcal{R}}_*$$

where  $\tilde{d}_1, \tilde{d}_0, \tilde{\mathcal{R}}_*$  are defined in (8.70)-(8.72) and (recall also (2.2))

$$\mathcal{B}_1 h := -6\partial_x(\bar{v}h), \quad \mathcal{B}_2 h := -6\partial_x\{\bar{v}\Pi_S[(\partial_x^{-1}\bar{v})\partial_x^{-1}h] + h\pi_0[(\partial_x^{-1}\bar{v})^2]\} + 6\pi_0\{(\partial_x^{-1}\bar{v})\Pi_S[\bar{v}h]\}. \quad (8.77)$$

Note that  $\mathcal{B}_1$  and  $\mathcal{B}_2$  are the linear Hamiltonian vector fields of  $H_S^\perp$  generated, respectively, by the Hamiltonian  $z \mapsto 3 \int_{\mathbb{T}} v z^2$  in (3.6), and the fourth order Birkhoff Hamiltonian  $\mathcal{H}_{4,2}$  in (3.7) at  $v = \bar{v}$ .

We transform  $\mathcal{L}_3$  by a symplectic operator  $\Phi_1 : H_{S^\perp}^s(\mathbb{T}^{\nu+1}) \rightarrow H_{S^\perp}^s(\mathbb{T}^{\nu+1})$  of the form

$$\Phi_1 := \exp(\varepsilon A_1) = I_{H_S^\perp} + \varepsilon A_1 + \varepsilon^2 \frac{A_1^2}{2} + \varepsilon^3 \hat{A}_1, \quad \hat{A}_1 := \sum_{k \geq 3} \frac{\varepsilon^{k-3}}{k!} A_1^k, \quad (8.78)$$

where  $A_1(\varphi)h = \sum_{j, j' \in S^c} (A_1)_{j'}^{j'}(\varphi) h_{j'} e^{ijx}$  is a Hamiltonian vector field. The map  $\Phi_1$  is symplectic, because it is the time-1 flow of a Hamiltonian vector field. Therefore

$$\begin{aligned} & \mathcal{L}_3 \Phi_1 - \Phi_1 \Pi_S^\perp (\mathcal{D}_\omega + m_3 \partial_{xxx}) \Pi_S^\perp \\ &= \Pi_S^\perp (\varepsilon \{\mathcal{D}_\omega A_1 + m_3 [\partial_{xxx}, A_1] + \mathcal{B}_1\} + \varepsilon^2 \{\mathcal{B}_1 A_1 + \mathcal{B}_2 + \frac{1}{2} m_3 [\partial_{xxx}, A_1^2] + \frac{1}{2} (\mathcal{D}_\omega A_1^2)\} + \tilde{d}_1 \partial_x + R_3) \Pi_S^\perp \end{aligned} \quad (8.79)$$

where

$$R_3 := \tilde{d}_1 \partial_x (\Phi_1 - I) + \tilde{d}_0 \Phi_1 + \tilde{\mathcal{R}}_* \Phi_1 + \varepsilon^2 \mathcal{B}_2 (\Phi_1 - I) + \varepsilon^3 \{\mathcal{D}_\omega \hat{A}_1 + m_3 [\partial_{xxx}, \hat{A}_1] + \frac{1}{2} \mathcal{B}_1 A_1^2 + \varepsilon \mathcal{B}_1 \hat{A}_1\}. \quad (8.80)$$

**Remark 8.14.**  $R_3$  has no longer the form (7.7). However  $R_3 = O(\partial_x^0)$  because  $A_1 = O(\partial_x^{-1})$  (see Lemma 8.19), and therefore  $\Phi_1 - I_{H_S^\perp} = O(\partial_x^{-1})$ . Moreover the matrix decay norm of  $R_3$  is  $o(\varepsilon^2)$ .  $\square$

In order to eliminate the order  $\varepsilon$  from (8.79), we choose

$$(A_1)_{j'}^{j'}(l) := \begin{cases} -\frac{(\mathcal{B}_1)_{j'}^{j'}(l)}{i(\bar{\omega} \cdot l + m_3(j'^3 - j^3))} & \text{if } \bar{\omega} \cdot l + j'^3 - j^3 \neq 0, \\ 0 & \text{otherwise,} \end{cases} \quad j, j' \in S^c, l \in \mathbb{Z}^\nu. \quad (8.81)$$

This definition is well posed. Indeed, by (8.77) and (8.57),

$$(\mathcal{B}_1)_{j'}^{j'}(l) := \begin{cases} -6ij \sqrt{\xi_{j-j'}} & \text{if } j - j' \in S, \quad l = \ell(j - j') \\ 0 & \text{otherwise.} \end{cases} \quad (8.82)$$

In particular  $(\mathcal{B}_1)_{j'}^{j'}(l) = 0$  unless  $|l| \leq 1$ . Thus, for  $\bar{\omega} \cdot l + j'^3 - j^3 \neq 0$ , the denominators in (8.81) satisfy

$$\begin{aligned} |\bar{\omega} \cdot l + m_3(j'^3 - j^3)| &= |m_3(\bar{\omega} \cdot l + j'^3 - j^3) + (\omega - m_3 \bar{\omega}) \cdot l| \\ &\geq |m_3| |\bar{\omega} \cdot l + j'^3 - j^3| - |\omega - m_3 \bar{\omega}| |l| \geq 1/2, \quad \forall |l| \leq 1, \end{aligned} \quad (8.83)$$

for  $\varepsilon$  small, because the non zero integer  $|\bar{\omega} \cdot l + j'^3 - j^3| \geq 1$ , (8.50), and  $\omega = \bar{\omega} + O(\varepsilon^2)$ .

$A_1$  defined in (8.81) is a Hamiltonian vector field as  $\mathcal{B}_1$ .

**Remark 8.15.** This is a general fact: the denominators  $\delta_{l,j,k} := i(\bar{\omega} \cdot l + m_3(k^3 - j^3))$  satisfy  $\overline{\delta_{l,j,k}} = \delta_{-l,k,j}$  and an operator  $G(\varphi)$  is self-adjoint if and only if its matrix elements satisfy  $G_j^k(l) = G_k^j(-l)$ , see [2]-Remark 4.5. In a more intrinsic way, we could solve the homological equation of this Birkhoff step directly for the Hamiltonian function whose flow generates  $\Phi_1$ .  $\square$

**Lemma 8.16.** *If  $j, j' \in S^c$ ,  $j - j' \in S$ ,  $l = \ell(j - j')$ , then  $\bar{\omega} \cdot l + j'^3 - j^3 = 3jj'(j' - j) \neq 0$ .*

*Proof.* We have  $\bar{\omega} \cdot l = \bar{\omega} \cdot \ell(j - j') = (j - j')^3$  because  $j - j' \in S$  (see (2.10) and (8.58)). Note that  $j, j' \neq 0$  because  $j, j' \in S^c$ , and  $j - j' \neq 0$  because  $j - j' \in S$ .  $\square$

**Corollary 8.17.** *Let  $j, j' \in S^c$ . If  $\bar{\omega} \cdot l + j'^3 - j^3 = 0$  then  $(\mathcal{B}_1)_{j'}^{j'}(l) = 0$ .*

*Proof.* If  $(\mathcal{B}_1)_{j'}^{j'}(l) \neq 0$  then  $j - j' \in S, l = \ell(j - j')$  by (8.82). Hence  $\bar{\omega} \cdot l + j'^3 - j^3 \neq 0$  by Lemma 8.16.  $\square$

By (8.81) and the previous corollary, the term of order  $\varepsilon$  in (8.79) is

$$\Pi_S^\perp (\mathcal{D}_\omega A_1 + m_3 [\partial_{xxx}, A_1] + \mathcal{B}_1) \Pi_S^\perp = 0. \quad (8.84)$$

We now estimate the transformation  $A_1$ .

**Lemma 8.18.** (i) For all  $l \in \mathbb{Z}^\nu, j, j' \in S^c$ ,

$$|(A_1)_{j'}^{j'}(l)| \leq C(|j| + |j'|)^{-1}, \quad |(A_1)_{j'}^{j'}(l)|^{\text{lip}} \leq \varepsilon^{-2}(|j| + |j'|)^{-1}. \quad (8.85)$$

(ii)  $(A_1)_{j'}^{j'}(l) = 0$  for all  $l \in \mathbb{Z}^\nu, j, j' \in S^c$  such that  $|j - j'| > C_S$ , where  $C_S := \max\{|j| : j \in S\}$ .

*Proof.* (i) We already noted that  $(A_1)_{j'}^{j'}(l) = 0, \forall |l| > 1$ . Since  $|\omega| \leq |\bar{\omega}| + 1$ , one has, for  $|l| \leq 1, j \neq j'$ ,

$$|\omega \cdot l + m_3(j'^3 - j^3)| \geq |m_3||j'^3 - j^3| - |\omega \cdot l| \geq \frac{1}{4}(j'^2 + j^2) - |\omega| \geq \frac{1}{8}(j'^2 + j^2), \quad \forall (j'^2 + j^2) \geq C,$$

for some constant  $C > 0$ . Moreover, recalling that also (8.83) holds, we deduce that for  $j \neq j'$ ,

$$(A_1)_{j'}^{j'}(l) \neq 0 \quad \Rightarrow \quad |\omega \cdot l + m_3(j'^3 - j^3)| \geq c(|j| + |j'|)^2. \quad (8.86)$$

On the other hand, if  $j = j', j \in S^c$ , the matrix  $(A_1)_j^j(l) = 0, \forall l \in \mathbb{Z}^\nu$ , because  $(\mathcal{B}_1)_j^j(l) = 0$  by (8.82) (recall that  $0 \notin S$ ). Hence (8.86) holds for all  $j, j'$ . By (8.81), (8.86), (8.82) we deduce the first bound in (8.85). The Lipschitz bound follows similarly (use also  $|j - j'| \leq C_S$ ). (ii) follows by (8.81)-(8.82).  $\square$

The previous lemma means that  $A = O(|\partial_x|^{-1})$ . More precisely we deduce that

**Lemma 8.19.**  $|A_1 \partial_x|_s^{\text{Lip}(\gamma)} + |\partial_x A_1|_s^{\text{Lip}(\gamma)} \leq C(s)$ .

*Proof.* Recalling the definition of the (space-time) matrix norm in (2.23), since  $(A_1)_{j_1}^{j_2}(l) = 0$  outside the set of indices  $|l| \leq 1, |j_1 - j_2| \leq C_S$ , we have

$$|\partial_x A_1|_s^2 = \sum_{|l| \leq 1, |j| \leq C_S} \left( \sup_{|j_1 - j_2 = j} |j_1| |(A_1)_{j_1}^{j_2}(l)| \right)^2 \langle l, j \rangle^{2s} \leq C(s)$$

by Lemma 8.18. The estimates for  $|A_1 \partial_x|_s$  and the Lipschitz bounds follow similarly.  $\square$

It follows that the symplectic map  $\Phi_1$  in (8.78) is invertible for  $\varepsilon$  small, with inverse

$$\Phi_1^{-1} = \exp(-\varepsilon A_1) = I_{H_S} + \varepsilon \check{A}_1, \quad \check{A}_1 := \sum_{n \geq 1} \frac{\varepsilon^{n-1}}{n!} (-A_1)^n, \quad |\check{A}_1 \partial_x|_s^{\text{Lip}(\gamma)} + |\partial_x \check{A}_1|_s^{\text{Lip}(\gamma)} \leq C(s). \quad (8.87)$$

Since  $A_1$  solves the homological equation (8.84), the  $\varepsilon$ -term in (8.79) is zero, and, with a straightforward calculation, the  $\varepsilon^2$ -term simplifies to  $\mathcal{B}_2 + \frac{1}{2}[\mathcal{B}_1, A_1]$ . We obtain the Hamiltonian operator

$$\mathcal{L}_4 := \Phi_1^{-1} \mathcal{L}_3 \Phi_1 = \Pi_S^\perp (\mathcal{D}_\omega + m_3 \partial_{xxx} + \tilde{d}_1 \partial_x + \varepsilon^2 \{\mathcal{B}_2 + \frac{1}{2}[\mathcal{B}_1, A_1]\} + \tilde{R}_4) \Pi_S^\perp \quad (8.88)$$

$$\tilde{R}_4 := (\Phi_1^{-1} - I) \Pi_S^\perp [\varepsilon^2 (\mathcal{B}_2 + \frac{1}{2}[\mathcal{B}_1, A_1]) + \tilde{d}_1 \partial_x] + \Phi_1^{-1} \Pi_S^\perp R_3. \quad (8.89)$$

We split  $A_1$  defined in (8.81), (8.82) into  $A_1 = \bar{A}_1 + \tilde{A}_1$  where, for all  $j, j' \in S^c, l \in \mathbb{Z}^\nu$ ,

$$(\bar{A}_1)_j^{j'}(l) := \frac{6j \sqrt{\xi_{j-j'}}}{\bar{\omega} \cdot l + j'^3 - j^3} \quad \text{if } \bar{\omega} \cdot l + j'^3 - j^3 \neq 0, \quad j - j' \in S, \quad l = \ell(j - j'), \quad (8.90)$$

and  $(\bar{A}_1)_j^{j'}(l) := 0$  otherwise. By Lemma 8.16, for all  $j, j' \in S^c$ ,  $l \in \mathbb{Z}^\nu$ ,  $(\bar{A}_1)_j^{j'}(l) = \frac{2\sqrt{\xi_{j-j'}}}{j'(j'-j)}$  if  $j - j' \in S$ ,  $l = \ell(j - j')$ , and  $(\bar{A}_1)_j^{j'}(l) = 0$  otherwise, namely (recall the definition of  $\bar{v}$  in (8.57))

$$\bar{A}_1 h = 2\Pi_S^\perp [(\partial_x^{-1}\bar{v})(\partial_x^{-1}h)], \quad \forall h \in H_{S^\perp}^s(\mathbb{T}^{\nu+1}). \quad (8.91)$$

The difference is

$$(\tilde{A}_1)_j^{j'}(l) = (A_1 - \bar{A}_1)_j^{j'}(l) = -\frac{6j\sqrt{\xi_{j-j'}}\{(\omega - \bar{\omega}) \cdot l + (m_3 - 1)(j'^3 - j^3)\}}{(\omega \cdot l + m_3(j'^3 - j^3))(\bar{\omega} \cdot l + j'^3 - j^3)} \quad (8.92)$$

for  $j, j' \in S^c$ ,  $j - j' \in S$ ,  $l = \ell(j - j')$ , and  $(\tilde{A}_1)_j^{j'}(l) = 0$  otherwise. Then, by (8.88),

$$\mathcal{L}_4 = \Pi_S^\perp (\mathcal{D}_\omega + m_3 \partial_{xxx} + \tilde{d}_1 \partial_x + \varepsilon^2 T + R_4) \Pi_S^\perp, \quad (8.93)$$

where

$$T := \mathcal{B}_2 + \frac{1}{2}[\mathcal{B}_1, \bar{A}_1], \quad R_4 := \frac{\varepsilon^2}{2}[\mathcal{B}_1, \tilde{A}_1] + \tilde{R}_4. \quad (8.94)$$

The operator  $T$  is Hamiltonian as  $\mathcal{B}_2$ ,  $\mathcal{B}_1$ ,  $\bar{A}_1$  (the commutator of two Hamiltonian vector fields is Hamiltonian).

**Lemma 8.20.** *There is  $\sigma = \sigma(\nu, \tau) > 0$  (possibly larger than in Lemma 8.13) such that*

$$|R_4|_s^{\text{Lip}(\gamma)} \leq_s \varepsilon^5 \gamma^{-1} + \varepsilon \|\mathcal{J}_\delta\|_{s+\sigma}^{\text{Lip}(\gamma)}, \quad |\partial_i R_4[\tilde{v}]|_s \leq_s \varepsilon (\|\tilde{v}\|_{s+\sigma} + \|\mathcal{J}_\delta\|_{s+\sigma} \|\tilde{v}\|_{s_0+\sigma}). \quad (8.95)$$

*Proof.* We first estimate  $[\mathcal{B}_1, \tilde{A}_1] = (\mathcal{B}_1 \partial_x^{-1})(\partial_x \tilde{A}_1) - (\tilde{A}_1 \partial_x)(\partial_x^{-1} \mathcal{B}_1)$ . By (8.92),  $|\omega - \bar{\omega}| \leq C\varepsilon^2$  (as  $\omega \in \Omega_\varepsilon$  in (5.2)) and (8.50), arguing as in Lemmata 8.18, 8.19, we deduce that  $|\tilde{A}_1 \partial_x|_s^{\text{Lip}(\gamma)} + |\partial_x \tilde{A}_1|_s^{\text{Lip}(\gamma)} \leq_s \varepsilon^2$ . By (8.77) the norm  $|\mathcal{B}_1 \partial_x^{-1}|_s^{\text{Lip}(\gamma)} + |\partial_x^{-1} \mathcal{B}_1|_s^{\text{Lip}(\gamma)} \leq C(s)$ . Hence  $\varepsilon^2 |[\mathcal{B}_1, \tilde{A}_1]|_s^{\text{Lip}(\gamma)} \leq_s \varepsilon^4$ . Finally (8.94), (8.89), (8.87), (8.80), (8.75), (8.76), and the interpolation estimate (2.20) imply (8.95).  $\square$

## 8.5 Linear Birkhoff normal form. Step 2

The goal of this section is to remove the term  $\varepsilon^2 T$  from the operator  $\mathcal{L}_4$  defined in (8.93). We conjugate the Hamiltonian operator  $\mathcal{L}_4$  via a symplectic map

$$\Phi_2 := \exp(\varepsilon^2 A_2) = I_{H_S^\perp} + \varepsilon^2 A_2 + \varepsilon^4 \hat{A}_2, \quad \hat{A}_2 := \sum_{k \geq 2} \frac{\varepsilon^{2(k-2)}}{k!} A_2^k \quad (8.96)$$

where  $A_2(\varphi) = \sum_{j, j' \in S^c} (A_2)_j^{j'}(\varphi) h_{j'} e^{ijx}$  is a Hamiltonian vector field. We compute

$$\mathcal{L}_4 \Phi_2 - \Phi_2 \Pi_S^\perp (\mathcal{D}_\omega + m_3 \partial_{xxx}) \Pi_S^\perp = \Pi_S^\perp (\varepsilon^2 \{\mathcal{D}_\omega A_2 + m_3 [\partial_{xxx}, A_2] + T\} + \tilde{d}_1 \partial_x + \tilde{R}_5) \Pi_S^\perp, \quad (8.97)$$

$$\tilde{R}_5 := \Pi_S^\perp \{\varepsilon^4 ((\mathcal{D}_\omega \hat{A}_2) + m_3 [\partial_{xxx}, \hat{A}_2]) + (\tilde{d}_1 \partial_x + \varepsilon^2 T)(\Phi_2 - I) + R_4 \Phi_2\} \Pi_S^\perp. \quad (8.98)$$

We define

$$(A_2)_j^{j'}(l) := -\frac{T_j^{j'}(l)}{i(\omega \cdot l + m_3(j'^3 - j^3))} \quad \text{if } \bar{\omega} \cdot l + j'^3 - j^3 \neq 0; \quad (A_2)_j^{j'}(l) := 0 \quad \text{otherwise.} \quad (8.99)$$

This definition is well posed. Indeed, by (8.94), (8.82), (8.90), (8.77), the matrix entries  $T_j^{j'}(l) = 0$  for all  $|j - j'| > 2C_S$ ,  $l \in \mathbb{Z}^\nu$ , where  $C_S := \max\{|j|, j \in S\}$ . Also  $T_j^{j'}(l) = 0$  for all  $j, j' \in S^c$ ,  $|l| > 2$  (see also (8.100), (8.103), (8.104) below). Thus, arguing as in (8.83), if  $\bar{\omega} \cdot l + j'^3 - j^3 \neq 0$ , then  $|\omega \cdot l + m_3(j'^3 - j^3)| \geq 1/2$ . The operator  $A_2$  is a Hamiltonian vector field because  $T$  is Hamiltonian and by Remark 8.15.

Now we prove that the Birkhoff map  $\Phi_2$  removes completely the term  $\varepsilon^2 T$ .

**Lemma 8.21.** *Let  $j, j' \in S^c$ . If  $\bar{\omega} \cdot l + j'^3 - j^3 = 0$ , then  $T_j^{j'}(l) = 0$ .*

*Proof.* By (8.77), (8.91) we get  $\mathcal{B}_1 \bar{A}_1 h = -12\partial_x \{ \bar{v} \Pi_S^\perp [(\partial_x^{-1} \bar{v})(\partial_x^{-1} h)] \}$ ,  $\bar{A}_1 \mathcal{B}_1 h = -12\Pi_S^\perp [(\partial_x^{-1} \bar{v}) \Pi_S^\perp (\bar{v} h)]$  for all  $h \in H_{\bar{S}^\perp}^s$ , whence, recalling (8.57), for all  $j, j' \in S^c$ ,  $l \in \mathbb{Z}^\nu$ ,

$$([\mathcal{B}_1, \bar{A}_1]_j^{j'}(l) = 12i \sum_{\substack{j_1, j_2 \in S, j_1 + j_2 = j - j' \\ j' + j_2 \in S^c, \ell(j_1) + \ell(j_2) = l}} \frac{j j_1 - j' j_2}{j' j_1 j_2} \sqrt{\xi_{j_1} \xi_{j_2}}, \quad (8.100)$$

If  $([\mathcal{B}_1, \bar{A}_1]_j^{j'}(l) \neq 0$  there are  $j_1, j_2 \in S$  such that  $j_1 + j_2 = j - j'$ ,  $j' + j_2 \in S^c$ ,  $\ell(j_1) + \ell(j_2) = l$ . Then

$$\bar{\omega} \cdot l + j'^3 - j^3 = \bar{\omega} \cdot \ell(j_1) + \bar{\omega} \cdot \ell(j_2) + j'^3 - j^3 \stackrel{(8.58)}{=} j_1^3 + j_2^3 + j'^3 - j^3. \quad (8.101)$$

Thus, if  $\bar{\omega} \cdot l + j'^3 - j^3 = 0$ , Lemma 3.3 implies  $(j_1 + j_2)(j_1 + j')(j_2 + j') = 0$ . Now  $j_1 + j'$ ,  $j_2 + j' \neq 0$  because  $j_1, j_2 \in S$ ,  $j' \in S^c$  and  $S$  is symmetric. Hence  $j_1 + j_2 = 0$ , which implies  $j = j'$  and  $l = 0$  (the map  $\ell$  in (8.58) is odd). In conclusion, if  $\bar{\omega} \cdot l + j'^3 - j^3 = 0$ , the only nonzero matrix entry  $([\mathcal{B}_1, \bar{A}_1]_j^{j'}(l)$  is

$$([\mathcal{B}_1, \bar{A}_1]_j^j(0) \stackrel{(8.100)}{=} 24i \sum_{j_2 \in S, j_2 + j \in S^c} \xi_{j_2} j_2^{-1}. \quad (8.102)$$

Now we consider  $\mathcal{B}_2$  in (8.77). Split  $\mathcal{B}_2 = B_1 + B_2 + B_3$ , where  $B_1 h := -6\partial_x \{ \bar{v} \Pi_S [(\partial_x^{-1} \bar{v}) \partial_x^{-1} h] \}$ ,  $B_2 h := -6\partial_x \{ h \pi_0 [(\partial_x^{-1} \bar{v})^2] \}$ ,  $B_3 h := 6\pi_0 \{ \Pi_S (\bar{v} h) \partial_x^{-1} \bar{v} \}$ . Their Fourier matrix representation is

$$(B_1)_j^{j'}(l) = 6ij \sum_{\substack{j_1, j_2 \in S, j_1 + j' \in S \\ j_1 + j_2 = j - j', \ell(j_1) + \ell(j_2) = l}} \frac{\sqrt{\xi_{j_1} \xi_{j_2}}}{j_1 j'}, \quad (B_2)_j^{j'}(l) = 6ij \sum_{\substack{j_1, j_2 \in S, j_1 + j_2 \neq 0 \\ j_1 + j_2 = j - j', \ell(j_1) + \ell(j_2) = l}} \frac{\sqrt{\xi_{j_1} \xi_{j_2}}}{j_1 j_2}, \quad (8.103)$$

$$(B_3)_j^{j'}(l) = 6 \sum_{\substack{j_1, j_2 \in S, j_1 + j' \in S \\ j_1 + j_2 = j - j', \ell(j_1) + \ell(j_2) = l}} \frac{\sqrt{\xi_{j_1} \xi_{j_2}}}{ij_2}, \quad j, j' \in S^c, l \in \mathbb{Z}^\nu. \quad (8.104)$$

We study the terms  $B_1, B_2, B_3$  separately. If  $(B_1)_j^{j'}(l) \neq 0$ , there are  $j_1, j_2 \in S$  such that  $j_1 + j_2 = j - j'$ ,  $j_1 + j' \in S$ ,  $l = \ell(j_1) + \ell(j_2)$  and (8.101) holds. Thus, if  $\bar{\omega} \cdot l + j'^3 - j^3 = 0$ , Lemma 3.3 implies  $(j_1 + j_2)(j_1 + j')(j_2 + j') = 0$ , and, since  $j' \in S^c$  and  $S$  is symmetric, the only possibility is  $j_1 + j_2 = 0$ . Hence  $j = j'$ ,  $l = 0$ . In conclusion, if  $\bar{\omega} \cdot l + j'^3 - j^3 = 0$ , the only nonzero matrix element  $(B_1)_j^{j'}(l)$  is

$$(B_1)_j^j(0) = 6i \sum_{j_1 \in S, j_1 + j \in S} \xi_{j_1} j_1^{-1}. \quad (8.105)$$

By the same arguments, if  $(B_2)_j^{j'}(l) \neq 0$  and  $\bar{\omega} \cdot l + j'^3 - j^3 = 0$  we find  $(j_1 + j_2)(j_1 + j')(j_2 + j') = 0$ , which is impossible because also  $j_1 + j_2 \neq 0$ . Finally, arguing as for  $B_1$ , if  $\bar{\omega} \cdot l + j'^3 - j^3 = 0$ , then the only nonzero matrix element  $(B_3)_j^{j'}(l)$  is

$$(B_3)_j^j(0) = 6i \sum_{j_1 \in S, j_1 + j \in S} \xi_{j_1} j_1^{-1}. \quad (8.106)$$

From (8.102), (8.105), (8.106) we deduce that, if  $\bar{\omega} \cdot l + j'^3 - j^3 = 0$ , then the only non zero elements  $(\frac{1}{2}[\mathcal{B}_1, \bar{A}_1] + B_1 + B_3)_j^{j'}(l)$  must be for  $(l, j, j') = (0, j, j)$ . In this case, we get

$$\frac{1}{2}([\mathcal{B}_1, \bar{A}_1]_j^j(0) + (B_1)_j^j(0) + (B_3)_j^j(0) = 12i \sum_{\substack{j_1 \in S \\ j_1 + j \in S^c}} \frac{\xi_{j_1}}{j_1} + 12i \sum_{\substack{j_1 \in S \\ j_1 + j \in S}} \frac{\xi_{j_1}}{j_1} = 12i \sum_{j_1 \in S} \frac{\xi_{j_1}}{j_1} = 0 \quad (8.107)$$

because the case  $j_1 + j = 0$  is impossible ( $j_1 \in S$ ,  $j' \in S^c$  and  $S$  is symmetric), and the function  $S \ni j_1 \rightarrow \xi_{j_1}/j_1 \in \mathbb{R}$  is odd. The lemma follows by (8.94), (8.107).  $\square$

The choice of  $A_2$  in (8.99) and Lemma 8.21 imply that

$$\Pi_S^\perp (\mathcal{D}_\omega A_2 + m_3 [\partial_{xxx}, A_2] + T) \Pi_S^\perp = 0. \quad (8.108)$$

**Lemma 8.22.**  $|\partial_x A_2|_s^{\text{Lip}(\gamma)} + |A_2 \partial_x|_s^{\text{Lip}(\gamma)} \leq C(s)$ .

*Proof.* First we prove that the diagonal elements  $T_j^j(l) = 0$  for all  $l \in \mathbb{Z}^\nu$ . For  $l = 0$ , we have already proved that  $T_j^j(0) = 0$  (apply Lemma 8.21 with  $j = j'$ ,  $l = 0$ ). Moreover, in each term  $[\mathcal{B}_1, \bar{A}_1]$ ,  $B_1$ ,  $B_2$ ,  $B_3$  (see (8.100), (8.103), (8.104)) the sum is over  $j_1 + j_2 = j - j'$ ,  $l = \ell(j_1) + \ell(j_2)$ . If  $j = j'$ , then  $j_1 + j_2 = 0$ , and  $l = 0$ . Thus  $T_j^j(l) = T_j^j(0) = 0$ . For the off-diagonal terms  $j \neq j'$  we argue as in Lemmata 8.18, 8.19, using that all the denominators  $|\omega \cdot l + m_3(j'^3 - j^3)| \geq c(|j| + |j'|)^2$ .  $\square$

For  $\varepsilon$  small, the map  $\Phi_2$  in (8.96) is invertible and  $\Phi_2 = \exp(-\varepsilon^2 A_2)$ . Therefore (8.97), (8.108) imply

$$\mathcal{L}_5 := \Phi_2^{-1} \mathcal{L}_4 \Phi_2 = \Pi_S^\perp (\mathcal{D}_\omega + m_3 \partial_{xxx} + \tilde{d}_1 \partial_x + R_5) \Pi_S^\perp, \quad (8.109)$$

$$R_5 := (\Phi_2^{-1} - I) \Pi_S^\perp \tilde{d}_1 \partial_x + \Phi_2^{-1} \Pi_S^\perp \tilde{R}_5. \quad (8.110)$$

Since  $A_2$  is a Hamiltonian vector field, the map  $\Phi_2$  is symplectic and so  $\mathcal{L}_5$  is Hamiltonian.

**Lemma 8.23.**  $R_5$  satisfies the same estimates (8.95) as  $R_4$  (with a possibly larger  $\sigma$ ).

*Proof.* Use (8.110), Lemma 8.22, (8.75), (8.98), (8.95) and the interpolation inequalities (2.18), (2.20).  $\square$

## 8.6 Descent method

The goal of this section is to transform  $\mathcal{L}_5$  in (8.109) so that the coefficient of  $\partial_x$  becomes constant. We conjugate  $\mathcal{L}_5$  via a symplectic map of the form

$$\mathcal{S} := \exp(\Pi_S^\perp (w \partial_x^{-1})) \Pi_S^\perp = \Pi_S^\perp (I + w \partial_x^{-1}) \Pi_S^\perp + \widehat{\mathcal{S}}, \quad \widehat{\mathcal{S}} := \sum_{k \geq 2} \frac{1}{k!} [\Pi_S^\perp (w \partial_x^{-1})]^k \Pi_S^\perp, \quad (8.111)$$

where  $w : \mathbb{T}^{\nu+1} \rightarrow \mathbb{R}$  is a function. Note that  $\Pi_S^\perp (w \partial_x^{-1}) \Pi_S^\perp$  is the Hamiltonian vector field generated by  $-\frac{1}{2} \int_{\mathbb{T}} w (\partial_x^{-1} h)^2 dx$ ,  $h \in H_S^\perp$ . Recalling (2.2), we calculate

$$\mathcal{L}_5 \mathcal{S} - \mathcal{S} \Pi_S^\perp (\mathcal{D}_\omega + m_3 \partial_{xxx} + m_1 \partial_x) \Pi_S^\perp = \Pi_S^\perp (3m_3 w_x + \tilde{d}_1 - m_1) \partial_x \Pi_S^\perp + \tilde{R}_6, \quad (8.112)$$

$$\begin{aligned} \tilde{R}_6 := & \Pi_S^\perp \{ (3m_3 w_{xx} + \tilde{d}_1 \Pi_S^\perp w - m_1 w) \pi_0 + ((\mathcal{D}_\omega w) + m_3 w_{xxx} + \tilde{d}_1 \Pi_S^\perp w_x) \partial_x^{-1} + (\mathcal{D}_\omega \widehat{\mathcal{S}}) \\ & + m_3 [\partial_{xxx}, \widehat{\mathcal{S}}] + \tilde{d}_1 \partial_x \widehat{\mathcal{S}} - m_1 \widehat{\mathcal{S}} \partial_x + R_5 \mathcal{S} \} \Pi_S^\perp \end{aligned}$$

where  $\tilde{R}_6$  collects all the terms of order at most  $\partial_x^0$ . By Remark 8.12, we solve  $3m_3 w_x + \tilde{d}_1 - m_1 = 0$  by choosing  $w := -(3m_3)^{-1} \partial_x^{-1} (\tilde{d}_1 - m_1)$ . For  $\varepsilon$  small, the operator  $\mathcal{S}$  is invertible and, by (8.112),

$$\mathcal{L}_6 := \mathcal{S}^{-1} \mathcal{L}_5 \mathcal{S} = \Pi_S^\perp (\mathcal{D}_\omega + m_3 \partial_{xxx} + m_1 \partial_x) \Pi_S^\perp + R_6, \quad R_6 := \mathcal{S}^{-1} \tilde{R}_6. \quad (8.113)$$

Since  $\mathcal{S}$  is symplectic,  $\mathcal{L}_6$  is Hamiltonian (recall Definition 2.2).

**Lemma 8.24.** *There is  $\sigma = \sigma(\nu, \tau) > 0$  (possibly larger than in Lemma 8.23) such that*

$$|\mathcal{S}^{\pm 1} - I|_s^{\text{Lip}(\gamma)} \leq_s \varepsilon^5 \gamma^{-1} + \varepsilon \|\mathcal{J}_\delta\|_{s+\sigma}^{\text{Lip}(\gamma)}, \quad |\partial_i \mathcal{S}^{\pm 1}[\hat{v}]|_s \leq_s \varepsilon (\|\hat{v}\|_{s+\sigma} + \|\mathcal{J}_\delta\|_{s+\sigma} \|\hat{v}\|_{s_0+\sigma}).$$

*The remainder  $R_6$  satisfies the same estimates (8.95) as  $R_4$ .*

*Proof.* By (8.75), (8.73), (8.50),  $\|w\|_s^{\text{Lip}(\gamma)} \leq_s \varepsilon^5 \gamma^{-1} + \varepsilon \|\mathcal{J}_\delta\|_{s+\sigma}^{\text{Lip}(\gamma)}$ , and the lemma follows by (8.111). Since  $\widehat{\mathcal{S}} = O(\partial_x^{-2})$  the commutator  $[\partial_{xxx}, \widehat{\mathcal{S}}] = O(\partial_x^0)$  and  $\|[\partial_{xxx}, \widehat{\mathcal{S}}]\|_s^{\text{Lip}(\gamma)} \leq_s \|w\|_{s_0+3}^{\text{Lip}(\gamma)} \|w\|_{s+3}^{\text{Lip}(\gamma)}$ .  $\square$



## 8.7 KAM reducibility and inversion of $\mathcal{L}_\omega$

The coefficients  $m_3, m_1$  of the operator  $\mathcal{L}_6$  in (8.113) are constants, and the remainder  $R_6$  is a bounded operator of order  $\partial_x^0$  with small matrix decay norm, see (8.116). Then we can diagonalize  $\mathcal{L}_6$  by applying the iterative KAM reducibility Theorem 4.2 in [2] along the sequence of scales

$$N_n := N_0^{\chi^n}, \quad n = 0, 1, 2, \dots, \quad \chi := 3/2, \quad N_0 > 0. \quad (8.114)$$

In section 9, the initial  $N_0$  will (slightly) increase to infinity as  $\varepsilon \rightarrow 0$ , see (9.5). The required smallness condition (see (4.14) in [2]) is (written in the present notations)

$$N_0^{C_0} |R_6|_{s_0+\beta}^{\text{Lip}(\gamma)} \gamma^{-1} \leq 1 \quad (8.115)$$

where  $\beta := 7\tau + 6$  (see (4.1) in [2]),  $\tau$  is the diophantine exponent in (5.4) and (8.120), and the constant  $C_0 := C_0(\tau, \nu) > 0$  is fixed in Theorem 4.2 in [2]. By Lemma 8.24, the remainder  $R_6$  satisfies the bound (8.95), and using (7.8) we get (recall (5.10))

$$|R_6|_{s_0+\beta}^{\text{Lip}(\gamma)} \leq C\varepsilon^{7-2b}\gamma^{-1} = C\varepsilon^{3-2a}, \quad |R_6|_{s_0+\beta}^{\text{Lip}(\gamma)} \gamma^{-1} \leq C\varepsilon^{1-3a}. \quad (8.116)$$

We use that  $\mu$  in (7.8) is assumed to satisfy  $\mu \geq \sigma + \beta$  where  $\sigma := \sigma(\tau, \nu)$  is given in Lemma 8.24.

**Theorem 8.25. (Reducibility)** *Assume that  $\omega \mapsto i_\delta(\omega)$  is a Lipschitz function defined on some subset  $\Omega_o \subset \Omega_\varepsilon$  (recall (5.2)), satisfying (7.8) with  $\mu \geq \sigma + \beta$  where  $\sigma := \sigma(\tau, \nu)$  is given in Lemma 8.24 and  $\beta := 7\tau + 6$ . Then there exists  $\delta_0 \in (0, 1)$  such that, if*

$$N_0^{C_0} \varepsilon^{7-2b} \gamma^{-2} = N_0^{C_0} \varepsilon^{1-3a} \leq \delta_0, \quad \gamma := \varepsilon^{2+a}, \quad a \in (0, 1/6), \quad (8.117)$$

then:

(i) **(Eigenvalues)**. *For all  $\omega \in \Omega_\varepsilon$  there exists a sequence*

$$\mu_j^\infty(\omega) := \mu_j^\infty(\omega, i_\delta(\omega)) := i(-\tilde{m}_3(\omega)j^3 + \tilde{m}_1(\omega)j) + r_j^\infty(\omega), \quad j \in S^c, \quad (8.118)$$

where  $\tilde{m}_3, \tilde{m}_1$  coincide with the coefficients  $m_3, m_1$  of  $\mathcal{L}_6$  in (8.113) for all  $\omega \in \Omega_o$ , and

$$|\tilde{m}_3 - 1|^{\text{Lip}(\gamma)} + |\tilde{m}_1|^{\text{Lip}(\gamma)} \leq C\varepsilon^4, \quad |r_j^\infty|^{\text{Lip}(\gamma)} \leq C\varepsilon^{3-2a}, \quad \forall j \in S^c, \quad (8.119)$$

for some  $C > 0$ . All the eigenvalues  $\mu_j^\infty$  are purely imaginary. We define, for convenience,  $\mu_0^\infty(\omega) := 0$ .

(ii) **(Conjugacy)**. *For all  $\omega$  in the set*

$$\Omega_\infty^{2\gamma} := \Omega_\infty^{2\gamma}(i_\delta) := \left\{ \omega \in \Omega_o : |\omega \cdot l + \mu_j^\infty(\omega) - \mu_k^\infty(\omega)| \geq \frac{2\gamma|j^3 - k^3|}{\langle l \rangle^\tau}, \forall l \in \mathbb{Z}^\nu, j, k \in S^c \cup \{0\} \right\} \quad (8.120)$$

there is a real, bounded, invertible linear operator  $\Phi_\infty(\omega) : H_{S^\perp}^s(\mathbb{T}^{\nu+1}) \rightarrow H_{S^\perp}^s(\mathbb{T}^{\nu+1})$ , with bounded inverse  $\Phi_\infty^{-1}(\omega)$ , that conjugates  $\mathcal{L}_6$  in (8.113) to constant coefficients, namely

$$\mathcal{L}_\infty(\omega) := \Phi_\infty^{-1}(\omega) \circ \mathcal{L}_6(\omega) \circ \Phi_\infty(\omega) = \omega \cdot \partial_\varphi + \mathcal{D}_\infty(\omega), \quad \mathcal{D}_\infty(\omega) := \text{diag}_{j \in S^c} \{ \mu_j^\infty(\omega) \}. \quad (8.121)$$

The transformations  $\Phi_\infty, \Phi_\infty^{-1}$  are close to the identity in matrix decay norm, with

$$|\Phi_\infty - I|_{s, \Omega_\infty^{2\gamma}}^{\text{Lip}(\gamma)} + |\Phi_\infty^{-1} - I|_{s, \Omega_\infty^{2\gamma}}^{\text{Lip}(\gamma)} \leq_s \varepsilon^5 \gamma^{-2} + \varepsilon \gamma^{-1} \|\mathcal{J}_\delta\|_{s+\sigma}^{\text{Lip}(\gamma)}. \quad (8.122)$$

Moreover  $\Phi_\infty, \Phi_\infty^{-1}$  are symplectic, and  $\mathcal{L}_\infty$  is a Hamiltonian operator.

*Proof.* The proof is the same as the one of Theorem 4.1 in [2], which is based on Theorem 4.2, Corollaries 4.1, 4.2 and Lemmata 4.1, 4.2 of [2]. A difference is that here  $\omega \in \mathbb{R}^\nu$ , while in [2] the parameter  $\lambda \in \mathbb{R}$  is one-dimensional. The proof is the same because Kirszbraum's Theorem on Lipschitz extension of functions also holds in  $\mathbb{R}^\nu$  (see, e.g., Lemma A.2 in [25]). The bound (8.122) follows by Corollary 4.1 of [2] and the estimate of  $R_6$  in Lemma 8.24. We also use the estimates (8.50), (8.73) for  $\partial_i m_3, \partial_i m_1$  which correspond to (3.64) in [2]. Another difference is that here the sites  $j \in S^c \subset \mathbb{Z} \setminus \{0\}$  unlike in [2] where  $j \in \mathbb{Z}$ . We have defined  $\mu_0^\infty := 0$  so that also the first Melnikov conditions (8.123) are included in the definition of  $\Omega_\infty^{2\gamma}$ .  $\square$

**Remark 8.26.** Theorem 4.2 in [2] also provides the Lipschitz dependence of the (approximate) eigenvalues  $\mu_j^n$  with respect to the unknown  $i_0(\varphi)$ , which is used for the measure estimate Lemma 9.3.  $\square$

All the parameters  $\omega \in \Omega_\infty^{2\gamma}$  satisfy (specialize (8.120) for  $k = 0$ )

$$|i\omega \cdot l + \mu_j^\infty(\omega)| \geq 2\gamma|j|^3 \langle l \rangle^{-\tau}, \quad \forall l \in \mathbb{Z}^\nu, j \in S^c, \quad (8.123)$$

and the diagonal operator  $\mathcal{L}_\infty$  is invertible.

In the following theorem we finally verify the inversion assumption (6.32) for  $\mathcal{L}_\omega$ .

**Theorem 8.27. (Inversion of  $\mathcal{L}_\omega$ )** *Assume the hypotheses of Theorem 8.25 and (8.117). Then there exists  $\sigma_1 := \sigma_1(\tau, \nu) > 0$  such that,  $\forall \omega \in \Omega_\infty^{2\gamma}(i_\delta)$  (see (8.120)), for any function  $g \in H_{S^\perp}^{s+\sigma_1}(\mathbb{T}^{\nu+1})$  the equation  $\mathcal{L}_\omega h = g$  has a solution  $h = \mathcal{L}_\omega^{-1}g \in H_{S^\perp}^s(\mathbb{T}^{\nu+1})$ , satisfying*

$$\begin{aligned} \|\mathcal{L}_\omega^{-1}g\|_s^{\text{Lip}(\gamma)} &\leq \gamma^{-1}(\|g\|_{s+\sigma_1}^{\text{Lip}(\gamma)} + \varepsilon\gamma^{-1}\|\mathcal{J}_\delta\|_{s+\sigma_1}^{\text{Lip}(\gamma)}\|g\|_{s_0}^{\text{Lip}(\gamma)}) \\ &\leq \gamma^{-1}(\|g\|_{s+\sigma_1}^{\text{Lip}(\gamma)} + \varepsilon\gamma^{-1}\{\|\mathcal{J}_0\|_{s+\sigma_1+\sigma}^{\text{Lip}(\gamma)} + \gamma^{-1}\|\mathcal{J}_0\|_{s_0+\sigma}^{\text{Lip}(\gamma)}\|Z\|_{s+\sigma_1+\sigma}^{\text{Lip}(\gamma)}\}\|g\|_{s_0}^{\text{Lip}(\gamma)}). \end{aligned} \quad (8.124)$$

*Proof.* Collecting Theorem 8.25 with the results of sections 8.1-8.6, we have obtained the (semi)-conjugation of the operator  $\mathcal{L}_\omega$  (defined in (7.34)) to  $\mathcal{L}_\infty$  (defined in (8.121)), namely

$$\mathcal{L}_\omega = \mathcal{M}_1 \mathcal{L}_\infty \mathcal{M}_2^{-1}, \quad \mathcal{M}_1 := \Phi B \rho \mathcal{T} \Phi_1 \Phi_2 \mathcal{S} \Phi_\infty, \quad \mathcal{M}_2 := \Phi B \mathcal{T} \Phi_1 \Phi_2 \mathcal{S} \Phi_\infty, \quad (8.125)$$

where  $\rho$  means the multiplication operator by the function  $\rho$  defined in (8.41). By (8.123) and Lemma 4.2 of [2] we deduce that  $\|\mathcal{L}_\infty^{-1}g\|_s^{\text{Lip}(\gamma)} \leq \gamma^{-1}\|g\|_{s+2\tau+1}^{\text{Lip}(\gamma)}$ . In order to estimate  $\mathcal{M}_2, \mathcal{M}_1^{-1}$ , we recall that the composition of tame maps is tame, see Lemma 6.5 in [2]. Now,  $\Phi, \Phi^{-1}$  are estimated in Lemma 8.5,  $B, B^{-1}$  and  $\rho$  in Lemma 8.7,  $\mathcal{T}, \mathcal{T}^{-1}$  in Lemma 8.13. The decay norms  $|\Phi_1|_s^{\text{Lip}(\gamma)}, |\Phi_1^{-1}|_s^{\text{Lip}(\gamma)}, |\Phi_2|_s^{\text{Lip}(\gamma)}, |\Phi_2^{-1}|_s^{\text{Lip}(\gamma)} \leq C(s)$  by Lemmata 8.19, 8.22. The decay norm of  $\mathcal{S}, \mathcal{S}^{-1}$  is estimated in Lemma 8.24, and  $\Phi_\infty, \Phi_\infty^{-1}$  in (8.122). The decay norm controls the Sobolev norm by (2.21). Thus, by (8.125),

$$\|\mathcal{M}_2 h\|_s^{\text{Lip}(\gamma)} + \|\mathcal{M}_1^{-1} h\|_s^{\text{Lip}(\gamma)} \leq \|h\|_{s+3}^{\text{Lip}(\gamma)} + \varepsilon\gamma^{-1}\|\mathcal{J}_\delta\|_{s+\sigma+3}^{\text{Lip}(\gamma)}\|h\|_{s_0}^{\text{Lip}(\gamma)},$$

and (8.124) follows. The last inequality in (8.124) follows by (6.9) and (6.4).  $\square$

## 9 The Nash-Moser nonlinear iteration

In this section we prove Theorem 5.1. It will be a consequence of the Nash-Moser Theorem 9.1 below.

Consider the finite-dimensional subspaces

$$E_n := \{\mathcal{J}(\varphi) = (\Theta, y, z)(\varphi) : \Theta = \Pi_n \Theta, y = \Pi_n y, z = \Pi_n z\}$$

where  $N_n := N_0^{\chi_n}$  are introduced in (8.114), and  $\Pi_n$  are the projectors (which, with a small abuse of notation, we denote with the same symbol)

$$\begin{aligned} \Pi_n \Theta(\varphi) &:= \sum_{|l| < N_n} \Theta_l e^{il \cdot \varphi}, \quad \Pi_n y(\varphi) := \sum_{|l| < N_n} y_l e^{il \cdot \varphi}, \quad \text{where } \Theta(\varphi) = \sum_{l \in \mathbb{Z}^\nu} \Theta_l e^{il \cdot \varphi}, \quad y(\varphi) = \sum_{l \in \mathbb{Z}^\nu} y_l e^{il \cdot \varphi}, \\ \Pi_n z(\varphi, x) &:= \sum_{|(l,j)| < N_n} z_{lj} e^{i(l \cdot \varphi + jx)}, \quad \text{where } z(\varphi, x) = \sum_{l \in \mathbb{Z}^\nu, j \in S^c} z_{lj} e^{i(l \cdot \varphi + jx)}. \end{aligned} \quad (9.1)$$

We define  $\Pi_n^\perp := I - \Pi_n$ . The classical smoothing properties hold: for all  $\alpha, s \geq 0$ ,

$$\|\Pi_n \mathcal{J}\|_{s+\alpha}^{\text{Lip}(\gamma)} \leq N_n^\alpha \|\mathcal{J}\|_s^{\text{Lip}(\gamma)}, \quad \forall \mathcal{J}(\omega) \in H^s, \quad \|\Pi_n^\perp \mathcal{J}\|_s^{\text{Lip}(\gamma)} \leq N_n^{-\alpha} \|\mathcal{J}\|_{s+\alpha}^{\text{Lip}(\gamma)}, \quad \forall \mathcal{J}(\omega) \in H^{s+\alpha}. \quad (9.2)$$

We define the constants

$$\mu_1 := 3\mu + 9, \quad \alpha := 3\mu_1 + 1, \quad \alpha_1 := (\alpha - 3\mu)/2, \quad (9.3)$$

$$\kappa := 3(\mu_1 + \rho^{-1}) + 1, \quad \beta_1 := 6\mu_1 + 3\rho^{-1} + 3, \quad 0 < \rho < \frac{1-3a}{C_1(1+a)}, \quad (9.4)$$

where  $\mu := \mu(\tau, \nu)$  is the ‘‘loss of regularity’’ defined in Theorem 6.10 (see (6.40)) and  $C_1$  is fixed below.

**Theorem 9.1. (Nash-Moser)** Assume that  $f \in C^q$  with  $q > S := s_0 + \beta_1 + \mu + 3$ . Let  $\tau \geq \nu + 2$ . Then there exist  $C_1 > \max\{\mu_1 + \alpha, C_0\}$  (where  $C_0 := C_0(\tau, \nu)$  is the one in Theorem 8.25),  $\delta_0 := \delta_0(\tau, \nu) > 0$  such that, if

$$N_0^{C_1} \varepsilon^{b_*+1} \gamma^{-2} < \delta_0, \quad \gamma := \varepsilon^{2+a} = \varepsilon^{2b}, \quad N_0 := (\varepsilon \gamma^{-1})^\rho, \quad b_* := 6 - 2b, \quad (9.5)$$

then, for all  $n \geq 0$ :

(P1) $_n$  there exists a function  $(\mathfrak{J}_n, \zeta_n) : \mathcal{G}_n \subseteq \Omega_\varepsilon \rightarrow E_{n-1} \times \mathbb{R}^\nu$ ,  $\omega \mapsto (\mathfrak{J}_n(\omega), \zeta_n(\omega))$ ,  $(\mathfrak{J}_0, \zeta_0) := 0$ ,  $E_{-1} := \{0\}$ , satisfying  $|\zeta_n|^{\text{Lip}(\gamma)} \leq C \|\mathcal{F}(U_n)\|_{s_0}^{\text{Lip}(\gamma)}$ ,

$$\|\mathfrak{J}_n\|_{s_0+\mu}^{\text{Lip}(\gamma)} \leq C_* \varepsilon^{b_*} \gamma^{-1}, \quad \|\mathcal{F}(U_n)\|_{s_0+\mu+3}^{\text{Lip}(\gamma)} \leq C_* \varepsilon^{b_*}, \quad (9.6)$$

where  $U_n := (i_n, \zeta_n)$  with  $i_n(\varphi) = (\varphi, 0, 0) + \mathfrak{J}_n(\varphi)$ . The sets  $\mathcal{G}_n$  are defined inductively by:

$$\mathcal{G}_0 := \{\omega \in \Omega_\varepsilon : |\omega \cdot l| \geq 2\gamma \langle l \rangle^{-\tau}, \forall l \in \mathbb{Z}^\nu \setminus \{0\}\},$$

$$\mathcal{G}_{n+1} := \left\{ \omega \in \mathcal{G}_n : |\omega \cdot l + \mu_j^\infty(i_n) - \mu_k^\infty(i_n)| \geq \frac{2\gamma_n |j^3 - k^3|}{\langle l \rangle^\tau}, \forall j, k \in S^c \cup \{0\}, l \in \mathbb{Z}^\nu \right\}, \quad (9.7)$$

where  $\gamma_n := \gamma(1 + 2^{-n})$  and  $\mu_j^\infty(\omega) := \mu_j^\infty(\omega, i_n(\omega))$  are defined in (8.118) (and  $\mu_0^\infty(\omega) = 0$ ).

The differences  $\widehat{\mathfrak{J}}_n := \mathfrak{J}_n - \mathfrak{J}_{n-1}$  (where we set  $\widehat{\mathfrak{J}}_0 := 0$ ) is defined on  $\mathcal{G}_n$ , and satisfy

$$\|\widehat{\mathfrak{J}}_1\|_{s_0+\mu}^{\text{Lip}(\gamma)} \leq C_* \varepsilon^{b_*} \gamma^{-1}, \quad \|\widehat{\mathfrak{J}}_n\|_{s_0+\mu}^{\text{Lip}(\gamma)} \leq C_* \varepsilon^{b_*} \gamma^{-1} N_{n-1}^{-\alpha_1}, \quad \forall n > 1. \quad (9.8)$$

(P2) $_n$   $\|\mathcal{F}(U_n)\|_{s_0}^{\text{Lip}(\gamma)} \leq C_* \varepsilon^{b_*} N_{n-1}^{-\alpha}$  where we set  $N_{-1} := 1$ .

(P3) $_n$  (High norms).  $\|\mathfrak{J}_n\|_{s_0+\beta_1}^{\text{Lip}(\gamma)} \leq C_* \varepsilon^{b_*} \gamma^{-1} N_{n-1}^\kappa$  and  $\|\mathcal{F}(U_n)\|_{s_0+\beta_1}^{\text{Lip}(\gamma)} \leq C_* \varepsilon^{b_*} N_{n-1}^\kappa$ .

(P4) $_n$  (Measure). The measure of the ‘‘Cantor-like’’ sets  $\mathcal{G}_n$  satisfies

$$|\Omega_\varepsilon \setminus \mathcal{G}_0| \leq C_* \varepsilon^{2(\nu-1)} \gamma, \quad |\mathcal{G}_n \setminus \mathcal{G}_{n+1}| \leq C_* \varepsilon^{2(\nu-1)} \gamma N_{n-1}^{-1}. \quad (9.9)$$

All the Lip norms are defined on  $\mathcal{G}_n$ , namely  $\|\cdot\|_s^{\text{Lip}(\gamma)} = \|\cdot\|_{s, \mathcal{G}_n}^{\text{Lip}(\gamma)}$ .

*Proof.* To simplify notations, in this proof we denote  $\|\cdot\|_s^{\text{Lip}(\gamma)}$  by  $\|\cdot\|$ . We first prove (P1, 2, 3) $_n$ .

STEP 1: *Proof of (P1, 2, 3) $_0$ .* Recalling (5.6) we have  $\|\mathcal{F}(U_0)\|_s = \|\mathcal{F}(\varphi, 0, 0, 0)\|_s = \|X_P(\varphi, 0, 0)\|_s \leq_s \varepsilon^{6-2b}$  by (5.15). Hence (recall that  $b_* = 6 - 2b$ ) the smallness conditions in (P1) $_0$ –(P3) $_0$  hold taking  $C_* := C_*(s_0 + \beta_1)$  large enough.

STEP 2: Assume that (P1, 2, 3) $_n$  hold for some  $n \geq 0$ , and prove (P1, 2, 3) $_{n+1}$ . By (9.5) and (9.4),

$$N_0^{C_1} \varepsilon^{b_*+1} \gamma^{-2} = N_0^{C_1} \varepsilon^{1-3a} = \varepsilon^{1-3a-\rho C_1(1+a)} < \delta_0$$

for  $\varepsilon$  small enough, and the smallness condition (8.117) holds. Moreover (9.6) imply (6.4) (and so (7.8)) and Theorem 8.27 applies. Hence the operator  $\mathcal{L}_\omega := \mathcal{L}_\omega(\omega, i_n(\omega))$  defined in (6.31) is invertible for all  $\omega \in \mathcal{G}_{n+1}$  and the last estimate in (8.124) holds. This means that the assumption (6.32) of Theorem 6.10 is verified with  $\Omega_\infty = \mathcal{G}_{n+1}$ . By Theorem 6.10 there exists an approximate inverse  $\mathbf{T}_n(\omega) := \mathbf{T}_0(\omega, i_n(\omega))$  of the linearized operator  $L_n(\omega) := d_{i, \zeta} \mathcal{F}(\omega, i_n(\omega))$ , satisfying (6.40). Thus, using also (9.5), (9.2), (9.6),

$$\|\mathbf{T}_n g\|_s \leq_s \gamma^{-1} (\|g\|_{s+\mu} + \varepsilon \gamma^{-1} \{\|\mathfrak{J}_n\|_{s+\mu} + \gamma^{-1} \|\mathfrak{J}_n\|_{s_0+\mu} \|\mathcal{F}(U_n)\|_{s+\mu}\} \|g\|_{s_0+\mu}) \quad (9.10)$$

$$\|\mathbf{T}_n g\|_{s_0} \leq_{s_0} \gamma^{-1} \|g\|_{s_0+\mu} \quad (9.11)$$

and, by (6.41), using also (9.6), (9.5), (9.2),

$$\begin{aligned} \|(L_n \circ \mathbf{T}_n - I)g\|_s &\leq_s \gamma^{-1}(\|\mathcal{F}(U_n)\|_{s_0+\mu}\|g\|_{s+\mu} + \|\mathcal{F}(U_n)\|_{s+\mu}\|g\|_{s_0+\mu} \\ &\quad + \varepsilon\gamma^{-1}\|\mathfrak{J}_n\|_{s+\mu}\|\mathcal{F}(U_n)\|_{s_0+\mu}\|g\|_{s_0+\mu}), \end{aligned} \quad (9.12)$$

$$\begin{aligned} \|(L_n \circ \mathbf{T}_n - I)g\|_{s_0} &\leq_{s_0} \gamma^{-1}\|\mathcal{F}(U_n)\|_{s_0+\mu}\|g\|_{s_0+\mu} \\ &\leq_{s_0} \gamma^{-1}(\|\Pi_n\mathcal{F}(U_n)\|_{s_0+\mu} + \|\Pi_n^\perp\mathcal{F}(U_n)\|_{s_0+\mu})\|g\|_{s_0+\mu} \\ &\leq_{s_0} N_n^\mu\gamma^{-1}(\|\mathcal{F}(U_n)\|_{s_0} + N_n^{-\beta_1}\|\mathcal{F}(U_n)\|_{s_0+\beta_1})\|g\|_{s_0+\mu}. \end{aligned} \quad (9.13)$$

Then, for all  $\omega \in \mathcal{G}_{n+1}$ ,  $n \geq 0$ , we define

$$U_{n+1} := U_n + H_{n+1}, \quad H_{n+1} := (\widehat{\mathfrak{J}}_{n+1}, \widehat{\zeta}_{n+1}) := -\widetilde{\Pi}_n \mathbf{T}_n \Pi_n \mathcal{F}(U_n) \in E_n \times \mathbb{R}^\nu, \quad (9.14)$$

where  $\widetilde{\Pi}_n(\mathfrak{J}, \zeta) := (\Pi_n \mathfrak{J}, \zeta)$  with  $\Pi_n$  in (9.1). Since  $L_n := d_{i,\zeta} \mathcal{F}(i_n)$ , we write  $\mathcal{F}(U_{n+1}) = \mathcal{F}(U_n) + L_n H_{n+1} + Q_n$ , where

$$Q_n := Q(U_n, H_{n+1}), \quad Q(U_n, H) := \mathcal{F}(U_n + H) - \mathcal{F}(U_n) - L_n H, \quad H \in E_n \times \mathbb{R}^\nu. \quad (9.15)$$

Then, by the definition of  $H_{n+1}$  in (9.14), and writing  $\widetilde{\Pi}_n^\perp(\mathfrak{J}, \zeta) := (\Pi_n^\perp \mathfrak{J}, 0)$ , we have

$$\begin{aligned} \mathcal{F}(U_{n+1}) &= \mathcal{F}(U_n) - L_n \widetilde{\Pi}_n \mathbf{T}_n \Pi_n \mathcal{F}(U_n) + Q_n = \mathcal{F}(U_n) - L_n \mathbf{T}_n \Pi_n \mathcal{F}(U_n) + L_n \widetilde{\Pi}_n^\perp \mathbf{T}_n \Pi_n \mathcal{F}(U_n) + Q_n \\ &= \mathcal{F}(U_n) - \Pi_n L_n \mathbf{T}_n \Pi_n \mathcal{F}(U_n) + (L_n \widetilde{\Pi}_n^\perp - \Pi_n^\perp L_n) \mathbf{T}_n \Pi_n \mathcal{F}(U_n) + Q_n \\ &= \Pi_n^\perp \mathcal{F}(U_n) + R_n + Q_n + Q'_n \end{aligned} \quad (9.16)$$

where

$$R_n := (L_n \widetilde{\Pi}_n^\perp - \Pi_n^\perp L_n) \mathbf{T}_n \Pi_n \mathcal{F}(U_n), \quad Q'_n := -\Pi_n (L_n \mathbf{T}_n - I) \Pi_n \mathcal{F}(U_n). \quad (9.17)$$

**Lemma 9.2.** *Define*

$$w_n := \varepsilon\gamma^{-2}\|\mathcal{F}(U_n)\|_{s_0}, \quad B_n := \varepsilon\gamma^{-1}\|\mathfrak{J}_n\|_{s_0+\beta_1} + \varepsilon\gamma^{-2}\|\mathcal{F}(U_n)\|_{s_0+\beta_1}. \quad (9.18)$$

Then there exists  $K := K(s_0, \beta_1) > 0$  such that, for all  $n \geq 0$ , setting  $\mu_1 := 3\mu + 9$  (see (9.3)),

$$w_{n+1} \leq K N_n^{\mu_1 + \frac{1}{p} - \beta_1} B_n + K N_n^{\mu_1} w_n^2, \quad B_{n+1} \leq K N_n^{\mu_1 + \frac{1}{p}} B_n. \quad (9.19)$$

*Proof.* We estimate separately the terms  $Q_n$  in (9.15) and  $Q'_n, R_n$  in (9.17).

*Estimate of  $Q_n$ .* By (9.15), (5.6), (5.20) and (9.6), (9.2), we have the quadratic estimates

$$\|Q(U_n, H)\|_s \leq_s \varepsilon(\|\widehat{\mathfrak{J}}\|_{s+3}\|\widehat{\mathfrak{J}}\|_{s_0+3} + \|\mathfrak{J}_n\|_{s+3}\|\widehat{\mathfrak{J}}\|_{s_0+3}^2) \quad (9.20)$$

$$\|Q(U_n, H)\|_{s_0} \leq_{s_0} \varepsilon N_n^6 \|\widehat{\mathfrak{J}}\|_{s_0}^2, \quad \forall \widehat{\mathfrak{J}} \in E_n. \quad (9.21)$$

Now by the definition of  $H_{n+1}$  in (9.14) and (9.2), (9.10), (9.11), (9.6), we get

$$\begin{aligned} \|\widehat{\mathfrak{J}}_{n+1}\|_{s_0+\beta_1} &\leq_{s_0+\beta_1} N_n^\mu(\gamma^{-1}\|\mathcal{F}(U_n)\|_{s_0+\beta_1} + \varepsilon\gamma^{-2}\|\mathcal{F}(U_n)\|_{s_0+\mu}\{\|\mathfrak{J}_n\|_{s_0+\beta_1} + \gamma^{-1}\|\mathcal{F}(U_n)\|_{s_0+\beta_1}\}) \\ &\leq_{s_0+\beta_1} N_n^\mu(\gamma^{-1}\|\mathcal{F}(U_n)\|_{s_0+\beta_1} + \|\mathfrak{J}_n\|_{s_0+\beta_1}), \end{aligned} \quad (9.22)$$

$$\|\widehat{\mathfrak{J}}_{n+1}\|_{s_0} \leq_{s_0} \gamma^{-1} N_n^\mu \|\mathcal{F}(U_n)\|_{s_0}. \quad (9.23)$$

Then the term  $Q_n$  in (9.15) satisfies, by (9.20), (9.21), (9.22), (9.23), (9.5), (9.6),  $(\mathcal{P}2)_n$ , (9.3),

$$\|Q_n\|_{s_0+\beta_1} \leq_{s_0+\beta_1} N_n^{2\mu+9} \gamma(\gamma^{-1}\|\mathcal{F}(U_n)\|_{s_0+\beta_1} + \|\mathfrak{J}_n\|_{s_0+\beta_1}), \quad (9.24)$$

$$\|Q_n\|_{s_0} \leq_{s_0} N_n^{2\mu+6} \varepsilon\gamma^{-2} \|\mathcal{F}(U_n)\|_{s_0}^2. \quad (9.25)$$

*Estimate of  $\tilde{Q}'_n$ .* The bounds (9.12), (9.13), (9.2), (9.3), (9.6) imply

$$\|Q'_n\|_{s_0+\beta_1} \leq_{s_0+\beta_1} N_n^{2\mu} (\|\mathcal{F}(U_n)\|_{s_0+\beta_1} + \|\mathfrak{J}_n\|_{s_0+\beta_1} \|\mathcal{F}(U_n)\|_{s_0}), \quad (9.26)$$

$$\|Q'_n\|_{s_0} \leq_{s_0} \gamma^{-1} N_n^{2\mu} (\|\mathcal{F}(U_n)\|_{s_0} + N_n^{-\beta_1} \|\mathcal{F}(U_n)\|_{s_0+\beta_1}) \|\mathcal{F}(U_n)\|_{s_0}. \quad (9.27)$$

*Estimate of  $R_n$ .* For  $H := (\widehat{\mathcal{J}}, \widehat{\zeta})$  we have  $(L_n \widetilde{\Pi}_n^\perp - \Pi_n^\perp L_n)H = [\bar{D}_n, \Pi_n^\perp] \widehat{\mathcal{J}} = [\Pi_n, \bar{D}_n] \widehat{\mathcal{J}}$  where  $\bar{D}_n := d_i X_{H_\varepsilon}(i_n) + (0, 0, \partial_{xxx})$ . Thus Lemma 5.3, (9.6), (9.2) and (5.19) imply

$$\|(L_n \widetilde{\Pi}_n^\perp - \Pi_n^\perp L_n)H\|_{s_0} \leq_{s_0+\beta_1} \varepsilon N_n^{-\beta_1+\mu+3} (\|\widehat{\mathcal{J}}\|_{s_0+\beta_1-\mu} + \|\mathfrak{J}_n\|_{s_0+\beta_1-\mu} \|\widehat{\mathcal{J}}\|_{s_0+3}), \quad (9.28)$$

$$\|(L_n \widetilde{\Pi}_n^\perp - \Pi_n^\perp L_n)H\|_{s_0+\beta_1} \leq_s \varepsilon N_n^{\mu+3} (\|\widehat{\mathcal{J}}\|_{s_0+\beta_1-\mu} + \|\mathfrak{J}_n\|_{s_0+\beta_1-\mu} \|\widehat{\mathcal{J}}\|_{s_0+3}). \quad (9.29)$$

Hence, applying (9.10), (9.28), (9.29), (9.5), (9.6), (9.2), the term  $R_n$  defined in (9.17) satisfies

$$\|R_n\|_{s_0} \leq_{s_0+\beta_1} N_n^{\mu+6-\beta_1} (\varepsilon \gamma^{-1} \|\mathcal{F}(U_n)\|_{s_0+\beta_1} + \varepsilon \|\mathfrak{J}_n\|_{s_0+\beta_1}), \quad (9.30)$$

$$\|R_n\|_{s_0+\beta_1} \leq_{s_0+\beta_1} N_n^{\mu+6} (\varepsilon \gamma^{-1} \|\mathcal{F}(U_n)\|_{s_0+\beta_1} + \varepsilon \|\mathfrak{J}_n\|_{s_0+\beta_1}). \quad (9.31)$$

*Estimate of  $\mathcal{F}(U_{n+1})$ .* By (9.16) and (9.24), (9.25), (9.26), (9.27), (9.30), (9.31), (9.5), (9.6), we get

$$\|\mathcal{F}(U_{n+1})\|_{s_0} \leq_{s_0+\beta_1} N_n^{\mu_1-\beta_1} (\varepsilon \gamma^{-1} \|\mathcal{F}(U_n)\|_{s_0+\beta_1} + \varepsilon \|\mathfrak{J}_n\|_{s_0+\beta_1}) + N_n^{\mu_1} \varepsilon \gamma^{-2} \|\mathcal{F}(U_n)\|_{s_0}^2, \quad (9.32)$$

$$\|\mathcal{F}(U_{n+1})\|_{s_0+\beta_1} \leq_{s_0+\beta_1} N_n^{\mu_1} (\varepsilon \gamma^{-1} \|\mathcal{F}(U_n)\|_{s_0+\beta_1} + \varepsilon \|\mathfrak{J}_n\|_{s_0+\beta_1}), \quad (9.33)$$

where  $\mu_1 := 3\mu + 9$ .

*Estimate of  $\mathfrak{J}_{n+1}$ .* Using (9.22) the term  $\mathfrak{J}_{n+1} = \mathfrak{J}_n + \widehat{\mathfrak{J}}_{n+1}$  is bounded by

$$\|\mathfrak{J}_{n+1}\|_{s_0+\beta_1} \leq_{s_0+\beta_1} N_n^\mu (\|\mathfrak{J}_n\|_{s_0+\beta_1} + \gamma^{-1} \|\mathcal{F}(U_n)\|_{s_0+\beta_1}). \quad (9.34)$$

Finally, recalling (9.18), the inequalities (9.19) follow by (9.32)-(9.34), (9.6) and  $\varepsilon \gamma^{-1} = N_0^{1/\rho} \leq N_n^{1/\rho}$ .  $\square$

*Proof of  $(\mathcal{P}3)_{n+1}$ .* By (9.19) and  $(\mathcal{P}3)_n$ ,

$$B_{n+1} \leq K N_n^{\mu_1+\frac{1}{\rho}} B_n \leq 2C_* K \varepsilon^{b_*+1} \gamma^{-2} N_n^{\mu_1+\frac{1}{\rho}} N_{n-1}^\kappa \leq C_* \varepsilon^{b_*+1} \gamma^{-2} N_n^\kappa, \quad (9.35)$$

provided  $2K N_n^{\mu_1+\frac{1}{\rho}-\kappa} N_{n-1}^\kappa \leq 1$ ,  $\forall n \geq 0$ . This inequality holds by (9.4), taking  $N_0$  large enough (i.e  $\varepsilon$  small enough). By (9.18), the bound  $B_{n+1} \leq C_* \varepsilon^{b_*+1} \gamma^{-2} N_n^\kappa$  implies  $(\mathcal{P}3)_{n+1}$ .

*Proof of  $(\mathcal{P}2)_{n+1}$ .* Using (9.19), (9.18) and  $(\mathcal{P}2)_n, (\mathcal{P}3)_n$ , we get

$$w_{n+1} \leq K N_n^{\mu_1+\frac{1}{\rho}-\beta_1} B_n + K N_n^{\mu_1} w_n^2 \leq K N_n^{\mu_1+\frac{1}{\rho}-\beta_1} 2C_* \varepsilon^{b_*+1} \gamma^{-2} N_{n-1}^\kappa + K N_n^{\mu_1} (C_* \varepsilon^{b_*+1} \gamma^{-2} N_{n-1}^{-\alpha})^2$$

which is  $\leq C_* \varepsilon^{b_*+1} \gamma^{-2} N_n^{-\alpha}$  provided that

$$4K N_n^{\mu_1+\frac{1}{\rho}-\beta_1+\alpha} N_{n-1}^\kappa \leq 1, \quad 2K C_* \varepsilon^{b_*+1} \gamma^{-2} N_n^{\mu_1+\alpha} N_{n-1}^{-2\alpha} \leq 1, \quad \forall n \geq 0. \quad (9.36)$$

The inequalities in (9.36) hold by (9.3)-(9.4), (9.5),  $C_1 > \mu_1 + \alpha$ , taking  $\delta_0$  in (9.5) small enough. By (9.18), the inequality  $w_{n+1} \leq C_* \varepsilon^{b_*+1} \gamma^{-2} N_n^{-\alpha}$  implies  $(\mathcal{P}2)_{n+1}$ .

*Proof of  $(\mathcal{P}1)_{n+1}$ .* The bound (9.8) for  $\widehat{\mathcal{J}}_1$  follows by (9.14), (9.10) (for  $s = s_0 + \mu$ ) and  $\|\mathcal{F}(U_0)\|_{s_0+2\mu} = \|\mathcal{F}(\varphi, 0, 0, 0)\|_{s_0+2\mu} \leq_{s_0+2\mu} \varepsilon^{b_*}$ . The bound (9.8) for  $\widehat{\mathcal{J}}_{n+1}$  follows by (9.2), (9.23),  $(\mathcal{P}2)_n, (\mathcal{P}3)_n$ . It remains to prove that (9.6) holds at the step  $n+1$ . We have

$$\|\mathfrak{J}_{n+1}\|_{s_0+\mu} \leq \sum_{k=1}^{n+1} \|\widehat{\mathcal{J}}_k\|_{s_0+\mu} \leq C_* \varepsilon^{b_*} \gamma^{-1} \sum_{k \geq 1} N_{k-1}^{-\alpha_1} \leq C_* \varepsilon^{b_*} \gamma^{-1} \quad (9.37)$$

for  $N_0$  large enough, i.e.  $\varepsilon$  small. Moreover, using (9.2),  $(\mathcal{P}2)_{n+1}, (\mathcal{P}3)_{n+1}, (9.3)$ , we get

$$\begin{aligned} \|\mathcal{F}(U_{n+1})\|_{s_0+\mu+3} &\leq N_n^{\mu+3} \|\mathcal{F}(U_{n+1})\|_{s_0} + N_n^{\mu+3-\beta_1} \|\mathcal{F}(U_{n+1})\|_{s_0+\beta_1} \\ &\leq C_* \varepsilon^{b_*} N_n^{\mu+3-\alpha} + C_* \varepsilon^{b_*} N_n^{\mu+3-\beta_1+\kappa} \leq C_* \varepsilon^{b_*}, \end{aligned}$$

which is the second inequality in (9.6) at the step  $n + 1$ . The bound  $|\zeta_{n+1}|^{\text{Lip}(\gamma)} \leq C \|\mathcal{F}(U_{n+1})\|_{s_0}^{\text{Lip}(\gamma)}$  is a consequence of Lemma 6.1 (it is not inductive).

STEP 3: Prove  $(\mathcal{P}4)_n$  for all  $n \geq 0$ . For all  $n \geq 0$ ,

$$\mathcal{G}_n \setminus \mathcal{G}_{n+1} = \bigcup_{l \in \mathbb{Z}^\nu, j, k \in S^c \cup \{0\}} R_{ljk}(i_n) \quad (9.38)$$

where

$$R_{ljk}(i_n) := \{\omega \in \mathcal{G}_n : |\omega \cdot l + \mu_j^\infty(i_n) - \mu_k^\infty(i_n)| < 2\gamma_n |j^3 - k^3| \langle l \rangle^{-\tau}\}. \quad (9.39)$$

Notice that  $R_{ljk}(i_n) = \emptyset$  if  $j = k$ , so that we suppose in the sequel that  $j \neq k$ .

**Lemma 9.3.** For all  $n \geq 1$ ,  $|l| \leq N_{n-1}$ , the set  $R_{ljk}(i_n) \subseteq R_{ljk}(i_{n-1})$ .

*Proof.* Like Lemma 5.2 in [2] (with  $\omega$  in the role of  $\lambda\bar{\omega}$ , and  $N_{n-1}$  instead of  $N_n$ ).  $\square$

By definition,  $R_{ljk}(i_n) \subseteq \mathcal{G}_n$  (see (9.39)) and Lemma 9.3 implies that, for all  $n \geq 1$ ,  $|l| \leq N_{n-1}$ , the set  $R_{ljk}(i_n) \subseteq R_{ljk}(i_{n-1})$ . On the other hand  $R_{ljk}(i_{n-1}) \cap \mathcal{G}_n = \emptyset$  (see (9.7)). As a consequence, for all  $|l| \leq N_{n-1}$ ,  $R_{ljk}(i_n) = \emptyset$  and, by (9.38),

$$\mathcal{G}_n \setminus \mathcal{G}_{n+1} \subseteq \bigcup_{|l| > N_{n-1}, j, k \in S^c \cup \{0\}} R_{ljk}(i_n) \quad \forall n \geq 1. \quad (9.40)$$

**Lemma 9.4.** Let  $n \geq 0$ . If  $R_{ljk}(i_n) \neq \emptyset$  then  $|l| \geq C|j^3 - k^3| \geq \frac{1}{2}C(j^2 + k^2)$  for some  $C > 0$ .

*Proof.* Like Lemma 5.3 in [2]. The only difference is that  $\omega$  is not constrained to a fixed direction. Note also that  $|j^3 - k^3| \geq (j^2 + k^2)/2$ ,  $\forall j \neq k$ .  $\square$

By usual arguments (e.g. see Lemma 5.4 in [2]), using Lemma 9.4 and (8.119) we have:

**Lemma 9.5.** For all  $n \geq 0$ , the measure  $|R_{ljk}(i_n)| \leq C\varepsilon^{2(\nu-1)}\gamma \langle l \rangle^{-\tau}$ .

By (9.38) and Lemmata 9.4, 9.5 we get

$$|\mathcal{G}_0 \setminus \mathcal{G}_1| \leq \sum_{l \in \mathbb{Z}^\nu, |j|, |k| \leq C|l|^{1/2}} |R_{ljk}(i_0)| \leq \sum_{l \in \mathbb{Z}^\nu} \frac{C\varepsilon^{2(\nu-1)}\gamma}{\langle l \rangle^{\tau-1}} \leq C'\varepsilon^{2(\nu-1)}\gamma.$$

For  $n \geq 1$ , by (9.40),

$$|\mathcal{G}_n \setminus \mathcal{G}_{n+1}| \leq \sum_{|l| > N_{n-1}, |j|, |k| \leq C|l|^{1/2}} |R_{ljk}(i_n)| \leq \sum_{|l| > N_{n-1}} \frac{C\varepsilon^{2(\nu-1)}\gamma}{\langle l \rangle^{\tau-1}} \leq C'\varepsilon^{2(\nu-1)}\gamma N_{n-1}^{-1}$$

because  $\tau \geq \nu + 2$ . The estimate  $|\Omega_\varepsilon \setminus \mathcal{G}_0| \leq C\varepsilon^{2(\nu-1)}\gamma$  is elementary. Thus (9.9) is proved.  $\square$

**Proof of Theorem 5.1 concluded.** Theorem 9.1 implies that the sequence  $(\mathcal{I}_n, \zeta_n)$  is well defined for  $\omega \in \mathcal{G}_\infty := \bigcap_{n \geq 0} \mathcal{G}_n$ , that  $\mathcal{I}_n$  is a Cauchy sequence in  $\|\cdot\|_{s_0 + \mu, \mathcal{G}_\infty}^{\text{Lip}(\gamma)}$ , see (9.8), and  $|\zeta_n|^{\text{Lip}(\gamma)} \rightarrow 0$ . Therefore  $\mathcal{I}_n$  converges to a limit  $\mathcal{I}_\infty$  in norm  $\|\cdot\|_{s_0 + \mu, \mathcal{G}_\infty}^{\text{Lip}(\gamma)}$  and, by  $(\mathcal{P}2)_n$ , for all  $\omega \in \mathcal{G}_\infty$ ,  $i_\infty(\varphi) := (\varphi, 0, 0) + \mathcal{I}_\infty(\varphi)$ , is a solution of

$$\mathcal{F}(i_\infty, 0) = 0 \quad \text{with} \quad \|\mathcal{I}_\infty\|_{s_0 + \mu, \mathcal{G}_\infty}^{\text{Lip}(\gamma)} \leq C\varepsilon^{6-2b}\gamma^{-1}$$

by (9.6) (recall that  $b_* := 6 - 2b$ ). Therefore  $\varphi \mapsto i_\infty(\varphi)$  is an invariant torus for the Hamiltonian vector field  $X_{H_\varepsilon}$  (see (5.5)). By (9.9),

$$|\Omega_\varepsilon \setminus \mathcal{G}_\infty| \leq |\Omega_\varepsilon \setminus \mathcal{G}_0| + \sum_{n \geq 0} |\mathcal{G}_n \setminus \mathcal{G}_{n+1}| \leq 2C_*\varepsilon^{2(\nu-1)}\gamma + C_*\varepsilon^{2(\nu-1)}\gamma \sum_{n \geq 1} N_{n-1}^{-1} \leq C\varepsilon^{2(\nu-1)}\gamma.$$

The set  $\Omega_\varepsilon$  in (5.2) has measure  $|\Omega_\varepsilon| = O(\varepsilon^{2\nu})$ . Hence  $|\Omega_\varepsilon \setminus \mathcal{G}_\infty|/|\Omega_\varepsilon| \rightarrow 0$  as  $\varepsilon \rightarrow 0$  because  $\gamma = o(\varepsilon^2)$ , and therefore the measure of  $\mathcal{C}_\varepsilon := \mathcal{G}_\infty$  satisfies (5.11).

In order to complete the proof of Theorem 5.1 we show the linear stability of the solution  $i_\infty(\omega t)$ . By section 6 the system obtained linearizing the Hamiltonian vector field  $X_{H_\varepsilon}$  at a quasi-periodic solution  $i_\infty(\omega t)$  is conjugated to the linear Hamiltonian system

$$\begin{cases} \dot{\psi} & = K_{20}(\omega t)\eta + K_{11}^T(\omega t)w \\ \dot{\eta} & = 0 \\ \dot{w} - \partial_x K_{02}(\omega t)w & = \partial_x K_{11}(\omega t)\eta \end{cases} \quad (9.41)$$

(recall that the torus  $i_\infty$  is isotropic and the transformed nonlinear Hamiltonian system is (6.20) where  $K_{00}, K_{10}, K_{01} = 0$ , see Remark 6.5). In section 8 we have proved the reducibility of the linear system  $\dot{w} - \partial_x K_{02}(\omega t)w$ , conjugating the last equation in (9.41) to a diagonal system

$$\dot{v}_j + \mu_j^\infty v_j = f_j(\omega t), \quad j \in S^c, \quad \mu_j^\infty \in i\mathbb{R}, \quad (9.42)$$

see (8.121), and  $f(\varphi, x) = \sum_{j \in S^c} f_j(\varphi) e^{ijx} \in H_{S^\perp}^s(\mathbb{T}^{\nu+1})$ . Thus (9.41) is stable. Indeed the actions  $\eta(t) = \eta_0 \in \mathbb{R}, \forall t \in \mathbb{R}$ . Moreover the solutions of the non-homogeneous equation (9.42) are

$$v_j(t) = c_j e^{\mu_j^\infty t} + \tilde{v}_j(t), \quad \text{where} \quad \tilde{v}_j(t) := \sum_{l \in \mathbb{Z}^\nu} \frac{f_{jl} e^{i\omega \cdot l t}}{i\omega \cdot l + \mu_j^\infty}$$

is a quasi-periodic solution (recall that the first Melnikov conditions (8.123) hold at a solution). As a consequence (recall also  $\mu_j^\infty \in i\mathbb{R}$ ) the Sobolev norm of the solution of (9.42) with initial condition  $v(0) = \sum_{j \in S^c} v_j(0) e^{ijx} \in H^{s_0}(\mathbb{T}_x)$ ,  $s_0 < s$ , does not increase in time.  $\square$

**Construction of the set  $S$  of tangential sites.** We finally prove that, for any  $\nu \geq 1$ , the set  $S$  in (1.8) satisfying (S1)-(S2) can be constructed inductively with only a *finite* number of restriction at any step of the induction.

First, fix any integer  $\bar{j}_1 \geq 1$ . Then the set  $J_1 := \{\pm \bar{j}_1\}$  trivially satisfies (S1)-(S2). Then, assume that we have fixed  $n$  distinct positive integers  $\bar{j}_1, \dots, \bar{j}_n$ ,  $n \geq 1$ , such that the set  $J_n := \{\pm \bar{j}_1, \dots, \pm \bar{j}_n\}$  satisfies (S1)-(S2). We describe how to choose another positive integer  $\bar{j}_{n+1}$ , which is different from all  $j \in J_n$ , such that  $J_{n+1} := J_n \cup \{\pm \bar{j}_{n+1}\}$  also satisfies (S1), (S2).

Let us begin with analyzing (S1). A set of 3 elements  $j_1, j_2, j_3 \in J_{n+1}$  can be of these types: (i) all “old” elements  $j_1, j_2, j_3 \in J_n$ ; (ii) two “old” elements  $j_1, j_2 \in J_n$  and one “new” element  $j_3 = \sigma_3 \bar{j}_{n+1}$ ,  $\sigma_3 = \pm 1$ ; (iii) one “old” element  $j_1 \in J_n$  and two “new” elements  $j_2 = \sigma_2 \bar{j}_{n+1}$ ,  $j_3 = \sigma_3 \bar{j}_{n+1}$ , with  $\sigma_2, \sigma_3 = \pm 1$ ; (iv) all “new” elements  $j_i = \sigma_i \bar{j}_{n+1}$ ,  $\sigma_i = \pm 1$ ,  $i = 1, 2, 3$ .

In case (i), the sum  $j_1 + j_2 + j_3$  is nonzero by inductive assumption. In case (ii),  $j_1 + j_2 + j_3$  is nonzero provided  $\bar{j}_{n+1} \notin \{j_1 + j_2 : j_1, j_2 \in J_n\}$ , which is a finite set. In case (iii), for  $\sigma_2 + \sigma_3 = 0$  the sum  $j_1 + j_2 + j_3 = j_1$  is trivially nonzero because  $0 \notin J_n$ , while, for  $\sigma_2 + \sigma_3 \neq 0$ , the sum  $j_1 + j_2 + j_3 = j_1 + (\sigma_2 + \sigma_3) \bar{j}_{n+1} \neq 0$  if  $\bar{j}_{n+1} \notin \{\frac{1}{2}j : j \in J_n\}$ , which is a finite set. In case (iv), the sum  $j_1 + j_2 + j_3 = (\sigma_1 + \sigma_2 + \sigma_3) \bar{j}_{n+1} \neq 0$  because  $\bar{j}_{n+1} \geq 1$  and  $\sigma_1 + \sigma_2 + \sigma_3 \in \{\pm 1, \pm 3\}$ .

Now we study (S2) for the set  $J_{n+1}$ . Denote, in short,  $b := j_1^3 + j_2^3 + j_3^3 + j_4^3 - (j_1 + j_2 + j_3 + j_4)^3$ .

A set of 4 elements  $j_1, j_2, j_3, j_4 \in J_{n+1}$  can be of 5 types: (i) all “old” elements  $j_1, j_2, j_3, j_4 \in J_n$ ; (ii) three “old” elements  $j_1, j_2, j_3 \in J_n$  and one “new” element  $j_4 = \sigma_4 \bar{j}_{n+1}$ ,  $\sigma_4 = \pm 1$ ; (iii) two “old” element  $j_1, j_2 \in J_n$  and two “new” elements  $j_3 = \sigma_3 \bar{j}_{n+1}$ ,  $j_4 = \sigma_4 \bar{j}_{n+1}$ , with  $\sigma_3, \sigma_4 = \pm 1$ ; (iv) one “old” element  $j_1 \in J_n$  and three “new” elements  $j_i = \sigma_i \bar{j}_{n+1}$ ,  $\sigma_i = \pm 1$ ,  $i = 2, 3, 4$ ; (v) all “new” elements  $j_i = \sigma_i \bar{j}_{n+1}$ ,  $\sigma_i = \pm 1$ ,  $i = 1, 2, 3, 4$ .

In case (i),  $b \neq 0$  by inductive assumption.

In case (ii), assume that  $j_1 + j_2 + j_3 + j_4 \neq 0$ , and calculate

$$b = -3(j_1 + j_2 + j_3) \bar{j}_{n+1}^2 - 3(j_1 + j_2 + j_3)^2 \sigma_4 \bar{j}_{n+1} + [j_1^3 + j_2^3 + j_3^3 - (j_1 + j_2 + j_3)^3] =: p_{j_1, j_2, j_3, \sigma_4}(\bar{j}_{n+1}).$$

This is nonzero provided  $p_{j_1, j_2, j_3, \sigma_4}(\bar{j}_{n+1}) \neq 0$  for all  $j_1, j_2, j_3 \in J_n$ ,  $\sigma_4 = \pm 1$ . The polynomial  $p_{j_1, j_2, j_3, \sigma_4}$  is never identically zero because either the leading coefficient  $-3(j_1 + j_2 + j_3) \neq 0$  (and, if one uses (S3), this is always the case), or, if  $j_1 + j_2 + j_3 = 0$ , then  $j_1^3 + j_2^3 + j_3^3 \neq 0$  by (3.12) (using also that  $0 \notin J_n$ ).



In case (iii), assume that  $j_1 + \dots + j_4 = j_1 + j_2 + (\sigma_3 + \sigma_4)\bar{j}_{n+1} \neq 0$ , and calculate

$$b = -3\alpha\bar{j}_{n+1}^3 - 3\alpha^2(j_1 + j_2)\bar{j}_{n+1}^2 - 3(j_1 + j_2)^2\alpha\bar{j}_{n+1} - j_1j_2(j_1 + j_2) =: q_{j_1, j_2, \alpha}(\bar{j}_{n+1}),$$

where  $\alpha := \sigma_3 + \sigma_4$ . We impose that  $q_{j_1, j_2, \alpha}(\bar{j}_{n+1}) \neq 0$  for all  $j_1, j_2 \in J_n$ ,  $\alpha \in \{\pm 2, 0\}$ . The polynomial  $q_{j_1, j_2, \alpha}$  is never identically zero because either the leading coefficient  $-3\alpha \neq 0$ , or, for  $\alpha = 0$ , the constant term  $-j_1j_2(j_1 + j_2) \neq 0$  (recall that  $0 \notin J_n$  and  $j_1 + j_2 + \alpha\bar{j}_{n+1} \neq 0$ ).

In case (iv), assume that  $j_1 + \dots + j_4 = j_1 + \alpha\bar{j}_{n+1} \neq 0$ , where  $\alpha := \sigma_2 + \sigma_3 + \sigma_4 \in \{\pm 1, \pm 3\}$ , and calculate

$$b = \alpha\bar{j}_{n+1}r_{j_1, \alpha}(\bar{j}_{n+1}), \quad r_{j_1, \alpha}(x) := (1 - \alpha^2)x^2 - 3\alpha j_1 x - 3j_1^2.$$

The polynomial  $r_{j_1, \alpha}$  is never identically zero because  $j_1 \neq 0$ . We impose  $r_{j_1, \alpha}(\bar{j}_{n+1}) \neq 0$  for all  $j_1 \in J_n$ ,  $\alpha \in \{\pm 1, \pm 3\}$ .

In case (v), assume that  $j_1 + \dots + j_4 = \alpha\bar{j}_{n+1} \neq 0$ , with  $\alpha := \sigma_1 + \dots + \sigma_4 \neq 0$ , and calculate  $b = \alpha(1 - \alpha^2)\bar{j}_{n+1}^3$ . This is nonzero because  $\bar{j}_{n+1} \geq 1$  and  $\alpha \in \{\pm 2, \pm 4\}$ .

We have proved that, in choosing  $\bar{j}_{n+1}$ , there are only finitely many integers to avoid.

## References

- [1] Baldi P., *Periodic solutions of fully nonlinear autonomous equations of Benjamin-Ono type*, Ann. Inst. H. Poincaré (C) Anal. Non Linéaire 30 (2013), 33-77.
- [2] Baldi P., Berti M., Montalto R., *KAM for quasi-linear and fully nonlinear forced perturbations of Airy equation*, Math. Annalen, in print (published online 2014, DOI 10.1007/s00208-013-1001-7).
- [3] Berti M., Biasco P., Procesi M., *KAM theory for the Hamiltonian DNLS*, Ann. Sci. Éc. Norm. Supér. (4), Vol. 46, fascicule 2 (2013), 301-373.
- [4] Berti M., Biasco P., Procesi M., *KAM theory for the reversible derivative wave equation*, Arch. Rational Mech. Anal., 212, (2014), 905-955.
- [5] Berti M., Bolle P., *Quasi-periodic solutions with Sobolev regularity of NLS on  $\mathbb{T}^d$  with a multiplicative potential*, Eur. Jour. Math. 15 (2013), 229-286.
- [6] Berti M., Bolle P., *A Nash-Moser approach to KAM theory*, preprint 2014.
- [7] Berti M., Bolle P., *Quasi-periodic solutions for autonomous NLW on  $\mathbb{T}^d$  with a multiplicative potential*, in preparation.
- [8] Bourgain J., *Gibbs measures and quasi-periodic solutions for nonlinear Hamiltonian partial differential equations*, 23-43, Gelfand Math. Sem., Birkhäuser Boston, Boston, MA, 1996.
- [9] Bourgain J., *Green's function estimates for lattice Schrödinger operators and applications*, Annals of Mathematics Studies 158, Princeton University Press, Princeton, 2005.
- [10] Craig W., *Problèmes de petits diviseurs dans les équations aux dérivées partielles*, Panoramas et Synthèses, 9, Société Mathématique de France, Paris, 2000.
- [11] Craig W., Wayne C.E., *Newton's method and periodic solutions of nonlinear wave equation*, Comm. Pure Appl. Math. 46, 1409-1498, 1993.
- [12] Eliasson L.H., Kuksin S., *KAM for non-linear Schrödinger equation*, Annals of Math., 172 (2010), 371-435.
- [13] Geng J., Xu X., You J., *An infinite dimensional KAM theorem and its application to the two dimensional cubic Schrödinger equation*, Adv. Math. 226 (2011) 5361-5402.
- [14] Huang G., Kuksin S., *KdV equation under periodic boundary conditions and its perturbations*, preprint, <http://arxiv.org/abs/1309.1597>.

- [15] Iooss G., Plotnikov P.I., *Small divisor problem in the theory of three-dimensional water gravity waves*, Mem. Amer. Math. Soc. 200, no. 940 (2009).
- [16] Iooss G., Plotnikov P.I., *Asymmetrical three-dimensional travelling gravity waves*, Arch. Rational Mech. Anal. 200 no. 3, (2011), 789-880.
- [17] Iooss G., Plotnikov P.I., Toland J.F., *Standing waves on an infinitely deep perfect fluid under gravity*, Arch. Rational Mech. Anal. 177 no. 3, (2005), 367-478.
- [18] Lax P., *Development of singularities of solutions of nonlinear hyperbolic partial differential equations*, J. Mathematical Phys. 5 (1964), 611-613.
- [19] Liu J., Yuan X., *A KAM Theorem for Hamiltonian Partial Differential Equations with Unbounded Perturbations*, Comm. Math. Phys. 307 (3) (2011), 629-673.
- [20] Kappeler T., Pöschel J., *KAM and KdV*, Springer, 2003.
- [21] Klainerman S., Majda A., *Formation of singularities for wave equations including the nonlinear vibrating string*, Comm. Pure Appl. Math., 33, (1980), 241-263.
- [22] Kuksin S., *Hamiltonian perturbations of infinite-dimensional linear systems with imaginary spectrum*, Funktsional. Anal. i Prilozhen. 21, no. 3, 22-37, 95, 1987.
- [23] Kuksin S., *A KAM theorem for equations of the Korteweg-de Vries type*, Rev. Math. Phys., 10, 3, (1998), 1-64.
- [24] Kuksin S., *Analysis of Hamiltonian PDEs*, Oxford Lecture Series in Mathematics and its Applications, 19. Oxford University Press (2000).
- [25] Pöschel J., *A KAM-Theorem for some nonlinear PDEs*, Ann. Sc. Norm. Pisa, 23, (1996) 119-148.
- [26] Pöschel J., *Quasi-periodic solutions for a nonlinear wave equation*, Comment. Math. Helv., 71, no. 2, (1996) 269-296.
- [27] Procesi M., Procesi C., *A normal form for the Schrödinger equation with analytic non-linearities*, Comm. Math. Phys. 312 (2012), 501-557.
- [28] Procesi C., Procesi M., *A KAM algorithm for the completely resonant nonlinear Schrödinger equation*, preprint (2013).
- [29] Taylor M. E., *Pseudodifferential Operators and Nonlinear PDEs*, Progress in Mathematics, Birkhäuser, 1991.
- [30] Wang W. M., *Supercritical nonlinear Schrödinger equations I: quasi-periodic solutions*, preprint.
- [31] Wayne E., *Periodic and quasi-periodic solutions of nonlinear wave equations via KAM theory*, Comm. Math. Phys. 127, 479-528, 1990.
- [32] Zhang J., Gao M., Yuan X. *KAM tori for reversible partial differential equations*, Nonlinearity 24 (2011), 1189-1228.
- [33] Zehnder E., *Generalized implicit function theorems with applications to some small divisors problems I-II*, Comm. Pure Appl. Math. 28 (1975), 91-140, and 29 (1976), 49-113.

Pietro Baldi, Dipartimento di Matematica e Applicazioni “R. Caccioppoli”, Università di Napoli Federico II, Via Cintia, Monte S. Angelo, 80126, Napoli, Italy, [pietro.baldi@unina.it](mailto:pietro.baldi@unina.it).

Massimiliano Berti, Riccardo Montalto, SISSA, Via Bonomea 265, 34136, Trieste, Italy, [berti@sissa.it](mailto:berti@sissa.it), [riccardo.montalto@sissa.it](mailto:riccardo.montalto@sissa.it).